DIANION OF DIETHYL SUCCINATE: REACTIONS WITH ALKYLATING AGENTS AND CARBONYL COMPOUNDS

Manat POHMAKOTR,^{*} Vichai REUTRAKUL, Tipparat PHONGPRADIT, and Arayan CHANSRI Department of Chemistry, Faculty of Science, Mahidol University, Rama VI Road, Bangkok 4, Thailand

Dianion of diethyl succinate reacted readily with alkylating agents and carbonyl compounds to give α , β -dialkylsubstituted diethyl succinate and paraconic esters respectively. The conversion of some paraconic esters into 4-carboethoxycyclopentenones was also demonstrated.

Dianions of numerous classes of organic compounds have been reported and these species played important roles in organic syntheses due to their high regioselectivity with electrophiles.¹ Recently, P.J. Garatt and co-workers indicated that vicinal dianions of α,β -disubstituted succinic esters could undergo addition reactions with various electrophiles.² In the course of our exploration of synthetic pathways to lignan systems of types <u>1</u> and <u>2</u>, which are classes of natural products exhibiting interesting biological activities,³ we needed a synthetic equivalent containing four-carbon unit of type <u>3</u> which possesses nucleophilic sites at C-2 and C-3, as shown in Scheme I.

Scheme I



Recent report by M.W. Rathke⁴ on the alkylation of the dianion <u>4</u> with alkyl halides prompted us to report our results in this area. The reaction of one equivalent of <u>4</u> with **two** equivalents of halides, aliphatic aldehydes and ketones at -78° C gave good yields of alkylated products <u>5</u> and paraconic esters <u>6</u> respectively. The use of excess aldehydes or ketones in the reaction gave no diadduct and only the paraconic esters $\underline{6}$ could be isolated. The sequential treatment of the dianion $\underline{4}$ with cyclohexanone (or acetone) (1 equiv.) and methyl iodide (or allyl bromide) (1 equiv.) gave only trace amount of the expected diadduct, the major product being the paraconic esters.



The reaction of <u>4</u> with aryl aldehydes under standard conditions gave trace amount of the expected paraconic esters of the type <u>6</u>. However, upon addition of equivalent of 1M THF solution of $2nCl_2$ at -78° to the reaction mixture resulted in moderate yields of the expected paraconic esters. The results are summarized in <u>Table I</u>.

Alkylating agents or Carbonyl compounds	Products	Yields (%) ^a	Diastereomeric ratio ^b
СНЗІ	$\underline{5a}$, R^1 = CH ₃	85	_c
CH ₂ =CHCH ₂ Br	$\underline{5b}$, $R^1 = CH_2 = CHCH_2$	80	_c
C ₆ H ₅ CH ₂ Br	$5c$, $R^1 = C_6 H_5 C H_2$	84	_c
снзсно	<u>6a</u> , $R^2 = H$; $R^3 = CH_3$	61	1:1.3 ^d
<u>n</u> -C ₃ H ₇ CHO	<u>6b</u> , $R^2 = H$; $R^3 = \underline{n} - C_3 H_7$	59	1:2 ^d
<u>n</u> -C ₆ H ₁₃ CH0	<u>6e</u> , $R^2 = H$; $R^3 = \underline{n} - C_6 H_{13}$	58	1:1 ^d
снзсоснз	<u>6d</u> , $R^2 = R^3 = CH_3$	86	-
сн _з сосн ₂ сн ₃	$\underline{6e}$, R^2 = CH_3 ; R^3 = CH_3CH_2	83	-
Cyclopentanone	<u>6f</u> , $R^2 - R^3 = -(CH_2)_4$ -	81	-

Table I (cont.)

Alkylating agents or Carbonyl Compounds	Products	Yields (%) ^a	Diastereomeric ratio ^b
Cyclohexanone	<u>6g</u> , $R^2 - R^3 = -(CH_2)_5$ -	84	-
Cyclooctanone	<u>6h</u> , $R^2 - R^3 = -(CH_2)_7 -$	59	-
<u>р</u> -сн ₃ ос ₆ н ₄ сно	<u>6i</u> , $R^2 = H$; $R^3 = p-CH_3OC_6H_4$	70	1:1.5 ^e
<u>т</u> -сн ₃ ос ₆ н ₄ сно	<u>6j</u> , $R^2 = H$; $R^3 = \underline{m} - CH_3 OC_6 H_4$	43	1:1.5 ^e

a. Isolated yields by preparative layer chromatography (PLC).

b. The structure of diastereomers were determined by ${}^{1}\text{H-NMR}^{8}$.

c. Ratio could not be determined either by PLC or 1 H-NMR.

d. Ratio was determined by PLC separation.

e. Ratio was determined by ¹H-NMR.

Our results demonstrated the synthetic potential of the dianion $\underline{4}$ for the preparations of disubstituted succinic esters and paraconic esters. These types of paraconic esters are of special interest due to their versatile utility as intermediates in the synthesis of natural products.^{5,7} Thus, the lactones <u>6e-h</u> had been converted into cyclopentenone derivatives <u>7a-d</u> by treating with phosphorus pentoxide/methanesulfonic acid mixture.⁶ The results are summarized in <u>Scheme II</u>. Compound <u>7a</u> was an important synthetic intermediate for the preparation of (±)-Methylenomycin A, an antibiotic from the culture filtrate of a streptomycete, which had previously been reported.⁷

Scheme II



Further exploratory investigations including the syntheses of the lignans types $\underline{1}$ and $\underline{2}$ are now in progress.

REFERENCES

- See for examples: E.M. Kaiser, J.D. Petty, and P.L.A. Knutson, Synthesis, 509 (1981);
 D. Seebach and M. Pohmakotr, Tetrahedron, <u>37</u>, 4147 (1981); D. Seebach, M. Pohmakotr,
 Ch. Schregenberger, B. Weidmann, R.S. Mali, and S. Pohmakotr, Helv.Chim.Acta, <u>65</u> (1982);
 R.S. Mali, M. Pohmakotr, B. Weidmann, and D. Seebach, Liebigs Ann. Chem. 2272 (1981).
- K.G. Bilyard, P.J. Garatt, and R. Zahler, Synthesis, 389 (1980) and references cited therein;
 C.W. Doecke and P.J. Garatt, J.C.S. Chem.Comm., 873 (1981); K.G. Bilyard and P.J. Garratt,
 Tetrahedron Lett., <u>22</u>, 1755 (1981).
- P. Brownbridge and T.H. Chan, Tetrahedron Lett., <u>21</u>, 3427 (1980); A. Pelter, P. Satyanarayana, and R.S. Ward, ibid., <u>22</u>, 1549 (1981) and references cited therein.
- 4. N.R. Long and M.W. Rathke, Synth. Comm., <u>11</u>, 687 (1981).
- 5. S. Kano, S. Shiyuya, and T. Ebata, Heterocycles, 14, 661 (1980) and references cited therein.
- 6. P.E. Eaton, G.R. Carlson, and J.T. Lee, J.Org.Chem., <u>38</u>, 4071 (1973).
- J. Jernow, W. Tautz, P. Rosen, and J.F. Blout, J. Org. Chem., <u>44</u>, 4210 (1979); Y. Takahashi,
 K. Isobe, H. Hagiwara, H. Kosugi, and H. Uda, J.C.S. Chem.Comm., 714 (1981).
- 8. L.M. Jackman and S. Sternhell, Application of Nuclear Magnetic Resonance Spectroscopy in Organic Chemistry, Pergamon Press, London, 1969.

(Received March 13, 1982)