

131. 3-Lithiopyrroles by Halogen-Metal Interchange of 3-Bromo-1-(triisopropylsilyl)pyrroles. Synthesis of Verrucarin E and Other 3-Substituted Pyrroles¹⁾

Preliminary Communication

by Joseph M. Muchowski* and Reto Naef²⁾

Syntex Research, Institute of Organic Chemistry, Palo Alto, California 94304, U.S.A.

(15.III.84)

Summary

3-Lithio-1-(trimethylsilyl)pyrrole (**7**, *Scheme 2*), obtained by halogen-metal interchange from the 3-bromo compound **2**, reacted with various electrophilic reagents to provide products, which on fluoride ion desilylation, gave 3-substituted pyrroles in good overall yields. One such pyrrole **13** (*Scheme 3*), was converted into 2-formyl-3-octadecylpyrrole (**14**), reputed to be a metabolite of the marine sponge *Oscarella lobularis*.

3,4-Dibromo-1-(triisopropylsilyl)pyrrole (**5**) was efficiently transformed, by a process involving two consecutive bromine-lithium exchange reactions (*Scheme 4*), into the antibiotic verrucarin E **17**.

3-Substituted pyrroles have been synthesized by several processes, of variable efficiency which include acid-induced isomerization of the easily accessible α -isomers [1], β -substitution of *N*-(benzenesulfonyl)pyrrole with some electrophiles [2], and the electrophilic substitution of pyrroles bearing a removable deactivating 2-substituent [3] or a sterically demanding *N*-substituent [4] [5]. With regard to the last mentioned process, it was recently [5] shown that *N*-(triisopropylsilyl)pyrrole **1** undergoes highly selective kinetic electrophilic substitution at C(3) and that the silyl moiety is readily removed from the products obtained thereby with tetrabutylammonium fluoride [5] [6]³⁾. Among the examples reported was the reaction with *N*-bromosuccinimide which gave 3-

¹⁾ Contribution No. 667 from the Syntex Institute of Organic Chemistry.

²⁾ Syntex Research Post-Doctoral Fellow, 1983–1984.


³⁾ The 'strong steric screening' properties of the triisopropylsilyl group were first described in [6]. In [5], no acknowledgement of this important observation was made nor was due credit given for the development of the tetrabutylammonium fluoride scission of trialkylsilyl protecting groups [7]. We apologize to Prof. Corey and co-workers for this oversight. For other instances where the steric properties of this protecting group have profoundly affected the positional selectivity of a reaction, see [8].

bromo-1-(triisopropylsilyl)pyrrole **2** as the principal product (85:15 of β - vs. α -substitution at -80° to -20°). This particular result was of obvious importance because it was expected that **2** itself could serve as the source of a wide variety of 3-substituted pyrroles *via* the corresponding lithio compound. The results described herein confirm this expectation and demonstrate that di- and tribrominated derivatives of *N*-(triisopropylsilyl)pyrrole (**1**) also have a certain degree of synthetic utility.

Bromination of **1** with 1, 2, or 3 molar equiv. of *N*-bromosuccinimide at -78° in THF [9], gave a 97:3 mixture of **2** and the α -bromo isomer **3**, a 1:1 mixture of the 2,3- and 3,4-dibromopyrroles **4** and **5**, or the tribromopyrrole **6** (m.p. 46°) exclusively (*Scheme 1*)⁴. The 3,4-dibromo compound **5**, a solid (m.p. 77°), was easily separated from **4**, an oil, by low-temperature crystallization from pentane. Furthermore, the tribromopyrrole **6** was convertible, in high yield (see *below*) into **5**, thus making the latter compound readily accessible.

The formation of a substantial proportion of **4** in the above mixture is doubly noteworthy. Firstly, it demonstrates that the steric effect of the triisopropylsilyl moiety is not sufficient to entirely overcome the electronic directing effect (to C(2) [10]) of a bromo group at C(3). Secondly, it contrasts with the reported [4c] predominant formation of the 3,4-dibromo compound from *N*-tritylpyrrole and bromine.


Scheme 1



⁴) All new compounds were characterized by IR, 1H -NMR and mass spectra and had satisfactory elemental analyses.

The lithiation of 3-bromo-1-(triisopropylsilyl)pyrrole (**2**) was effected⁵⁾ by halogen-metal interchange with 1 equiv. of BuLi or 2 equiv. of *t*-BuLi in THF at -78° (0.5 h, *Scheme 2*). The 3-lithiopyrrole derivative **7** reacted with a broad spectrum of electrophilic reagents (see *Table*) which included alkyl halides (benzyl bromide gave bibenzyl, secondary alkyl halides were dehydrohalogenated), aldehydes, cyclohexanone, DMF, carbon dioxide, *etc.*, and in some instances, the products thus obtained were desilylated with tetrabutylammonium fluoride (THF, room temp., 5 min) to the 3-substituted pyrroles **9**. Therefore **7** is an effective, formal equivalent of 3-lithiopyrrole.

Table. 3-Substituted Pyrroles from 3-Lithio-1-(triisopropylsilyl)pyrrole



RLi (equiv.)	Electrophile	E in product	Yield [%]	M.p. [°C]
<i>t</i> -BuLi (2)	CH ₃ I	CH ₃	92	oil
<i>t</i> -BuLi (2)	C ₁₈ H ₃₇ I	C ₁₈ H ₃₇	88	glass
<i>t</i> -BuLi (2)	(CH ₃) ₃ SiCl	(CH ₃) ₃ Si	87	oil
BuLi (1)	CO ₂	COOH ^{a)}	88	162
<i>t</i> -BuLi (2)	HCHO gas	CH ₂ OH	73	37
BuLi (1)	C ₆ H ₅ CHO	CHOHC ₆ H ₅	48	132
<i>t</i> -BuLi (2)	Cyclohexanone	 ^{b)}	69	oil
BuLi (1)	DMF	CHO ^{c)}	82	oil
<i>t</i> -BuLi (2)	CH ₃ CONCH ₃ OCH ₃	COCH ₃ ^{d)}	61	69

^{a)} Desilylated (88%) to pyrrole-3-carboxylic acid, m.p. 149° ([19]: m.p. 150–150.5°).

^{b)} Dehydration effected by heating crude product at 150° for 45 min.

^{c)} Desilylated (86%) to pyrrole-3-carbaldehyde, m.p. 63° ([20]: m.p. 63–63.5°).

^{d)} Desilylated (84%) to 3-acetylpyrrole, m.p. 111–112° ([20]: m.p. 114–115°).


The mono-lithiation of the 3,4-dibromo and 2,3,4-tribromo compounds **5** and **6** was accomplished under conditions analogous to those used for the formation of **7**. The α -lithiated species **10** (*Scheme 2*), which reacted with the usual range of electrophiles⁶⁾, was formed essentially exclusively (> 99%) as deduced by protonolysis to 3,4-dibromo-1-(triisopropylsilyl)pyrrole (**5**) (85% yield).

The methodology described above was then utilized to synthesize a purported metabolite **14** of the marine sponge *Oscarella lobularis* [14] as well as the antibiotic verrucaric acid (**17**) [15]. Thus, for the former substance, 3-octadecyl-1-(triisopropylsilyl)pyrrole (**12**) (*Scheme 3*), obtained in nearly 90% yield from the lithio compound **7** and octadecyl iodide, was desilylated (76%) and the 3-substituted pyrrole **13** (m.p. 53°), obtained in this way, was formylated under *Vilsmeier-Haack* [16] conditions. A

⁵⁾ *N*-Benzyl-3-lithiopyrrole has been obtained from the bromo compound and metallic lithium [11] and 4,4-dimethyl-2-(*N*-methylpyrrol-2-yl)oxazoline can be selectively lithiated at C(3) [12]. The usefulness of these reagents is minimal because removal of the *N*-substituent is not possible. *N*-(Trimethylsilyl)pyrrole is metallated to a considerable extent at C(3) with excess *t*-BuLi in pentane [13].


⁶⁾ A detailed description of this study will be reported in the full paper.

Scheme 2




crystalline (m.p. $72-75^\circ$) 3:1 mixture (as yet not separable) of the supposed natural product **14** and the isomeric aldehyde **15** was isolated in 85% yield⁷⁾. To synthesize verrucaric acid **17**, the 3,4-dibromo compound **5** reacted sequentially, in one pot (Scheme 4), with $BuLi$ (1 equiv., $THF/-78^\circ$), N -methoxy- N -methylacetamide [**17**], t - $BuLi$ (2 equiv., $THF/-78^\circ$) and dry, gaseous formaldehyde, to produce the 3-acetyl-4-hydroxymethylpyrrole derivative **16** (m.p. 91°) in 58% overall yield, after chromatographic purification on silica gel. Desilylation of **16** gave crystalline verrucaric acid **17**, m.p. 90° ([15a]: m.p. $90.5-91^\circ$), the NMR spectral characteristic of which were identical to those published [15a]. This synthesis of **17** is considerably more efficient (44% yield overall) than two of those previously reported [15] [18] and compares not too unfavorably with the elegant and efficient three-step synthesis of Gossauer & Suhl [21].

Scheme 3



⁷⁾ **14**: 1H -NMR (300 MHz, CCl_4): 0.88 (t , $J = 6.5$, 3 H, CH_3CH_2); 1.25 (m , 30 H); 1.63 (m , 2 H, CH_2); 2.75 (m , $J = 7.6$, 2 H, CCH_2); 6.05 (d , $J = 2.5$, 1 H, $H-C(4)$); 7.03 (dd , $J(4,5) = 2.5$, $J(5,CHO) = 1.1$, 1 H, $H-C(5)$); 9.61 (d , $J = 1.1$, 1 H, CHO); 10.95 (br. s , 1 H, NH). The 1H -NMR spectrum of **14** is similar to but not identical with that reported in [14a] [0.9 (t , $J = 6$, 3 H); 1.26 (m , 2 H); 2.70 (t , $J = 6$, 2 H); 5.94 (d , $J = 2.4$, 1 H); 6.76 (d , $J = 2.4$, 1 H); 9.22 (s , 1 H); 11.0 (br. s , 1 H, NH)]. The latter parameters are, however, nearly identical with those in [14b] for the 2,5-disubstituted isomer, except that these authors observed a larger coupling constant ($J = 3.7$) for the ring protons and two-proton m at 1.68.

Scheme 4



The authors wish to acknowledge the spirited discussion of some aspects of this work with Dr. R. Clark and J. Berger, and the extra effort, expended on our behalf, by Dr. M. Maddox with regard to several of the NMR spectra.

REFERENCES

- [1] J. DeSales, R. Greenhouse & J. M. Muchowski, *J. Org. Chem.* **47**, 3668 (1982) and references therein.
- [2] R. X. Xu, H. J. Anderson, N. J. Gogan, C. E. Loader & R. McDonald, *Tetrahedron Lett.* **22**, 4899 (1981); J. Rokach, P. Hamel, M. Kakushima & G. M. Smith, *ibid.* **22**, 4901 (1981).
- [3] C. E. Loader & H. J. Anderson, *Can. J. Chem.* **59**, 2673 (1981) and references therein.
- [4] a) C. F. Candy, R. A. Jones & P. H. Wright, *J. Chem. Soc. C* **1970**, 2563; b) D. J. Chadwick, G. D. Meakins & C. A. Rhodes, *J. Chem. Res. (S)* **42**, (M) 0878 (1983); c) D. J. Chadwick & S. T. Hodgson, *J. Chem. Soc., Perkin Trans. 1* **1983**, 93.
- [5] J. M. Muchowski & D. R. Solas, *Tetrahedron Lett.* **24**, 3455 (1983).
- [6] E. J. Corey, H. Cho, C. Rücker & D. H. Hua, *Tetrahedron Lett.* **22**, 3455 (1981).
- [7] E. J. Corey & B. B. Snider, *J. Am. Chem. Soc.* **94**, 2549 (1972); E. J. Corey & A. Venkateswarly, *ibid.* **94**, 6190 (1972).
- [8] G. Nechtatal & D. A. Widdowson, *J. Chem. Soc., Chem. Commun.* **1982**, 467; N. F. Mastens & D. A. Widdowson, *ibid.* **1983**, 955.
- [9] H. M. Gilow & D. E. Burton, *J. Org. Chem.* **46**, 221 (1981); E. Aiello, G. Dattolo, G. Cirrincione, A. M. Almerico & I. D'Asidia, *J. Heterocycl. Chem.* **19**, 977 (1982).
- [10] R. A. Jones & G. P. Bean, *The Chemistry of Pyrroles*, Academic Press, London, 1977, pp. 151, 153.
- [11] H. J. Anderson & S. J. Griffiths, *Can. J. Chem.* **45**, 2227 (1967).
- [12] D. J. Chadwick, M. V. McKnight & R. Ngochindo, *J. Chem. Soc., Perkin Trans. 1* **1982**, 1343.
- [13] D. J. Chadwick & S. T. Hodgson, *J. Chem. Soc., Perkin Trans. 1* **1982**, 1833.
- [14] a) G. Cimino, S. de Stefano & L. Minale, *Experientia* **31**, 1387 (1975); b) see also, D. B. Stierle & J. D. Faulkner, *J. Org. Chem.* **45**, 4980 (1980).
- [15] a) E. Fetz & Ch. Tamm, *Helv. Chim. Acta* **49**, 349 (1966); b) P. Pfäffli & Ch. Tamm, *ibid.* **52**, 1911 (1969).
- [16] A. Vilsmeier & Haack, *Ber. Dtsch. Chem. Ges.* **60**, 119 (1927); M. R. de Maheas, *Bull. Soc. Chim. Fr.* **1962**, 1989.
- [17] M. S. Nahm & S. M. Weinreb, *Tetrahedron Lett.* **22**, 3815 (1981).
- [18] J. K. Groves, N. E. Cundasawmy & H. J. Anderson, *Can. J. Chem.* **51**, 1089 (1973).
- [19] H. Rapoport & C. D. Wilson, *J. Org. Chem.* **26**, 1102 (1961).
- [20] C. E. Loader & H. J. Anderson, *Tetrahedron* **25**, 3979 (1969).
- [21] A. Gossauer & K. Suhl, *Helv. Chim. Acta* **59**, 1698 (1976).