

Communications

Kinetic Resolution of Planar-Chiral Ferrocenes by Molybdenum-Catalyzed Enantioselective Metathesis

Masamichi Ogasawara,^{*,†} Susumu Watanabe,[†] Liyan Fan,[†] Kiyohiko Nakajima,[‡] and Tamotsu Takahashi^{*,†}




Catalysis Research Center and Graduate School of Pharmaceutical Sciences, Hokkaido University, and SORST, Japan Science and Technology Agency (JST), Kita-ku, Sapporo 001-0021, Japan, and Department of Chemistry, Aichi University of Education, Igaya, Kariya, Aichi, 448-8542, Japan

Received September 11, 2006

Summary: Kinetic resolution of planar-chiral 1,1'-diallylferrocene derivatives was realized by molybdenum-catalyzed asymmetric ring-closing metathesis with excellent enantioselectivity. This is the first example of highly enantioselective metal-catalyzed methods of preparing optically active planar-chiral metallocenes.

Planar-chiral ferrocenes are important chiral scaffolds in organic and organometallic chemistry.¹ Although such ferrocenes have been used in a variety of asymmetric reactions as ligands² or catalysts,³ their preparation in optically active form is still a challenging problem. Most nonracemic planar-chiral ferrocenes were obtained either by diastereoselective metalation utilizing chiral *ortho*-directing groups⁴ or by optical resolution of racemic compounds. Enantioselective lithiation of prochiral ferrocenes by a stoichiometric chiral base has also shown fair success.⁵ On the other hand, examples of *catalytic* asymmetric synthesis of planar-chiral ferrocene derivatives are extremely

Scheme 1. ARCM Kinetic Resolution of Planar-Chiral Ferrocenes and Metathesis Catalysts.



* To whom correspondence should be addressed. E-mail: ogasawara@cat.hokudai.ac.jp.

[†] Hokkaido University.

[‡] Aichi University of Education.

(1) For reviews of planar-chiral ferrocenes, see: (a) Wagner, G.; Herrmann, R. In *Ferrocenes*; Togni, A., Hayashi, T., Eds.; VCH: Weinheim, 1995; Chapter 4, p 173. (b) Togni, A. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1475. (c) Richards, C. J.; Locke, A. J. *Tetrahedron: Asymmetry* **1998**, *9*, 2377. See, also: (d) Halterman, R. L. *Chem. Rev.* **1992**, *92*, 965.

(2) (a) Hayashi, T. In *Ferrocenes*; Togni, A., Hayashi, T., Eds.; VCH: Weinheim, 1995; Chapter 2, p 105. (b) Togni, A. In *Metallocenes*; Togni, A., Halterman, R. L., Eds.; Wiley-VCH: Weinheim, 1998; Vol. 2, Chapter 11, p 685. (c) Colacot, T. J. *Chem. Rev.* **2003**, *103*, 3101. (d) Barbaro, P.; Bianchini, C.; Giambastiani, G.; Parisel, S. L. *Coord. Chem. Rev.* **2004**, *248*, 2131.

(3) (a) Butsugan, Y.; Araki, S.; Watanabe, M. In *Ferrocenes*; Togni, A., Hayashi, T., Eds.; VCH: Weinheim, 1995; Chapter 3, p 143. (b) Fu, G. C. *Acc. Chem. Res.* **2000**, *33*, 412. (c) Fu, G. C. *Acc. Chem. Res.* **2004**, *37*, 542.

(4) (a) Marquarding, D.; Klusacek, H.; Gokel, G.; Hoffmann, P.; Ugi, I. *J. Am. Chem. Soc.* **1970**, *92*, 5389. (b) Riant, O.; Samuel, O.; Kagan, H. B. *J. Am. Chem. Soc.* **1993**, *115*, 5835. (c) Rebiere, F.; Riant, O.; Ricard, L.; Kagan, H. B. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 568. (d) Richards, C. J.; Damalidis, T.; Hibbs, D. E.; Hursthouse, M. B. *Synlett* **1995**, *74*. (e) Sammakia, T.; Latham, H. A.; Schaad, D. R. *J. Org. Chem.* **1995**, *60*, 10. (f) Nishibayashi, Y.; Arikawa, Y.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1996**, *61*, 1172. (g) Enders, D.; Peters, R.; Lochtman, R.; Raabe, G. *Angew. Chem., Int. Ed.* **1999**, *38*, 2421.

(5) (a) Tsukazaki, M.; Tinkl, M.; Roglans, A.; Chapell, B. J.; Taylor, N. J.; Snieckus, V. *J. Am. Chem. Soc.* **1996**, *118*, 685. (b) Nishibayashi, Y.; Arikawa, Y.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1996**, *61*, 1172. (c) Laufer, R. S.; Veith, U.; Taylor, N. J.; Snieckus, V. *Org. Lett.* **2000**, *2*, 629. (d) Fukuda, T.; Imazato, K.; Iwao, M. *Tetrahedron Lett.* **2003**, *44*, 7503. (e) Metallinos, C.; Szillat, H.; Taylor, N. J.; Snieckus, V. *Adv. Synth. Catal.* **2003**, *345*, 370.

rare.^{6–8} Apart from several reports on enzymatic resolution of racemic ferrocenes,⁶ to the best of our knowledge, only two examples of catalytic induction of ferrocenyl planar chirality was reported so far with modest enantioselectivity.^{7,8}

Richards⁹ and we¹⁰ recently reported preparation of [4]-ferrocenophanes by Ru- or Mo-catalyzed metathesis reaction of 1,1'-diallylferrocenes.¹¹ Here, we report kinetic resolution of planar-chiral 1,1'-diallylferrocene derivatives by Mo-catalyzed asymmetric ring-closing metathesis (ARCM).¹² The reaction

(6) For representative examples, see: (a) Izumi, T.; Hino, T. *J. Chem. Technol. Biotechnol.* **1992**, *55*, 325. (b) Lambusta, D.; Nicolosi, G.; Patti, A.; Piattelli, M. *Tetrahedron Lett.* **1996**, *37*, 127. (c) Patti, A.; Lambusta, D.; Piattelli, M.; Nicolosi, G. *Tetrahedron: Asymmetry* **1998**, *9*, 3073.

(7) Siegel, S.; Schmalz, H.-G. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2456.

(8) Enantioselective lithiation of prochiral ferrocenes using alkylolithium in conjunction with substoichiometric quantities of chiral diamines was reported recently; see: Genet, C.; Canipa, S. J.; O'Brien, P.; Taylor, S. J. *Am. Chem. Soc.* **2006**, *128*, 9337.

(9) Locke, A. J.; Jones, C.; Richards, C. J. *J. Organomet. Chem.* **2001**, *637–639*, 669.

(10) (a) Ogasawara, M.; Nagano, T.; Hayashi, T. *J. Am. Chem. Soc.* **2002**, *124*, 9068; *J. Am. Chem. Soc.* **2002**, *124*, 12626. (b) Ogasawara, M.; Nagano, T.; Hayashi, T. *Organometallics* **2003**, *22*, 1174.

Table 1. Molybdenum-Catalyzed Asymmetric RCM Kinetic Resolution of Planar-Chiral Ferrocenes^a

entry	substrate 1 (mol/L) ^b	Mo* cat. (%)	conditions	yields (%) ^c 2/1/3	% ee 2/1 ^d	<i>k</i> _{rel} ^e
1	1a (0.1)	5	23 °C, 15 min	73/26/trace	<1/2	1.03
2	1b (0.1)	5	23 °C, 1 h	44/49/trace	12/9	1.4
3	1c (0.01)	10	80 °C, 24 h	26/53/16	66/29	6.4
4	1d (0.01)	5	70 °C, 24 h	0/54/38	−/7	
5	1e (0.1)	5	23 °C, 4 h	23/30/47	>99.5/78	>500
6	1e (0.005)	10	50 °C, 24 h	46/47/3	96/95	183
7	1f (0.005)	10	50 °C, 24 h	43/41/12	82/83	26
8	1g (0.005)	10	50 °C, 24 h	45/52/trace	97/81	165


^a The reaction was carried out in benzene in the presence of the molybdenum catalyst (*R*)-Mo*. ^bInitial concentration of the substrate. ^cIsolated yields of the products (**2** and **3**) and the recovered substrate (**1**). ^dEnantiomeric excess of the recovered **1** was determined after converting them into **2** by a reaction with the second-generation Grubbs' catalyst. ^eCalculated based on a first-order equation (ref 14).

proceeds very efficiently to afford planar-chiral ferrocenes in high yield with excellent enantiomeric enrichment.

The ferrocene substrates for this study (*rac*-**1**, see Scheme 1) possess a trisubstituted η^5 -(C₅H₂-1-allyl-2,4-R₂) ligand, which constructs a planar-chiral environment in the ferrocenes, and a monosubstituted η^5 -cyclopentadienyl ligand with an allylic side chain. A readily available chiral molybdenum species (*R*)-Mo* (Scheme 1)¹³ was chosen as an asymmetric metathesis catalyst.

Enantioselectivity in the ARCM kinetic resolution was strongly dependent on the structure of the allylic group in the monosubstituted cyclopentadienyl moiety. Treatment of **1a** with the Mo* catalyst (5 mol %) for 15 min in benzene at 23 °C yielded the corresponding ferrocenophane **2a** in 73%, and the unreacted **1a** was recovered in 26%. To our disappointment, however, the selectivity of the reaction was very low: **2a** was nearly racemic (<1% ee) and the recovered **1a** was only 2% ee (Table 1, entry 1). The RCM reaction of the ferrocene **1b**, which was with η^5 -(C₅H₄-crotyl), showed slightly better enantioselectivity and the ferrocenophane **2a** of 12% ee was obtained in 44% yield (entry 2). Introduction of an η^5 -(C₅H₄-cinnamyl) ligand in **1c** further improved enantioselectivity of the ARCM, although diluted conditions were required to suppress formation of metathesized homodimer **3** (as a mixture of stereoisomers). A reaction of **1c** at 80 °C (initial concentration of **1c** = 0.01 mol/L) gave **2a** (66% ee, 26% yield) and recovered **1c** (29% ee, 53%). The *k*_{rel} value ([rate of fast-reacting enantiomer]/[rate of slow-reacting enantiomer]) for this reaction was estimated to be 6.4¹⁴ (entry 3). A reaction of **1d**, which was with η^5 -C₅H₄CH₂CH=CMe₂, produced the dimer **3d** in 38% as a sole metathesis product and **2a** was not detected (entry 4). The recovered **1d** was only 7% ee; that is, kinetic resolution of **1d** at the dimerization was not very efficient.

It was found that the enantioselectivity was dramatically improved by introducing a η^5 -(C₅H₄-methallyl) moiety in the ferrocene substrates. The RCM reaction of **1e** at 23 °C (initial concentration of **1e** = 0.1 mol/L) proceeded with excellent enantioselectivity. The bridged ferrocene **2e** was obtained in

**Figure 1.** ORTEP drawing of (+)-(*R*)-**2e** with 40% thermal ellipsoids. All hydrogen atoms are omitted for clarity.

nearly enantiomerically pure form (>99.5% ee) in 23% yield, and unreacted **1e** of 78% ee was recovered in 30% yield. The *k*_{rel} for this reaction was >500¹⁴ (entry 5). Unfortunately, however, 47% of the substrate was converted to dimer **3e**. The dimer **3e**, which could be converted to the ferrocenophane **2e** by treatment with the second-generation Grubbs catalyst, was only 7% ee; that is, the dimer formation occurred with negligible enantioselectivity. As before, dimerization was minimized in the reaction under diluted conditions. A reaction of **1e** with initial concentration of 0.005 mol/L affords **2e** of 96% ee in 47% yield, and unreacted **1e** of 95% ee was recovered in 46% (entry 6). Although somewhat higher temperature (50 °C) was required to gain a reasonable reaction rate under the high-dilution conditions, enantioselectivity of the reaction was still excellent (*k*_{rel} = 183) and nearly perfect kinetic resolution of *rac*-**1e** was achieved. As shown in entries 7 and 8, similarly high levels of enantioselectivity and reaction efficiency were observed for kinetic resolution of **1f** (*k*_{rel} = 26) and **1g** (*k*_{rel} = 165) also under the optimized conditions.

Single-crystal X-ray crystallography revealed that absolute configuration of the ARCM product **2e** (Table 1, entry 5; >99.5% ee), which was dextrorotatory ([α]_D²⁰ +22 (c 0.50 in CHCl₃)), was *R*, as shown in Figure 1. The configurations of other ARCM products were assigned by analogy.

For highly enantioselective kinetic resolution of *rac*-**1**, an allyl group in a trisubstituted cyclopentadienyl needs to be more reactive than an allylic substituent in a monosubstituted cyclopentadienyl. When the (*R*)-Mo* species approaches *rac*-**1a**, initial metathesis occurs with the less crowded C₅H₄-allyl moiety preferentially to form a diastereomeric mixture of (*R*_a,*R*_p)- and (*R*_a,*S*_p)-**4** ("R_a" represents absolute configuration of the axially chiral biaryl moiety in Mo*) and formation of **5** is unfavorable (Scheme 2, top). Whereas the allyl group in C₅H₄-allyl is remote from the planar-chiral η^5 -C₅H₂Bu₂(allyl) moiety, initial me-


(11) For examples of metathesis reactions of metallocene substrates, see: (a) Hürlerlände, D.; Kleigrew, N.; Kehr, G.; Erker, G.; Fröhlich, R. *Eur. J. Inorg. Chem.* **2002**, 2633. (b) Sierra, J. C.; Hürlerlände, D.; Hill, M.; Kehr, G.; Erker, G.; Fröhlich, R. *Chem. Eur. J.* **2003**, 9, 3618. (c) Kuwabara, J.; Takeuchi, D.; Osakada, K. *Organometallics* **2005**, 24, 2705. (d) Bauer, E. B.; Gladysz, J. A. In *Handbook of Metathesis*; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003; Vol. 2, Chapter 2.11, p 403.

(12) For reviews of asymmetric olefin metathesis, see: (a) Hoveyda, A. H.; Schrock, R. R. *Chem. Eur. J.* **2001**, 7, 945. (b) Schrock, R. R.; Hoveyda, A. H. *Angew. Chem., Int. Ed.* **2003**, 42, 4592. (c) Hoveyda, A. H. In *Handbook of Metathesis*; Grubbs, R. H., Ed.; Wiley-VCH: Weinheim, 2003; Vol. 2, Chapter 2.3, p 128.

(13) Aeilts, S. L.; Cefalo, D. R.; Bonitatebus, P. J., Jr.; Houser, J. H.; Hoveyda, A. H.; Schrock, R. R. *Angew. Chem., Int. Ed.* **2001**, 40, 1452.

(14) (a) Kagan, H. B.; Fiaud, J. C. *Top. Stereochem.* **1998**, 18, 249. (b) Vedejs, E.; Jure, M. *Angew. Chem., Int. Ed.* **2005**, 44, 3974.

Scheme 2. Plausible Stereochemical Pathways of the Mo-Catalyzed ARCM Kinetic Resolution of 1



tathesis with (R) -Mo* takes place with negligible stereoselectivity. Following the RCM step in **4** is an intramolecular process, and thus both (R_a,R_p) - and (R_a,S_p) -**4** can be smoothly transformed into the corresponding ferrocenophanes to give **2a** in nearly

racemic form. On the other hand, with a less reactive methallyl group in a monosubstituted Cp in place of the allyl group, the reaction between (R) -Mo* and *rac*-**1e** takes place at the allyl group in the planar-chiral trisubstituted Cp preferentially and formation of **7** is minor. While (R_a,R_p) -**6** is transformed into (R) -**2e** smoothly via RCM, the epimeric (R_a,S_p) -**6** is forced to take an unfavorable conformation due to the steric repulsion between a t Bu group in η^5 -C₅H₂ t Bu₂(allyl) and a t Bu group in the chiral biaryl ligand in (R) -Mo* to liberate (S) -**1e** (Scheme 2, bottom).

In summary, we have developed an effective method for kinetic resolution of racemic planar-chiral ferrocene derivatives via molybdenum-catalyzed asymmetric ring-closing metathesis. This is the first example of highly enantioselective metal-catalyzed induction of planar chirality in metallocene substrates.

Acknowledgment. This work was partly supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan, and the Nagase Science and Technology Foundation (to M.O.).

Supporting Information Available: Detailed experimental procedures, compound characterization data, and crystallographic data (CIF file). This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM0608298