

Deprotonated 2-Benzylpyridine as a Bridging Ligand. Synthesis and Structure of the Binuclear Palladium(II) Complex $trans(N,N)-[Pd(\mu\text{-pyCHPh-N,CH})(\gamma\text{-picoline})Cl]_2 \cdot CH_2Cl_2$

ALLAN J. CANTY, NIGEL J. MINCHIN


Chemistry Department, University of Tasmania, Hobart, Tas. 7001, Australia

LUTZ M. ENGELHARDT and ALLAN H. WHITE

Department of Physical and Inorganic Chemistry, University of Western Australia, Nedlands, W.A. 6009, Australia

Received March 29, 1985

The initial report [1] that 2-benzylpyridine is readily orthometallated by palladium(II) acetate to give $[Pd(pyCH_2(C_6H_4)_N, C^1)(\mu\text{-O}_2\text{CMe})]_2$, with deprotonated 2-benzylpyridine bonded to palladium(II) as in (1), has led to extensive development of the organopalladium(II) chemistry of orthometallated 2-benzylpyridine [1–3].



As part of a study of the chemistry of anions of 2-pyridylmethanes [4], we have obtained a complex containing the moiety (2) on reaction of $PdCl_2(\gamma\text{-picoline})_2$ with 2-benzylpyridine deprotonated by *n*-butyllithium. As this represents a new binding mode for deprotonated 2-benzylpyridine, we report here the synthesis and structure of $trans(N,N)-[Pd(\mu\text{-pyCHPh-N,CH})(\gamma\text{-picoline})Cl]_2 \cdot CH_2Cl_2$ prior to investigation of its reactivity.

In a typical synthesis, lithiated 2-benzylpyridine in tetrahydrofuran was added to an equimolar quantity of $PdCl_2(\gamma\text{-picoline})_2$ in diethyl ether at -80°C under nitrogen. Benzene was added at ambient temperature, and on warming at 60°C for 1 h, hydrolysis gave an orange solid from the organic phase. Extraction with hexane gave a small quantity (*ca.* 4% yield) of $(pyCHPh)_2$, which has been obtained previously in 80% yield from reaction of lithiated 2-benzylpyridine with mercury(II) iodide [4]. Recrystallization of the remaining orange residue from dichloromethane/acetone gave orange-red crystals of the title compound in *ca.* 30% yield, containing traces of recrystallization solvents.




Fig. 1. ORTEP drawing of $trans(N,N)-[Pd(\mu\text{-pyCHPh-N,CH})(\gamma\text{-picoline})Cl]_2$ with selected atom numbering, bond distances (Å) and angles (deg) for the coordination plane of Pd [with corresponding values for Pd' in parentheses], and for the nitrogen and carbon atoms of $pyCHPh^-$ bonded to palladium: Pd–C7, 2.070(6) [2.079(5)]; Pd–N1, 2.046(4) [2.029(5)]; Pd–N2, 2.049(4) [2.041(5)]; Pd–Cl, 2.430(2) [2.446(2)]; C7'–Pd–N1, 91.7(2) [94.0(2)]; C7'–Pd–N2, 89.5(2) [88.6(2)]; N1–Pd–Cl, 91.1(1) [87.7(1)]; N2–Pd–Cl, 87.6(1) [89.9(1)]; C2–N1–Pd, 121.4(4) [124.0(4)]; C6–N1–Pd, 118.8(4) [119.7(5)]; C2'–C7'–Pd, 117.3(4) [112.9(4)]; C8'–C7'–Pd, 106.3(4) [112.1(4)]; C2–C7–C8, 115.5(4) [116.6(5)].

A single crystal X-ray study allowed unambiguous stereochemical assignment as the title compound (Fig. 1). Crystal data for $C_{37}H_{36}N_4Cl_2Pd_2$: monoclinic, space group $P2_1/c$ with $a = 13.967(2)$, $b = 13.996(3)$, $c = 18.886(2)$ Å; $\beta = 98.74(1)^\circ$; $V = 3649(1)$ Å³; $Z = 4$ (two dimers/cell); $\rho_{calc} = 1.623$ g cm^{−3}; $F(000) = 1784$ electrons; $\mu(Mo\text{--}K\alpha) = 12.9$ cm^{−1}. Intensity data were collected on a crystal of dimensions $0.3 \times 0.3 \times 0.1$ mm, within the limit $2\theta_{max} = 50^\circ$ using a Syntex $P2_1$ four-circle diffractometer in conventional $2\theta/\theta$ scan mode; 6446 independent reflections were collected, 4930 with $I > 3\sigma(I)$ being considered 'observed' and used in the 9×9 block diagonal least squares refinement after analytical absorption correction and solution of the structure by vector methods. Anisotropic thermal parameters were refined for the non-hydrogen atoms; hydrogen atoms were included at calculated positions (x, y, z, U_{iso}) and constrained. Residuals (F) at convergence were 0.037, 0.032 (R, R'), reflections weights being $(\sigma^2(F_o) + 0.0005(F_o)^2)^{-1}$. Neutral complex scattering factors were used [5]; computation used the XTAL 83 program system [6] implemented by Dr. S. R. Hall on a Perkin-Elmer 3240 computer.

The complex has minor differences in coordination geometry for the two palladium atoms (Fig. 1). Two 2-benzylpyridine ligands link the palladium atoms as shown schematically in (2), with the square-planar coordination of palladium based on a *trans*-arrangement of nitrogen donors, '*trans*-PdN₂CCl'. Bond angles for the palladium atoms range from 87.7(1)–94.0(2)°, with maximum deviation from the mean planes 'PdN₂CCl' observed for N2 [0.057(6) Å]. The palladium atoms Pd and Pd' are 3.0620(7) Å apart, and the structure of the complex is similar to that reported for the α -picolyl complex *trans*(P,N)-[Pd(μ -pyCH₂-N,CH)(PPh₃)Cl]₂, obtained by oxidative addition of pyCH₂Cl to Pd(PPh₃)₄ [7].

The complex Pd(PyCH₂(C₆H₄)-N,C¹)(3,5-lutidine)Cl·0.2 CH₂Cl₂ [1] may be compared directly with *trans*(N,N)-[Pd(μ -pyCHPh-N,CH)(γ -picoline)-Cl]₂·CH₂Cl₂, as both complexes have a '*trans*-PdN₂-CCl' coordination geometry, but with deprotonated 2-benzylpyridine present as (1) and (2), respectively. Work is in progress on a comparison of the reactivity

of the different coordination forms of deprotonated 2-benzylpyridine.

Acknowledgements

This work was supported by the University of Tasmania and the Australian Research Grants Scheme.

References

- 1 K. Hiraki, Y. Fuchita and K. Takechi, *Inorg. Chem.*, **20**, 4316 (1981).
- 2 A. D. Ryabov and G. M. Kazankov, *J. Organomet. Chem.*, **268**, 85 (1984).
- 3 A. D. Ryabov, *J. Organomet. Chem.*, **268**, 91 (1984).
- 4 A. J. Canty and N. J. Minchin, *Inorg. Chim. Acta*, **100**, L13 (1985).
- 5 'International Tables for X-Ray Crystallography, Vol. 4', Kynoch Press, Birmingham, 1974.
- 6 'The XTAL System', Technical Report TR-1364, Computer Science Center, University of Maryland, 1983.
- 7 K. Nakatsu, K. Kafuku and H. Yamaoka, *Inorg. Chim. Acta*, **54**, L69 (1981).