

N4	0,3971 (3)	0,2354 (4)	0,1038 (2)	2,19 (5)
C5	0,5086 (3)	0,2750 (5)	0,1563 (2)	2,44 (6)
O3	0,1683 (2)	0,1925 (4)	0,1316 (2)	3,02 (5)
O5	0,6253 (2)	0,2968 (5)	0,1294 (2)	3,67 (6)
C4	0,3950 (4)	0,2373 (7)	0,0056 (2)	3,38 (7)

Tableau 2. Paramètres géométriques (Å, °)

N1—N2	1,418 (3)	N4—C5	1,374 (4)
N1—C5	1,355 (4)	N4—C4	1,457 (4)
N2—C3	1,348 (4)	C5—O5	1,229 (4)
C3—N4	1,373 (4)	N1—H1	0,98 (4)
C3—O3	1,230 (4)	N2—H2	1,04 (5)
N2—N1—C5	107,2 (3)	N1—C5—N4	107,2 (3)
N1—N2—C3	107,3 (2)	N1—C5—O5	126,5 (3)
N2—C3—N4	107,3 (2)	N4—C5—O5	126,2 (3)
N2—C3—O3	127,0 (3)	N2—N1—H1	120 (3)
N4—C3—O3	125,7 (3)	C5—N1—H1	121 (2)
C3—N4—C5	109,4 (3)	N1—N2—H2	117 (3)
C3—N4—C4	125,0 (3)	C3—N2—H2	127 (3)
C5—N4—C4	125,2 (3)		

Tableau 3. Distances interatomiques (Å)

O3...H1 ⁱ	1,84 (4)	O3...N1 ⁱ	2,810 (4)
O5...H2 ⁱⁱ	1,80 (5)	O5...N2 ⁱⁱ	2,786 (4)

Codes de symétrie: (i) $x - \frac{1}{2}, y, \frac{1}{2} - z$; (ii) $\frac{1}{2} + x, y, \frac{1}{2} - z$.

Tableau 4. Déviation (Å) par rapport au plan moyen de C3, N4 et C5

	C3	N4	C5	N1	N2
Méthylurazole*	0,0	0,0	0,0	-0,060 (3)	0,110 (3)
Urazole†	0,0	0,0	0,0	-0,032 (3)	0,107 (3)

Equations cartésiennes des plans moyens:

$$* 0,1726x - 0,9818y + 0,0797z + 0,6998 = 0.$$

$$\dagger 0,9048x + 0,3781y - 0,1958z - 3,6833 = 0.$$

Les paramètres isotropiques de H1 et H2 dont les positions ont été affinées sont fixés à 4. Pour les autres, les valeurs ont été fixées 1,3 fois plus grand que celle de l'atome C4 auquel ils sont attachés.

La résolution de la structure, l'affinement et l'analyse des données cristallographiques ont été faites avec le programme SDP (Frenz, 1978) du CAD-4. La structure a été affinée par la méthode des moindres carrés minimisant la fonction $\sum w(|F_o| - |F_c|)^2$. La figure a été dessinée par ORTEPII (Johnson, 1976).

Ce travail est financé par le Ministère Français de la Coopération que nous remercions. Nous remercions aussi Mr Marcel Pierrot, Directeur de Recherche au Laboratoire de Bio-Inorganique Structurale de l'Université d'Aix-Marseille III, pour les facilités d'accès au diffractomètre et aux équipements informatiques.

Les listes des facteurs de structure, des facteurs d'agitation thermique anisotrope, des coordonnées des atomes d'hydrogène, des distances et angles des atomes d'hydrogène, ont été déposées au dépôt d'archives de l'UICr (Référence: PA1153). On peut en obtenir des copies en s'adressant à: The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, Angleterre.

Références

Agmon, I., Ashkenazi, P. & Kaftory, M. (1993). *Acta Cryst.* **C49**, 911–914.

Agmon, I., Kaftory, M., Nelsen, S. F. & Blackstock, S. C. (1986). *J. Am. Chem. Soc.* **108**, 4477–4484.

Belaj, F. (1992). *Acta Cryst.* **C48**, 1088–1090.

Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.

Frenz, B. A. (1978). *The Enraf-Nonius CAD-4 SDP – a Real-Time System for Concurrent X-ray Data Collection and Crystal Structure Determination. Computing in Crystallography*, édité par H. Schenk, R. Olthof-Hazekamp, H. van Koningsveld & G. C. Bassi, pp. 64–71. Version 2.2. VAXSDP. Delft Univ. Press.

Johnson, C. K. (1976). *ORTEPII*. Rapport ORNL-5138. Oak Ridge National Laboratory, Tennessee, EU.

Kaftory, M. (1980). *Acta Cryst.* **B36**, 597–606.

Nelsen, S. F., Petillo, A. P., Chang, H., Frigo, B. T., Dougherty, A. D. & Kaftory, M. (1991). *J. Org. Chem.* **56**, 613–618.

Acta Cryst. (1995). **C51**, 1442–1444

7-Hydroxy-4'-methoxyflavanone

ENGİN KENDİ AND SÜHEYLA ÖZBEY

Department of Physics Engineering, Hacettepe University, 06532 Beytepe, Ankara, Turkey

RAHMIYE ERTAN

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Ankara, 06100 Besevler, Ankara, Turkey

HOONG-KUN FUN AND BOON-CHUAN YIP

School of Physics, University Sains Malaysia, 11800 USM, Penang, Malaysia

(Received 12 August 1994; accepted 13 January 1995)


Abstract

The γ -pyrone ring of the title molecule [2,3-dihydro-7-hydroxy-2-(4-methoxyphenyl)-4H-1-benzopyran-4-one, C₁₆H₁₄O₄] adopts the *C*₂-sofa conformation; the C2 and C3 atoms show disorder. The 2-phenyl substituent is twisted out of the benzo- γ -pyrone ring. The C3—C2—C20—C21 torsion angle is 109.5(4)° and the C3'—C2'—C20—C21 torsion angle is 16.7(7)°.

Comment

Many flavonoid compounds have recognized therapeutic properties. The medicinal activity of flavonoids depends upon the structure and relative orientation of the various rings in the molecule (Cody, 1988). The most studied of these structures are the flavones. In contrast, only few single-crystal X-ray studies of flavanones have been reported so far (Mariezcurrena, 1978; Tomlin & Cantrell, 1990; Kendi, Özbey, Ertan, Fun & Yip, 1995). Flavanones have recognized spasmolytic, expectorant,


anti-ulcer, liver-protecting, anti-inflammatory and antimicrobial properties (Gabor, 1986). To investigate the spasmolytic activity, a series of flavanone ether derivatives, including the title compound, have been synthesized. The determination of the structure of the title compound, (I), was undertaken to study the conformation of the molecule and to contribute to the growing structural data becoming available for the characterization of structure–activity relationships.



The γ -pyrone ring (ring C) exhibits a distorted C_2 -sofa conformation. Atoms C2 and C3 are disordered, with occupancy factors of 0.506 (5) for the sites C2 and C3 and 0.494 (5) for the sites C2' and C3'. The distances of the disordered atoms from the C9–C10–C4–O1 plane for C2, C3, C2' and C3' are 0.473 (6), -0.332 (6), -0.327 (7) and 0.421 (8) Å, respectively. The C2–C3' bond does not exist. The chirality of the molecule is the same whether the partial atoms C2 and C3 are present or the alternative sites are occupied.

The phenyl ring (ring B) is planar; the maximum deviation from the plane is 0.014 Å for C22. The angle between this plane and that of the γ -pyrone ring is 65.2 (1)°. This angle is 70.8° in 5,7,4'-trimethoxyflavanone (Mariezcurrana, 1978), 103.0 (1)° in 7-hydroxyflavanone and 121.1 (2)° in (4-oxo-2-phenyl-2,3-dihydrobenzopyran-7-yl)oxyacetic acid ethyl ester (Kendi, Özbeý, Ertan, Fun & Yip, 1995). The methoxy group is slightly twisted with a C24–C23–O23–C11 torsion angle of -2.2 (5)°. The bond distances agree well with the normally accepted values. The C2–C3 bond in ring C is single [1.528 (7) Å] and C2'–C3' is 1.498 (7) Å. This bond is 1.511 (5) Å in 3-chloroflavanone (Tomlin & Cantrell, 1990) and 1.502 (5) Å in 7-hydroxyflavanone.

The structure is stabilized by intermolecular hydrogen bonds between O7 and O4 [O7–H7A 0.850 (2), H7A···O4 1.842 (2) Å, O7–H7A···O4 176.44 (19)°].



C6	0.4736 (2)	0.1077 (3)	1.14838 (14)	0.0639 (6)
C7	0.4273 (2)	-0.0328 (3)	1.08669 (13)	0.0546 (5)
O7	0.46299 (13)	-0.1800 (2)	1.13346 (11)	0.0705 (4)
C8	0.3493 (2)	-0.0178 (2)	0.98327 (13)	0.0527 (5)
C10	0.3638 (2)	0.2798 (2)	0.99995 (15)	0.0572 (5)
C11	-0.2110 (3)	0.1647 (4)	0.3367 (2)	0.1029 (10)
C21	0.1525 (2)	0.3363 (3)	0.6008 (2)	0.0841 (8)
C22	0.0748 (2)	0.3225 (4)	0.4986 (2)	0.0795 (7)
C23	-0.0312 (2)	0.2513 (3)	0.4725 (2)	0.0666 (6)
O23	-0.1017 (2)	0.2397 (2)	0.36774 (12)	0.0870 (5)
C24	-0.0602 (2)	0.2005 (4)	0.5492 (2)	0.0926 (9)
C25	0.0184 (3)	0.2177 (4)	0.6522 (2)	0.1135 (11)

† Site occupancy = 0.506 (5).

‡ Site occupancy = 0.494 (5).

Table 2. Selected geometric parameters (Å, °)

O1—C9	1.360 (2)	C20—C25	1.375 (4)
O1—C2	1.400 (5)	C5—C6	1.358 (3)
C2—C20	1.515 (5)	C5—C10	1.390 (3)
C2'—C20	1.595 (5)	C6—C7	1.403 (3)
C2—C3	1.528 (7)	C7—O7	1.346 (3)
C2'—C3'	1.498 (8)	C7—C8	1.376 (2)
C3—C4	1.576 (5)	C11—O23	1.412 (3)
C3'—C4	1.476 (7)	C21—C22	1.361 (3)
C4—O4	1.225 (3)	C22—C23	1.372 (3)
C4—C10	1.440 (3)	C23—C24	1.361 (4)
C9—C8	1.385 (3)	C23—O23	1.367 (3)
C9—C10	1.399 (3)	C24—C25	1.376 (3)
C20—C21	1.374 (4)		
C9—O1—C2	118.3 (2)	O1—C2'—C20	101.1 (3)
C9—O1—C2'	113.79 (3)	O1—C2'—C3'	113.5 (4)
O1—C2—C20	108.2 (3)	C4—C3'—C2'	104.7 (4)
O1—C2—C3	105.8 (4)	C6—C5—C10	122.2 (2)
C20—C2—C3	104.8 (4)	C5—C6—C7	119.1 (2)
C2—C3—C4	108.6 (4)	O7—C7—C8	122.6 (2)
O4—C4—C10	123.8 (2)	O7—C7—C6	116.9 (2)
O4—C4—C3	122.4 (3)	C8—C7—C6	120.6 (2)
C10—C4—C3	112.1 (3)	C7—C8—C9	119.3 (2)
C3'—C4—C10	116.1 (3)	C5—C10—C9	117.7 (2)
C3'—C4—O4	117.9 (3)	C5—C10—C4	122.6 (2)
O1—C9—C8	116.5 (2)	C9—C10—C4	119.7 (2)
O1—C9—C10	122.4 (2)	C22—C21—C20	121.5 (3)
C8—C9—C10	121.2 (2)	C21—C22—C23	120.1 (3)
C21—C20—C25	117.5 (2)	C24—C23—O23	124.7 (2)
C21—C20—C2	108.1 (3)	C24—C23—C22	119.8 (2)
C2'—C20—C21	132.9 (3)	O23—C23—C22	115.5 (2)
C2'—C20—C25	108.7 (3)	C23—O23—C11	117.7 (2)
C25—C20—C2	133.5 (3)	C23—C24—C25	119.4 (3)
C20—C2'—C3'	112.8 (4)	C20—C25—C24	121.7 (3)

Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1990b). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993).

The authors would like to thank the Malaysian Government and the Universiti Sains Malaysia for research grant R & D No. 123/3417/2201.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and bond distances and angles involving non-H atoms have been deposited with the IUCr (Reference: AB1220). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Cody, V. (1988). *Progress in Clinical and Biological Research*, Vol. 280, *Plant Flavonoids in Biology and Medicine II: Biochemical, Cellular and Medicinal Properties*, edited by V. Cody, E. Middleton, J. B. Harborne & A. Beretz, pp. 29–45. New York: Alan R. Liss.
- Gabor, M. (1986). *The Pharmacology of Benzopyrone Derivatives and Related Compounds*. Budapest: Akademiai-Kiado.
- Kendi, E., Özbey, S., Ertan, R., Fun, H. K. & Yip, B. C. (1995). *Acta Cryst.* C51. In the press.
- Mariezcurrera, R. A. (1978). *Acta Cryst.* B34, 2322–2324.
- Sheldrick, G. M. (1990a). *SHELXTL/PC*. Release 4.1. Siemens Crystallographic Research Systems, Madison, Wisconsin, USA.
- Sheldrick, G. M. (1990b). *Acta Cryst.* A46, 467–473.
- Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. Univ. of Göttingen, Germany.
- Tomlin, D. W. & Cantrell, J. S. (1990). *Acta Cryst.* C46, 519–521.
- Acta Cryst.* (1995). C51, 1444–1447
- ## Cholesteryl 4-(8-Octenyloxy)benzoate
- MICHAEL L. CHABINYC AND ALBERT V. FRATINI
- University of Dayton, Department of Chemistry,
300 College Park, Dayton, OH 45469-2357, USA*
- EDWARD P. SOCCI AND BARRY L. FARMER
- Department of Materials Science & Engineering,
University of Virginia, Charlottesville,
VA 22903-2442, USA*
- TIMOTHY J. BUNNING* AND WALTER W. ADAMS
- WL/MLPJ, Materials Directorate, Wright-Patterson
Air Force Base, OH 45433-7702, USA*
- (Received 11 July 1994; accepted 18 January 1995)
- ## Abstract
- The crystal structure of cholest-5-en-3-yl 4-(8-octenyloxy)benzoate, C₄₂H₆₄O₃, was determined by single-crystal X-ray diffraction. The benzoate moiety is twisted 64.6 (2)° from the least-squares plane through the tetracyclic core. Elongated molecules pack in an antiparallel fashion. Fully overlapping molecules are related by 2₁ screw symmetry.
- ## Comment
- The title structure, (I), is the second in a series of crystallographic studies of cholesteryl *n*-alkenebenzoate compounds (Socci, Farmer, Chabiny, Fratini, Bunning & Adams, 1995). When these compounds are attached to cyclosiloxane backbones, a class of intermediate molecular weight liquid crystals is produced (Bunning, 1992; Bunning, Klei, Samulski, Crane & Adams, 1993). Ordered transparent films of these materials have potential use as matrices for optical devices. An understanding of the solid-state structure of cholesteryl esters may provide an insight into the types of packing exhibited by these molecules in the liquid-crystalline state. Relationships between the crystal structures and the packing