J.C.S. Снем. Сомм., 1979

Synthesis of Retrochinensin; a New Naturally Occurring 4-Aryl-2,3-naphthalide Lignan

By SHIBNATH GHOSAL* and SHANTA BANERJEE

(Pharmaceutical Chemistry Research Laboratory, Department of Pharmaceutics, Banaras Hindu University, Varanasi-5, India)

Summary 2,3-Dichloro-5,6-dicyanobenzoquinone-induced oxidative cyclization of suchilactone (I) afforded retrochinensin (II), a new naturally occurring 4-aryl-2,3naphthalide lignan, in high yield; treatment of (I) with N-bromosuccimide on the other hand, afforded justicidin B (IV) in high yield.

THE ontogenic and seasonal variations of the lignan constituents of *Polygala chinensis*^{1,2} prompted us to examine the role, if any, of its major monoarylidenebutyrolactone lignan, suchilactone (I), in the formation of the congener 1- and 4aryl-2,3-naphthalide lignans. Previous reports³⁻⁵ suggested diarylidenebutyrolactone lignans as likely intermediates (*via* cyclic peroxides or β -apolignans) to arylnaphthalide lignans.

In the present study, two routes, (i) allylic bromination by N-bromosuccinimide (NBS) followed by dehydrobromination, and (ii) dehydrogenation by 2,3-dichloro-5,6dicyanobenzoquinone (DDQ), were envisaged for the synthesis of the diarylidene from (I). A three-step synthesis of (I) has been reported before.⁶

Refluxing (I) with NBS, in CCl₄, directly afforded justicidin B (IV) in high yield (72%). Dehydrogenation of (I) (1 mmol) with DDQ (1 mmol), in dioxan, on the other hand, gave a 4-aryl-2,3-naphthalide lignan (II), (34%), an unstable lignan-DDQ adduct (III), (15%), and unchanged (I) (43%). The lignan-DDQ adduct was assigned structure (III) on the basis of the following evidence. It showed u.v. maxima similar to those of (I); when the mass spectrum was recorded, it fragmented in the direct inlet system before exhibiting any molecular ion peak but characteristic fragment-ion peaks appeared at m/e 381 (rel. int., 1.2%), 377 (2.8) (species A) 366 (68, diarylidene), 232 (3.8), 231 (6), 228 (12.5), 227 (18) (DDQ-H₂), 217 (100, species B), 187 (5.5), 165 (3), 150 (2), 149 (5), and 121 (2). On treatment with HCl it afforded the corresponding diarylidene and DDQ-H₂. The synthesis of the diarylidene has been reported before.³ When kept in acetone solution, (III) was converted into diphyllin, presumably via the diarylidene, while refluxing in benzene with a further quantity (1:1) of DDQ afforded (II).

Compound (II)[†] crystallized from alcohol as colourless needles, m.p. 234 °C; $R_{\rm f}$ 0·38 ($C_{6}H_{6}$ -HOAc, 98:2), blue fluorescence under short-wave u.v. lamp; $\lambda_{\rm max}$ (log ϵ) 250 (4·61), 315 (3·95), and 350 infl. (3·57) nm; $\nu_{\rm max}$ (Nujol) 1766, 1635, and 938 cm⁻¹; δ (CDCl₃) 8·3 (1H, s, 1-H), 7·36—6·95 (5H, m, aromatic), 6·14 (2H, s, -OCH₂O-), 5·24 (2H, q, lactone -CH₂-), 4·02 (3H, OMe), and 3·91 (3H, OMe); *m/e* 364 (M⁺, 100%, C₂₁H₁₆O₆), 349 (4), 335 (7), 307 (3), and 306 (2). The yield of (II) was much improved (255 mg, 85%) when (I) (298 mg, 0·8 mmol) was refluxed (18 h) with DDO



(365 mg, 1.6 mmol) in anhydrous benzene. The product was obtained as a homogeneous entity by a single crystallization from alcohol.

[†] Munakata *et al.* (K. Munakata, S. Marumo, K. Ohta, and Y.-L. Chen, *Tetrahedron Letters*, 1967, 3821) also synthesized (II) by a multistep procedure and reported its m.p. (280 °C), u.v., and i.r. data; the spectral data are comparable to those given here. Rao *et al.* (B. V. G. Rao, L. R. Row, and P. Satyanarayana, *Indian J. Chem.* (B), 1978, 16, 68) recently obtained (II) in traces, while synthesizing chinensin² and reported only its m.p. (240 °C).

The method exemplifies one of the simplest synthetic processes for 4-aryl-2,3-naphthalide lignans. Incidentally, it provides the first laboratory analogy for the biogenesis of 'retro-disposed' lignans from monoarylidenebutyrolactone intermediates. Compound (II), which we named retrochinensin, has been recently found in Justicia prostata (Acanthaceae) along with a number of 4-aryl-2,3-naphthalide lignans.⁷

We thank Dr. B. C. Das, C.N.R.S., Gif-sur-Yvette, France, and Dr. R. K. Chaudhuri, Pharmazeutisches Institut, Bonn, Germany, for mass and n.m.r. spectra, and the U.G.C., New Delhi, India, for financial support.

(Received, 25th September 1978; Com. 1031.)

- ¹ S. Ghosal, R. B. P. S. Chauhan, and R. S. Srivastava, *Phytochemistry*, 1974, 13, 1933. ² S. Ghosal, R. B. P. S. Chauhan, and R. S. Srivastava, *Phytochemistry*, 1974, 13, 2281.

- S. Ghosai, K. D. S. Ghaudhan, and R. D. Ghrussatra, *Information formation and Pharm. Bull.* (Japan), 1972, 20, 624.
 Y.-T. Lin, T.-B. Lo, K.-T. Wang, and B. Weinstein, *Tetrahderon Letters*, 1967, 849.
 T. Momose, K. Kanai, T. Nakamura, and Y. Kuni, *Chem. and Pharm. Bull.* (Japan), 1977, 25, 2755.
 S. Ghosal and S. Banerjee, Abstracts, Convention of Chemists, Jaipur, 1977, 41.
 S. Chosal A. Chaudhuri and S. Banerjee in propagation.
- ⁷S. Ghosal, A. Chaudhuri, and S. Banerjee, in preparation.