## A Novel Intermolecular Synthesis of $\gamma$ -Lactones via Visible-Light Photoredox Catalysis

## Xiao-Jing Wei,<sup>†</sup> Deng-Tao Yang,<sup>†</sup> Lin Wang,<sup>†</sup> Tao Song,<sup>†</sup> Li-Zhu Wu,<sup>\*,‡</sup> and Qiang Liu<sup>\*,†</sup>

State Key Laboratory of Applied Organic Chemistry, Lanzhou University, Lanzhou 730000, PR China, and Key Laboratory of Photochemical Conversion and Optoelectronic Materials, Technical Institute of Physics and Chemistry, Chinese Academy of Sciences, Beijing 100190, PR China

liuqiang@lzu.edu.cn; lzwu@mail.ipc.ac.cn

## Received October 13, 2013



Direct  $\gamma$ -lactone formation via visible-light photoredox catalysis has been achieved efficiently including hydroxylalkylation of aromatic alkenes and transesterification. The present photocatalytic protocol has good regioselectivity and substrate compatibility, affording a novel way to intermolecular  $\gamma$ -lactone synthesis by the reaction of styrenes with  $\alpha$ -bromo esters in the absence of any external oxidants.

 $\gamma$ -Lactones are basic structural elements in many natural products and are prevalently used building blocks in agrichemicals, pharmaceuticals and organic materials.<sup>1</sup> A tremendous amount of effort has been devoted and much progress has been made to introduce the  $\gamma$ -lactone moiety into organic molecules,<sup>2</sup> among which the intra- or intermolecular reactions of alkenes with nucleophiles<sup>3</sup> or radical precursors<sup>4</sup> are most common approaches to construct the skeletons. However, stoichiometric oxidants and harsh reaction conditions are always employed to activate alkenyl moieties or produce radicals in the transformations. Moreover, the addition of oxidant tends to cause a crisis of undesired side reactions and may lead to pollution of the environment. As a result, an efficient and environmentally friendly method should be developed.

ORGANIC LETTERS

XXXX Vol. XX, No. XX

000-000

As an efficient and sustainable method to perform organic reactions, visible-light photoredox catalysis has attracted widespread research interest owing to its attractive features such as mild and green conditions, excellent functional group tolerance, and high reactivity.<sup>5</sup> In addition, as the main wave band of solar energy, visible light is generally readily available and can easily be used without the need of a special instrument or apparatus. Moreover, compared to ultraviolet irradiation, the transparent properties of most organic substrates for visible light minimize the side reactions.

Over these years, a variety of elegant chemical transformations via visible-light photoredox have been developed.<sup>6</sup> Seminal works by MacMillian,<sup>7a,b</sup> Stephenson,<sup>7c,d</sup> and Gagné<sup>7e</sup> et al. have demonstrated that activated haloalkanes such as carbonyl, benzyl, alkenyl, aryl, and heteroatom

<sup>&</sup>lt;sup>†</sup>Lanzhou University.

<sup>&</sup>lt;sup>‡</sup> Technical Institute of Physics and Chemistry.

 <sup>(1) (</sup>a) Nefkens, G. H. L.; Thuring, J. W. J. F.; Beenakkers, M. F. M.; Zwanenburg, B. J. Agric. Food Chem. **1997**, 45, 2273. (b) Mangnus, E. M.; Zwanenburg, B. J. Agric. Food Chem. **1992**, 40, 1066. (c) Fang, B.; Xie, X.; Zhao, C.; Jing, P.; Li, H.; Wang, Z.; Gu, J.; She, X. J. Org. Chem. **2013**, 78, 6338.

<sup>(2) (</sup>a) Gil, S.; Parra, M.; Rodriguez, P.; Segura, J. *Mini-Rev. Org. Chem.* **2009**, *6*, 345. (b) Cheng, X.-F.; Li, Y.; Su, Y.-M.; Yin, F.; Wang, J.-Y.; Sheng, J.; Vora, H. U.; Wang, X.-S.; Yu, J.-Q. *J. Am. Chem. Soc.* **2013**, *135*, 1236. (c) Newhouse, T. R.; Kaib, P. S. J.; Gross, A. W.; Corey, E. J. Org. Lett. **2013**, *15*, 1591.

<sup>(3) (</sup>a) Huang, L.; Jiang, H.; Qi, C.; Liu, X. J. Am. Chem. Soc. 2010, 132, 17652. (b) Xie, X.; Li, Y.; Fox, J. M. Org. Lett. 2013, 15, 1500.

<sup>(4) (</sup>a) Powell, L. H.; Docherty, P. H.; Hulcoop, D. G.; Kemmitt, P. D.; Burton, J. W. *Chem. Commun.* **2008**, 2559. (b) Wang, W.; Xu, M.-H.; Lei, X.-S.; Lin, G.-Q. *Org. Lett.* **2000**, *2*, 3773. (c) Iwahama, T.; Sakaguchi, S.; Ishii, Y. *Chem. Commun.* **2000**, 613.

<sup>(5)</sup> Reviews: (a) Narayanam, J. M. R.; Stephenson, C. R. J. Chem. Soc. Rev. 2011, 40, 102. (b) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Chem. Rev. 2013, 113, 5322. (c) Yoon, T. P.; Ischay, M. A.; Du, J. Nat. Chem. 2010, 2, 527. (d) Xuan, J.; Xiao, W.-J. Angew. Chem. 2012, 124, 6934. Angew. Chem., Int. Ed. 2012, 51, 6828. (e) Shi, L.; Xia, W. Chem. Soc. Rev. 2012, 41, 7687.

Table 1. Screening of Reaction Conditions<sup>a</sup>

Ph Ph 1a	+ Br CO <sub>2</sub> Et CO <sub>2</sub> Et <b>2a</b>	fac-Ir(ppy)₃ (0.5 mol %) hv = 450 nm solvent, additives	$Ph$ $CO_2Et$ Ph $3a$
entry		conditions	yield (%)
1	(CH <sub>3</sub> ) <sub>2</sub> CO:H	H <sub>2</sub> O = 9:1	28
2	DMF:H <sub>2</sub> O =	= 9:1	74
3	CH <sub>3</sub> CN:H <sub>2</sub> C	D = 9:1	80
4	CH <sub>3</sub> CN:H <sub>2</sub> C	D = 4:1	82
5	CH <sub>3</sub> CN:H <sub>2</sub> C	O = 4:1; LiBr (1.2 equiv)	79
6	CH <sub>3</sub> CN:H <sub>2</sub> C	O = 4:1; LiCl (1.2 equiv)	84
7	CH <sub>3</sub> CN:H <sub>2</sub> C	$O = 4:1; LiBF_4 (1.2 equiv)$	) 93
8	CH <sub>3</sub> CN:H <sub>2</sub> C	$O = 4:1; LiBF_4 (0.5 equiv)$	) 87

<sup>*a*</sup> Reaction conditions: a mixture of **1a** (0.20 mmol), **2a** (0.24 mmol) and photocatalyst *fac*-Ir(ppy)<sub>3</sub> (0.5 mol %) in solvent was irradiated with a 3 W blue LEDs at room temperature for 24 h. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> Irradiated without *fac*-Ir(ppy)<sub>3</sub>. <sup>*d*</sup> The reaction was conducted in the dark. <sup>*e*</sup> N.D. = not detected. <sup>*f*</sup> N.R. = no reaction.

CH<sub>3</sub>CN, LiBF<sub>4</sub> (1.2 equiv) (no H<sub>2</sub>O)

 $CH_3CN:H_2O = 4:1, LiBF_4 (1.2 equiv)$ 

 $CH_{3}CN:H_{2}O = 4:1, LiBF_{4}$  (1.2 equiv)

N.D.

N.R.

NR

halides are efficient precursors for the generation of carboncentered radicals under photoredox catalytic conditions. In some of these reactions both the quenching and the regeneration of the photocatalyst are promoted by the substrates or intermediates of the catalyzed reaction; therefore, external reductants or oxidants are no longer needed.<sup>7</sup>

With this in mind, we initiated the study by using 1,1diphenylethylene **1a** and  $\alpha$ -bromo diethyl malonate **2a** with catalytic amount of *fac*-Ir(ppy)<sub>3</sub>. It was encouraging to see that the aimed product **3a** was obtained in 28% yield after 24 h of irradiation (blue LEDs,  $\lambda = 450$  nm) at room temperature when the reaction was performed in a mixture of acetone and water (9:1 v/v) (Table 1, entry 1). The replacement of acetone by *N*,*N*-dimethylformamide (DMF) or acetonitrile resulted in a significant improvement of the yield (Table 1, entries 2 and 3), and acetonitrile was chosen as the ideal organic solvent for the reaction. Next, we investigated the amount of water for the transformation; the result proved that acetonitrile/water (4:1) gained a better result (Table 1, entry 4). Considering the activation

 $9^e$ 

 $10^{c,f}$ 

 $11^{d,f}$ 

**Table 2.** Scope of Visible-Light-Induced  $\gamma$ -Lactones Synthesis with  $\alpha$ -Bromo Esters  $\mathbf{2}^a$ 

Ph Ph 1a	$ \begin{array}{c} \operatorname{Br} \stackrel{R^{1}}{\underset{CO_{2}}{\operatorname{R}^{3}}} \operatorname{R}^{2} \stackrel{fau}{\underset{CO_{2}}{\operatorname{R}^{3}}} \\ 2 \end{array} ( $	c-lr(ppy) <sub>3</sub> (0.5 mol %) hv = 450  nm CH <sub>3</sub> CN/H <sub>2</sub> O, LiBF <sub>4</sub>	%) →	$Ph$ $R^1$ $R^2$ $R^1$ $R^2$ $R^3$
entry	$\alpha$ -bromo esters	product		yield $(\%)^b$
1	$EtO_2C$ Br $CO_2Et$ 2a	EtO <sub>2</sub> C O Ph	3a	93
2	MeO <sub>2</sub> C Br CO <sub>2</sub> Me 2b	MeO <sub>2</sub> C O O Ph	3b	95
3	$EtO_2C$ Br $CO_2Et$ 2c	EtO <sub>2</sub> C OPh	3c	98
4	$\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{C}}}}}_{Br}^{Br}}_{CO_2Et}$	O Ph Ph	3d	94
5	$Ph Br CO_2Et 2e$	Ph O O Ph	3e	94
6	F Br CO <sub>2</sub> Et 2f	Ph O Ph	3f	92
7	$F \xrightarrow{F} Br CO_2Et 2g$	F O O Ph	3g	60
8	Br CO <sub>2</sub> Et 2h	O O Ph	3h	99

<sup>*a*</sup> Reaction conditions: a mixture of **1** (0.20 mmol), **2** (0.24 mmol), LiBF<sub>4</sub> (0.24 mmol) and photocatalyst *fac*-Ir(ppy)<sub>3</sub> (0.5 mol %) in CH<sub>3</sub>CN/H<sub>2</sub>O (4:1 v/v) was irradiated with a 3 W blue LEDs at room temperature for 24 h. <sup>*b*</sup> Isolated yield.

of carbonyl moiety may be beneficial for the production of  $\gamma$ -lactones, a brief screen of lithium salts additives was performed (Table 1, entries 5–7). The best result was achieved when 1.2 equiv of LiBF<sub>4</sub> was introduced to the system. In this case we obtained the desired  $\gamma$ -lactone **3a** in 93% isolated yield. In addition, the attempt to reduce the amout of LiBF<sub>4</sub> slightly decreased the yield (Table 1, entry 8). It was worth noting that no product was detected when the reaction was carried out in dry acetonitrile (Table 1, entry 9). Moreover, when the reaction was carried out in the absence of *fac*-Ir(ppy)<sub>3</sub> (Table 1, entry 10) or in the dark (Table 1, entry 11), no conversion could be observed, which indicated that the reaction was indeed a visible-light-driven photoredox process.

On submission to the optimized conditions, we next investigated the scope of  $\alpha$ -bromo esters **2** for the reaction.  $\alpha$ -Bromo dimethyl malonate **2b** can also react with **1a** to produce **3b** in excellent yield (Table 2, entry 2). Moreover, this kind of reaction was successfully applied to ethyl

<sup>(6)</sup> Selected recent examples: (a) Nagib, D. A.; MacMillan, D. W. C. *Nature* **2011**, *480*, 224. (b) Ischay, M. A.; Anzovino, M. E.; Du, J.; Yoon, T. P. *J. Am. Chem. Soc.* **2008**, *130*, 12886. (c) Cai, S.; Zhao, X.; Wang, X.; Liu, Q.; Li, Z.; Wang, D. Z. *Angew. Chem., Int. Ed.* **2012**, *51*, 8050. (d) Hari, D. P.; Schroll, P.; König, B. *J. Am. Chem. Soc.* **2012**, *134*, 2958. (e) Sun, H.; Yang, C.; Gao, F.; Li, Z.; Xia, W. *Org. Lett.* **2013**, *15*, 624. (f) Zhu, S.; Das, A.; Bui, L.; Zhou, H.; Curran, D. P.; Rueping, M. J. Am. Chem. Soc. **2013**, *135*, 1823. (g) Ye, Y.; Sanford, M. S. *J. Am. Chem. Soc.* **2012**, *134*, 9034.

<sup>(7) (</sup>a) Nicewicz, D. A.; MacMillan, D. W. C. Science 2008, 322, 77.
(b) Shih, H.-W.; Vander Wal, M. N.; Grange, R. L.; MacMillan, D. W. C. J. Am. Chem. Soc. 2010, 132, 13600. (c) Narayanam, J. M. R.; Tucker, J. W.; Stephenson, C. R. J. J. Am. Chem. Soc. 2009, 131, 8756. (d) Nguyen, J. D.; D'Amato, E. M.; Narayanam, J. M. R.; Stephenson, C. R. J. Nat. Chem. 2012, 4, 854. (e) Andrews, R. S.; Becker, J. J.; Gagné, M. R. Angew. Chem., Int. Ed. 2010, 49, 7274. (f) Liu, Q.; Yi, H.; Liu, J.; Yang, Y.; Zhang, X.; Zeng, Z.; Lei, A. Chem.—Eur. J. 2013, 19, 5120. (g) Su, Y.; Zhang, L.; Jiao, N. Org. Lett. 2011, 13, 2168.

Table 3. Reaction of 1a with Various Alkyl Bromoacetates<sup>a</sup>

*	→ <sup>Ph</sup> Ph + <b>1a</b>	Br CO <sub>2</sub> R <b>2</b>	fac-lr(ppy) <sub>3</sub> (0.5 mol %) $h_{V} = 450 \text{ nm}$ CH <sub>3</sub> CN/H <sub>2</sub> O, LiBF <sub>4</sub>	· Ph Ph <b>3i</b>
entry		bro	yield (%) <sup>l</sup>	
1		<b>2i</b> (R	99	
<b>2</b>		<b>2j</b> (R	99	
3		<b>2k</b> (I	$29 \ (67)^c$	
4		<b>21</b> (R	70	
<b>5</b>		<b>2m</b> (	75	
6		<b>2n</b> (I	$R = p - NO_2 Ph$	23

 $^a$  The reaction conditions as described in Table 2.  $^b$  Isolated yields.  $^c$  Yield in bracket performed with 2.0 equiv of 2,6-lutidine.

bromoacetate with substituted groups such as methyl and phenyl group. The corresponding  $\gamma$ -lactone products 3c-3ewere obtained in excellent yields (98–94%). Reaction of 2fand 2g proceed effectively, and lactones with fluorine group such as 3f, 3g might be helpful in the pharmaceutical and agrochemical fields (Table 2, entries 6 and 7). Ethyl 2-bromo-2-methylpropanoate 2h was also smoothly transformed to corresponding lactone with excellent yield (Table 2, entry 8).

We next studied the reactivities of bromoacetates with various alkoxyl groups<sup>8</sup> in the visible-light-promoted  $\gamma$ -lactone synthesis. The results showed that methyl and ethyl 2-bromoacetate were completely converted into  $\gamma$ -lactones, respectively (Table 3, entries 1 and 2). Trimethyl-silyl and phenyl 2-bromoacetate could also produce the same product **3i**, even though the reaction proceeded less efficiently than the former cases (Table 3, entries 4 and 5). Finally, we found the *tert*-butyl and *p*-nitrophenyl just could afford small amount of target compound (Table 3, entries 3 and 6), but in the case of *tert*-butyl 2-bromoacetate the yield could be improved to 67% by addition of a base 2,6-lutidine.

Having found a broad range of  $\alpha$ -carbonyl alkyl bromides, the substrate scope with regard to substituted styrene was examined. As shown in Table 4, both styrene and *p*-methoxy-substituted styrene could obtain the  $\gamma$ -lactones with good yields (Table 4, entries 1 and 2). Not surprisingly, the  $\alpha$  and  $\beta$  methyl substituted styrene yielded the aimed products with good to excellent yields (Table 4, entries 3 and 4). Biaryl lactone **3n** was obtained in similar way (Table 4, entry 5). It is noteworthy that when conjugated benzyl diene (*E*)-buta-1,3-dien-1-ylbenzene **1g** was used, the desired product **3o** was isolated in 78% yield with exclusive regioselectivity toward the terminal olefin (Table 4, entry 6). At the same time, benzocycloalkene (Table 4, entries 7 and 8) gave **Table 4.** Scope of Visible-Light-Induced  $\gamma$ -Lactones Synthesis with Styrenes  $\mathbf{1}^a$ 



<sup>*a*</sup> The reaction conditions as described in Table 2. <sup>*b*</sup> Isolated yield. <sup>*c*</sup> dr and E/Z were determined by <sup>1</sup>H NMR spectrum.

the tricyclic  $\gamma$ -lactones with moderate to good yields, which had been employed as starting materials for the synthesis of strigolactone analogues.<sup>1a</sup>

Since both bromoacetates and styrenes are transparent to 450-nm photons (see the Supporting Information), only the photocatalyst *fac*-Ir(ppy)<sub>3</sub> can be excited under the

<sup>(8)</sup> Yoo, D.-W.; Han, J.-H.; Nam, S. H.; Kim, H. J.; Kim, C.; Lee, J. K. Inorg. Chem. Commun. 2006, 9, 654.

<sup>(9) (</sup>a) Maslak, V.; Matović, R.; Saičić, R. N. *Tetrahedron* 2004, 60, 8957.
(b) Bercot, E. A.; Kindrachuk, D. E.; Rovis, T. Org. Lett. 2005, 7, 107.
(c) Iwahama, T.; Sakaguchi, S.; Ishii, Y. Chem. Commun. 2000, 613.
(d) Ko, Y. O.; Chun, Y. S.; Kim, Y.; Kim, S. J.; Shin, H.; Lee, S.-g. Tetrahedron Lett. 2010, 51, 6893.

<sup>(10)</sup> For representative alkene difunctionalization examples with photoredox catalysis, see: (a) Yasu, Y.; Koike, T.; Akita, M. Org. Lett. **2013**, *15*, 2136. (b) Fumagalli, G.; Boyd, S.; Greaney, M. F. Org. Lett. **2013**, *15*, 4398. (c) Sahoo, B.; Hopkinson, M. N.; Glorius, F. J. Am. Chem. Soc. **2013**, *135*, 5505. (d) Yasu, Y.; Koike, T.; Akita, M. Angew. Chem., Int. Ed. **2012**, *51*, 9567. (e) Wilger, D. J.; Gesmundo, N. J.; Nicewicz, D. A. Chem. Sci. **2013**, *4*, 3160. (f) Yoo, W.-J.; Kobayashi, S. Green Chem. **2013**, *15*, 1844.



**Figure 1.** Photoluminescence quenching experiments. (A) Luminescence quenching of *fac*-Ir(ppy)<sub>3</sub> ( $2.00 \times 10^{-4}$  M) with progressive addition of bromoacetate **2a**. (B) Luminescence quenching of *fac*-Ir(ppy)<sub>3</sub> ( $2.00 \times 10^{-4}$  M) with progressive addition of styrene **1a**.

irradiation of blue LEDs (blue LEDs,  $\lambda = 450$  nm). As shown in Figure 1, excitation of the characteristic absorption of fac-Ir(ppy)<sub>3</sub> resulted in a maximal photoluminescence at 513 nm in acetonitrile/water (4:1, v/v) solution. which was quenched by bromoacetate 2a with a rate constant of  $6.00 \times 10^4 \,\text{L mol}^{-1}$ . Although the luminescence of fac-Ir(ppy)<sub>3</sub> can also be quenched by styrene 1a, the rate was smaller  $(2.90 \times 10^4 \text{ L mol}^{-1})$ . As the spectral overlap of absorption of bromoacetate 2a and photoluminescence of fac-Ir(ppy)<sub>3</sub> is rather small, the energy transfer from the excited fac-Ir(ppy)<sub>3</sub> to **2a** would be negligible. The photoluminescence quenching is therefore attributed to the electron transfer from the excited fac-Ir(ppy)<sub>3</sub> to bromoacetate 2a. The ET procedure generates electron-deficient radical 4 along with  $\hat{fac}$ -Ir<sup>IV</sup>(ppy)<sub>3</sub>. Then intermolecular  $\pi$ -addition of the radical 4 to styrene 1a produces a benzylic radical **5**. Further ET from radical **5** to fac-Ir<sup>IV</sup>(ppy)<sub>3</sub> affords carbocation  $\mathbf{6}$  and regenerates the photocatalyst. The nucleophilic attack of the carbocation 6 by water results in the hydroxyalkylation of styrene, which was proved by H<sub>2</sub><sup>18</sup>O labeling experiment (see the Supporting Information). Finally, a spontaneous intramolecular transesterfication promoted by LiBF<sub>4</sub> provided the desired  $\gamma$ -lactone **3a** (Scheme 1).<sup>9</sup>

Scheme 1. Proposed Mechanism Cycle



In summary, we have developed a catalytic, externaloxidant-free method for the construction of aryl-substituted  $\gamma$ -lactone via photoredox catalysis for the first time. The highlight of this operationally simple reaction is the avoidance of stoichiometric oxidants, low catalyst loading, and the generation of  $\gamma$ -lactone in high yields under mild conditions. Moreover, the photocatalytic transformation is also a beneficial supplement for alkene difunctionalization with photoredox catalysis.<sup>10</sup> Further explorations on the method to build  $\delta$ -lactone mediated by visible light is under investigation in our laboratory.

Acknowledgment. We are grateful for financial support of NSFC (No. 21172102, 21090343), the Ministry of Science and Technology of China (2013CB834804), and the Fundamental Research Funds for the Central Universities.

**Supporting Information Available.** Experimental procedure, characterization data for the products and UV/vis absorption spectra. This material is available free of charge via the Internet at http://pubs.acs.org.

The authors declare no competing financial interest.