


This article was downloaded by: [FU Berlin]

On: 11 May 2015, At: 10:41

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Carbohydrate Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcar20>

The Tmsotf-Promoted “One Pot” β -Glycosidations of Peracetylated Chitobiose

Biao Yu ^a, Qinqin Ouyang ^a, Chuan Li ^a & Yongzheng Hui ^a

^a State Key Laboratory of Bio-organic and Natural Products Chemistry, Shanghai Institute of Organic Chemistry, Academia Sinica 354 Fenling Lu, Shanghai, 200032, P.R. China

Published online: 22 Aug 2006.

To cite this article: Biao Yu, Qinqin Ouyang, Chuan Li & Yongzheng Hui (1996) The Tmsotf-Promoted “One Pot” β -Glycosidations of Peracetylated Chitobiose, Journal of Carbohydrate Chemistry, 15:3, 297-302, DOI: [10.1080/07328309608005654](https://doi.org/10.1080/07328309608005654)

To link to this article: <http://dx.doi.org/10.1080/07328309608005654>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the “Content”) contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

THE TMSOTf-PROMOTED "ONE POT" β -GLYCOSIDATIONS OF PERACETYLATED CHITOBIOSE

Biao Yu, Qinqin Ouyang, Chuan Li, Yongzheng Hui*

State Key Laboratory of Bio-organic and Natural Products Chemistry
Shanghai Institute of Organic Chemistry, Academia Sinica
354 Fenling Lu, Shanghai 200032, P.R.China

Received August 8, 1995 - Final Form December 11, 1995

ABSTRACT

β -Glycosidations of peracetylated chitobiose with such alcohols as methyl, allyl, benzyl, isopropyl, *tert*-butyl alcohol and 1,2:3,4-*O*-diisopropylidene-1-*O*- α -D-galactoside were carried out in good yields by employing TMSOTf as a promoter. The reaction proceeded through the oxazoline intermediate.

INTRODUCTION


The β -(1 \rightarrow 4) linked *N*-acetylglucosamine chitooligosaccharide fragments are widely distributed in many biologically important compounds. The inner core region of cell surface *N*-glycoproteins consists of a chitobiose substructure.¹ Chitobiose is also found as repeating units of the bacterial cell wall peptidoglycan.² On the other hand, chitooligosacchrides play significant roles in eliciting defense related responses in various plants.³ The tetra- and pentachitosaccharides are the backbone of the nodulation factors on legume roots.⁴ Furthermore, some chitooligosaccharides and their derivatives possess immunostimulating and antitumor activities.⁵

The synthetic approaches to these compounds have relied mainly on a variety of coupling reactions of glucosamine derivatives.⁶ A facile way to achieve these biologically interesting compounds is to employ peracetylated chitobiose as a starting material,^{7,8} which is readily accessible from chemical or enzymatic degradation of chitin.^{8,9} Only based on an efficient glycosidation reaction, can the peracetylated chitobiose be changed into proper acceptors and donors. The glycosidation of 2-acetamido-2-deoxy sugars mainly rely on the oxazoline method.¹⁰ By treatment with TMSOTf (trimethylsilyl triflate), the peracetylated chitobiose was easily changed to its oxazoline in high yield, the glycosidation reactions of this oxazoline with such simple alcohols as methanol, 2-propen-1-ol, and 2,2,2-trichloroethanol in the presence of trifluoromethanesulfonic acid as catalyst gave the corresponding β -glycosides in 76%, 53%, and 40% yield, respectively.⁸ On the other hand, the glycosidations of 2-methyl (3,4,6-tri-*O*-acetyl-1,2-dideoxy- α -D-glucopyranose [2,1-d]-2-oxazoline¹¹ and 2-methyl-4-*O*-(2,3,4,6-tetra-*O*-acetyl- β -D-galactosyl)-3,6-di-*O*-acetyl-1,2-dideoxy- α -D-glucopyranose [2,1-d]-2-oxazoline¹² can be promoted by TMSOTf.

We report here facile TMSOTf-promoted “one pot” β -glycosidation reactions of peracetylated chitobiose **1**.

RESULTS AND DISCUSSION

Peracetylated chitobiose was transformed completely into its oxazoline derivative by treatment with TMSOTf (1.1 equiv) at 50 °C overnight. After the mixture was cooled to room temperature, it was stirred under dry 4 Å molecular sieve for a period before the alcohol was added, otherwise, the yield of glycoside was drastically lowered. The results are summarized in the Scheme. The yields of glycosides were good with all alcohol glycosyl acceptors we employed, including isopropyl and *tert*-butyl alcohols. Only the β -anomers were found in their ¹H NMR spectra. On the other hand, the intermediate oxazoline can be separated and purified by silica gel chromatography,¹³ which can be also converted into corresponding glycosides in good yield by employing TMSOTf (0.2 equiv) as catalyst. By employing FeCl₃¹⁴ or BF₃OEt₂¹⁵ as catalyst, the yields of the glycosides were less than 35% with allyl and benzyl alcohol as the glycosyl acceptors, and no glycoside product was observed with isopropyl alcohol as the glycosyl acceptor.



Scheme

EXPERIMENTAL

General Methods. Solvents were purified in the usual way. Melting points are uncorrected. Optical rotations given in units of $10^{-1}\text{deg cm}^2\text{g}^{-1}$, were determined with a Perkin-Elmer Model 241 MC polarimeter at 22 °C. IR spectra were measured with IR-440 spectrometer. ^1H NMR spectra were recorded on Bruker AM-300 spectrometers using TMS as internal standard. Mass spectra were taken on HP5989A, and VG Quattro MS/MS spectrometers. TLC was performed using silica gel plates F254 (Merk). Column chromatography was performed using silica gel, particle size 10-40 μm , purchased from Qindao Ocean Chemical Factory.

Typical Procedure. To a suspension of peracetylated chitobiose **1** (2.0 g, 2.96 mmol) in dry $\text{ClCH}_2\text{CH}_2\text{Cl}$ (40 mL), was added TMSOTf (0.63 mL, 3.26 mmol). After the mixture was stirred at 50 °C overnight, the resulting solution was cooled to room temperature under N_2 and dry 4Å MS (1 g) was added. The mixture was stirred for 30 min at room temperature and then alcohol (3.0 equiv) was added. After the mixture was stirred for another 2 days, it was neutralized with Et_3N , then filtered. The solid was washed with CH_2Cl_2 -MeOH (10:1) several times. The filtrate and washings were combined, and concentrated. The residue was chromatographed on silica gel with CH_2Cl_2 /MeOH (50:1 v/v) as the eluent to give the product (75%-95%).

Methyl 2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy- β -D-glucopyranosyl)- β -D-glucopyranoside (2). R_f 0.36 (10:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$); mp 282–284 °C (from MeOH); $[\alpha]_D^{22}$ -55.9° (c 0.30, CHCl_3) [Lit.⁸ mp 284 °C (from MeOH), $[\alpha]_D^{22}$ -44° (c 0.55, CHCl_3)]; ^1H NMR (CDCl_3) δ 5.99 (d, 1H, $J=9.0$ Hz, N'-H), 5.74 (d, 1H, $J=9.5$ Hz, N-H), 5.19 (t, 1H, $J=9.5$ Hz, H-3'), 5.11 (t, 1H, $J=8.3$ Hz, H-3), 5.06 (t, 1H, $J=9.6$ Hz, H-4'), 4.56 (d, 1H, $J=8.4$ Hz, H-1'), 4.40–4.35 (m, 3H), 4.06 (m, 3H), 3.90 (dd, 1H, $J=8.7, 9.4$ Hz, H-2'), 3.74 (t, 1H, $J=8.5$ Hz, H-4), 3.64 (m, 2H), 3.47 (s, 3H, -OMe), 2.14–1.95 (m, 21H, 7 \times Ac); EI-MS: m/z 649 (0.8, M^+), 617 (0.5, M^+ -OMe), 330 (51), 288 (12).

Allyl 2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy- β -D-glucopyranosyl)- β -D-glucopyranoside (3). R_f 0.40 (10:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$); mp 253–254 °C (from EtOH); $[\alpha]_D^{22}$ -56.3° (c 0.38 CHCl_3) [Lit.⁸ mp 254 °C (from EtOH), $[\alpha]_D^{22}$ -42° (c 0.43, CHCl_3)]; ^1H NMR (CDCl_3) δ 6.04 (d, 1H, $J=9.0$ Hz, N'-H), 5.85 (m, 1H, $-\text{CH}_2\text{CH}=\text{CH}_2$), 5.78 (d, 1H, $J=9.2$ Hz, N-H), 5.26 (m, 2H, $-\text{CH}_2\text{CH}=\text{CH}_2$), 5.19 (t, 1H, $J=9.5$ Hz, H-3'), 5.16 (t, 1H, $J=9.1$ Hz, H-3), 5.06 (t, 1H, $J=9.8$ Hz, H-4'), 4.57 (d, 1H, $J=8.3$ Hz, H-1'), 4.51 (d, 1H, $J=7.6$ Hz, H-1), 4.40–4.22 (m, 4H), 4.05 (m, 3H), 3.89 (dd, 1H, $J=8.7, 9.3$ Hz, H-2'), 3.74 (t, 1H, $J=8.2$ Hz, H-4), 3.65 (m, 2H), 2.14–1.95 (m, 21H, 7 \times Ac); EI-MS: m/z 675 (1.6, M^+), 617 (1.2, M^+ - $\text{OCH}_2\text{CH}=\text{CH}_2$), 330 (65), 288 (3).

Benzyl 2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy- β -D-glucopyranosyl)- β -D-glucopyranoside (4). R_f 0.44 (10:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$); mp 264–265 °C (from MeOH); $[\alpha]_D^{22}$ -46.3° (c 0.24, CHCl_3); IR (KBr) 3350, 1740 (OAc), 1660 (NAc), 1540, 1360, 1230, 1040; ^1H NMR (CDCl_3) δ 7.30 (m, 5H, -OBn), 6.21 (d, 1H, $J=7.8$ Hz, N'-H), 5.79 (d, 1H, $J=7.9$ Hz, N-H), 5.18 (t, 1H, $J=9.5$ Hz, H-3'), 5.04 (m, 2H), 4.87 (d, 1H, $J=11.9$ Hz, $-\text{OCH}_2\text{Ph}$), 4.59–4.34 (m, 6H), 4.12 (m, 1H), 4.03 (d, 1H, $J=12.4$ Hz, $-\text{OCH}_2\text{Ph}$), 3.90 (dd, 1H, $J=8.5, 9.3$ Hz, H-2'), 3.75 (m, 1H, H-4), 3.64 (m, 2H), 2.16–1.93 (m, 21H, 7 \times Ac); EI-MS: m/z 726 (0.5, M^++1), 617 (0.4, M^+ -OBn), 330 (57), 288 (17).

Anal. Calcd for $\text{C}_{33}\text{H}_{44}\text{N}_2\text{O}_{16}$: C, 54.68; H, 6.13; N, 3.87. Found: C, 54.35; H, 5.86; N, 3.60.

Isopropyl 2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy-β-D-glucopyranosyl)-β-D-glucopyranoside (5). R_f 0.41 (10:1 CH₂Cl₂/MeOH); [α]_D²² -41.4° (*c* 0.35, CH₂Cl₂); IR (KBr) 3300 (N-H), 1750 (OAc), 1680 (NAc), 1540, 1370, 1230, 1040; ¹H NMR (CDCl₃+CD₃OD) δ 5.17 (dd, 1H, *J*=9.4, 10.3 Hz, H-3'), 5.08 (dd, 1H, *J*=8.6, 10.3 Hz, H-3), 5.00 (t, 1H, *J*=9.5 Hz, H-4'), 4.56 (d, 1H, *J*=8.3 Hz, H-1'), 4.54 (d, 1H, *J*=8.2 Hz, H-1), 4.03 (m, 2H), 4.12 (q, 1H, *J*=5.8 Hz), 4.03 (dd, 1H, *J*=2.0, 12.2 Hz), 3.87 (m, 2H), 3.67 (m, 3H), 3.38 (m, 1H, -OCHMe₂), 2.11-2.00 (m, 15H, 5×OAc), 1.92, 1.91 (2s, 6H, 2×NAc), 1.20 (d, 3H, *J*=6.2 Hz, -OCHMe₂), 1.13 (d, 3H, *J*=6.1 Hz, -OCHMe₂); FAB-MS: *m/z* 677 (M⁺+1), 618, 330, 288.

***tert*-Butyl 2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy-β-D-glucopyranosyl)-β-D-glucopyranoside (6).** R_f 0.42 (10:1 CH₂Cl₂/MeOH); mp 258-259 °C(dec.) (from MeOH); [α]_D²² -24.8° (*c* 0.5, CH₂Cl₂); IR (KBr) 3300(N-H), 1740 (OAc), 1660 (NAc), 1360, 1220, 1040; ¹H NMR (CDCl₃) δ 6.04 (d, 1H, *J*=7.4 Hz, N'-H), 5.66 (broad, 1H, N-H), 5.28 (m, 2H), 5.03 (t, 1H, *J*=9.2 Hz), 4.72 (d, 1H, *J*=8.8 Hz, H-1'), 4.67 (d, 1H, *J*=7.5 Hz, H-1), 4.37 (m, 2H), 4.22 (m, 1H), 4.05 (d, 1H, *J*=12.2 Hz), 3.68 (m, 5H), 2.10-2.00 (m, 15H, 5×OAc), 1.94 (s, 6H, 2×NAc), 1.21 (s, 9H, -C(CH₃)₃); FAB-MS: *m/z* 691 (M⁺+1), 618, 330, 288, 228.

6-*O*-[2-Acetamido-3,6-di-*O*-acetyl-2-deoxy-4-*O*-(2-acetamido-3,4,6-tri-*O*-acetyl-2-deoxy-β-D-glucopyranosyl)-β-D-glucopyranosyl]-1,2:3,4-di-*O*-isopropylidene-α-D-galactopyranose (7). R_f 0.48 (10:1 CH₂Cl₂/MeOH); [α]_D²² -63.7° (*c* 0.42, CHCl₃); IR (KBr) 3350, 1740 (OAc), 1660 (NAc), 1360, 1220, 1050; ¹H NMR (CDCl₃) δ 5.97 (d, 1H, *J*=8.5 Hz, N'-H), 5.76 (d, 1H, *J*=8.8 Hz, N-H), 5.33 (d, 1H, *J*=4.9 Hz, H-1), 5.10 (m, 3H), 4.61-4.35 (m, 7H), 4.15-3.97 (m, 6H), 3.68 (m, 4H), 2.13-1.95 (m, 21H, 7×Ac), 1.50, 1.44, 1.32, 1.31 (4s, 12H, 2×isopropylidene); FAB-MS: *m/z* 877 (1.0, M⁺), 617 (1.5), 330 (10), 288 (20).

ACKNOWLEDGMENTS

The authors are grateful to the *National Climbing Program of China* for financial support of this work.

REFERENCES AND NOTES

1. J. Montreuil, *Adv. Carbohydr. Chem. Biochem.*, **37**, 157 (1980).
2. a) C. Merser, P. Sinay, A. Adam, *Biochem. Biophys. Res. Commun.*, **66**, 1316 (1975); b) S. Kusumoto, Y. Tarumi, K. Ikenaka, T. Shiba, *Bull. Chem. Soc. Jpn.*, **49**, 533 (1976).
3. A. Darvill, C. Augur, C. Bergmann, R. W. Carlson, J. -J. Cheong, S. Eberhard, M. G. Hahn, V.-M. Ló, V. Marfà, B. Meyer, D. Mohnen, M. A. O'Neill, M. D. Spiro, H. van Halbeek, W. S. York and P. Albersheim, *Glycobiology*, **2**, 181 (1992).
4. a) P. Lerouge, P. Roche, C. Faucher, F. Maillet, G. Truchet, J. C. Prome, J. Denarie, *Nature*, **344**, 781 (1990); b) R. F. Fisher, S. R. Long, *Nature*, **357**, 655 (1992).
5. a) K. Suzuki, T. Mikami, Y. Okawa, A. Tokoro, S. Suzuki, M. Suzuki, *Carbohydr. Res.*, **151**, 403 (1986); b) V. I. Gorbach, I. N. Krasikova, P. A. Luk'yanov, Y. N. Loenko, T. F. Solov'eva, Y. S. Ovodov, V. V. Deev, A. A. Pimenov, *ibid.*, **260**, 73 (1994).
6. a) H. Kuyama, Y. Nakahara, T. Nukada, Y. Ito, Y. Nakahara, T. Ogawa, *ibid.*, **243**, C1 (1993); b) C. Lamberth, J. O. Nagy, C. Kasper, M. D. Bednarski, *J. Carbohydr. Chem.*, **13**, 819 (1994).
7. S.-I. Nishimura, H. Kuzuhara, *Carbohydr. Res.*, **206**, 207 (1990).
8. S.-I. Nishimura, H. Kuzuhara, Y. Takiguchi, K. Shimahara, *ibid.*, **194**, 223 (1989).
9. C. Bosso, J. Defaye, A. Domard, A. Gadelle, C. Pedersen, *ibid.*, **156**, 57 (1986).
10. J. Banoub, P. Boullanger, D. Lafont, *Chem. Rev.*, **92**, 1167 (1992).
11. a) T. Ogawa, K. Beppu, S. Nakabayashi, *Carbohydr. Res.*, **93**, C6 (1981); b) M. Hollosi, E. Kollat, I. Laczko, K. F. Medzihradszky, J. Thurin, L. Otvos, Jr., *Tetrahedron Lett.*, **32**, 1531 (1991); c) P. C. Boldt, M. H. M. G. Schumacher-Wandersleb, M. G. Peter, *ibid.*, **32**, 1413 (1991).
12. J. Dahman, G. Guosspeilius, A. -C. Larsson, T. Lave, G. Noorl, K. Palsson, *Carbohydr. Res.*, **138**, 17 (1985).
13. Peracetylated chitobiose oxazoline was prepared by treatment of peracetylated chitobiose **1** with TMSOTf, as described by H. Kuzuhara et al.⁸. After the reaction was completed, the mixture was poured into ice-cold aqueous sodium hydrogen-carbonate, extracted with CH₂Cl₂. The CH₂Cl₂ layer was washed with water, dried, and concentrated. The residue was chromatographed on silica gel with 50:1(v/v) CH₂Cl₂-MeOH as the eluent, yield 95%.
14. M. Kiso, L. Anderson, *Carbohydr. Res.*, **72**, C12, C15 (1979).
15. G. Arsequell, L. Krippner, R. A. Dwek, S. Y. C. Wong, *J. Chem. Soc., Chem. Commun.*, 2383 (1994).