

## Carbon–Carbon Bond Formation promoted by Electrophiles and the Mechanism of the Fischer–Tropsch Reaction

Andrew Nutton, Amelio Vázquez de Miguel, Kiyoshi Isobe, and Peter M. Maitlis\*

Department of Chemistry, The University, Sheffield S3 7HF, U.K.

Decomposition of *trans*-[(C<sub>5</sub>Me<sub>5</sub>RhMe)<sub>2</sub>(μ-CH<sub>2</sub>)<sub>2</sub>] takes place at 20 °C in the presence of 1-electron oxidisers to give methane, ethylene, and notably propylene, by coupling of C<sub>1</sub> ligands (thermally the same reactions only occur at 250 °C); the analogy to Fischer–Tropsch reactions on rhodium surfaces is noted.

The promotion of C–C coupling reactions, in addition to the formation of R–E, by the action of electrophiles (E<sup>+</sup>) on alkyl metal complexes has been known for some time.<sup>1</sup> Kochi and others have pointed out that for a number of complexes of the type [L<sub>n</sub>MR<sub>2</sub>], (M = Fe<sup>II</sup>, Ni<sup>II</sup>, Co<sup>III</sup>, *etc.*) reductive elimination to give R–R proceeds most easily if a one-electron oxidation occurs first; this step can be induced by a large number of electrophiles.<sup>2</sup>

We now report, (i) that C–C coupling is promoted by electrophiles even in much higher oxidation state complexes, (ii) that electrophiles will also promote the coupling of three C<sub>1</sub> ligands on a metal complex, and (iii) on the relevance of these observations to current models for Fischer–Tropsch reactions on metal surfaces.


The Ir<sup>V</sup> complex C<sub>3</sub>Me<sub>5</sub>IrMe<sub>4</sub><sup>3</sup> is rather stable; it decomposes on pyrolysis, yielding methane (≥99%), but temperatures of >200 °C are needed to get useful rates and high degrees of decomposition. By contrast, Na<sub>2</sub>IrCl<sub>6</sub> in MeCN causes a fast (stoichiometric) reaction even at 20 °C that gives ethylene, ethane, and methane in a 6:6:88 ratio, together with some methyl chloride. Although the reaction is

complex some electrophile promoted coupling has clearly occurred.

More informative are the reactions of the *trans*-dimethyldi-μ-methylenedirhodium(IV) complex (1).<sup>4,5</sup> With iodine (and bromine) a two-stage reaction occurs, yielding methyl halide and the dihalogeno-di-μ-methylene complex (2),<sup>6</sup> which reacts with more X<sub>2</sub> to eliminate ethylene and give (3) (Scheme 1).

Acids react in a similar two-step process. Methane is of course the main product but some ethylene and ethane, and, particularly notably, propylene are also obtained (Table 1). The proportion of propylene (and of ethylene and ethane) increases when other (oxidizing) electrophiles are used. The reaction with Na<sub>2</sub>IrCl<sub>6</sub> in MeCN is especially noteworthy since it gives a very similar distribution of C<sub>1</sub>, C<sub>2</sub>, and C<sub>3</sub> products to those obtained by pyrolysing (1).<sup>†</sup> However, in

<sup>†</sup> Propylene only becomes a significant product in the thermal decomposition at *higher* temperatures suggesting that there is a substantial activation barrier towards the 3 × C<sub>1</sub> coupling process.



Scheme 1

**Table 1.** Products from the decomposition of [(C<sub>5</sub>Me<sub>5</sub>RhCH<sub>3</sub>)<sub>2</sub>(μ-CH<sub>2</sub>)<sub>2</sub>], (1).

Conditions <sup>a</sup>	μmol of products <sup>b</sup>				Decomp. <sup>c</sup> %
	CH <sub>4</sub>	C <sub>2</sub> H <sub>4</sub>	C <sub>2</sub> H <sub>6</sub>	C <sub>3</sub> H <sub>6</sub>	
HBf <sub>4</sub> -C <sub>6</sub> H <sub>6</sub>	26.4	3.4	0.9	0.5	49
AgBF <sub>4</sub> -C <sub>6</sub> H <sub>6</sub>	11.6	6.4	3.4	0.7	45
AgBF <sub>4</sub> -MeCN	12.7	2.0	12.6	1.3	61
(NH <sub>4</sub> ) <sub>4</sub> Ce(SO <sub>4</sub> ) <sub>4</sub> ·2H <sub>2</sub> O-MeCN	12.8	6.4	0.6	3.2	41
Na <sub>2</sub> IrCl <sub>6</sub> ·2H <sub>2</sub> O-MeCN <sup>d</sup>	20.6	3.0	0.5	4.6	57
Pyrolysis at 350 °C <sup>e</sup>	19.6	4.8	0.2	5.2	61

<sup>a</sup> Samples of (1) (10 mg) were treated with 5–6 equiv. of the named reagent in *ca.* 1 cm<sup>3</sup> solvent in 3 cm<sup>3</sup> sealed flasks at 20 °C. Reaction was instantaneous in most cases (as shown by colour changes or deposition of metal) and was completed by briefly heating to 50 °C. *cis*-(1) behaved similarly. <sup>b</sup> Analysed by g.c. on Poropak Q at 100 °C. The identities of the gases were checked by mass spectrometry which also showed no significant incorporation of deuterium from either (1) labelled on the C<sub>5</sub>Me<sub>5</sub> ring or from CD<sub>3</sub>CN solvent. <sup>c</sup> Calculation based upon the theoretical maximum amount of decomposition from 18.7 μmol of (1) being 4 × 18.7 (*i.e.* 74.8) μmol equiv. of C<sub>1</sub>. <sup>d</sup> Small amounts of methyl chloride and propane were also detected. <sup>e</sup> A sample of 25 μg of complex was pyrolysed in the inlet of a Carlo Erba gas chromatograph at 350 °C and the gases were analysed. The numbers of μmol of products given have been scaled up to allow direct comparison. A small amount of propane was also detected.

Several decomposition and coupling reactions are occurring simultaneously, the significant one of which is the formation of propylene. One possible mode by which this occurs is a linking of one CH<sub>3</sub> and the two CH<sub>2</sub> ligands in an 'oxidized' form of the complex to give a C<sub>3</sub> ligand which β-eliminates propylene.<sup>†‡§</sup>

We have already noted<sup>4</sup> the appropriateness of complex (1) (and its *cis*-isomer) as models for the methylene polymerization stage in one mechanism for the Fischer-Tropsch reaction on metal surfaces.<sup>7</sup> One intriguing feature of the heterogeneous Fischer-Tropsch reactions is the large but poorly understood influence of promoters. The results presented here suggest that in the heterogeneous Fischer-Tropsch reaction the surface on which the metal is supported may play a hitherto unsuspected role as an electron acceptor in the polymerization stage. This is in addition to the roles already noted where the surface can act as Lewis acid and oxygen acceptor in the CO activation and reduction steps.<sup>8</sup> Particular point is given to this observation since a number of workers have noted that certain specific rhodium plus surface oxide combinations lead to enhanced rates of methanation of CO-H<sub>2</sub> mixtures and, more interestingly, to the enhanced formation of higher hydrocarbons and oxygenates when compared to rhodium metal alone.<sup>9</sup> It has been suggested that these effects are due to 'strong metal-support interactions'<sup>10</sup> in which the (oxide) support actually acts as an oxidant of the metal. Although the heterogeneous metal catalyst system is very much more complex and has many more steps than the rather simple model offered by complex (1), the similarity of behaviour is encouraging.

We thank B.P. Chemicals Ltd, the S.E.R.C., and the Spanish Ministry of Education for support and Mr. P. R. Ashton and Mr. D. G. Andrews for technical assistance.

Received, 18th November 1982; Com. 1318

## References

- See, for example, U. Krüerke, C. Hoogzand, and W. Hübel, *Chem. Ber.*, 1961, **94**, 2817; U. Krüerke and W. Hübel, *ibid.*, p. 2829.
- Summarized in J. K. Kochi, 'Organometallic Mechanisms and Catalysis,' Academic Press, New York, 1978, pp. 282, 351–352.
- K. Isobe, P. M. Bailey, and P. M. Maitlis, *J. Chem. Soc., Chem. Commun.*, 1981, 808.
- K. Isobe, D. G. Andrews, B. E. Mann, and P. M. Maitlis, *J. Chem. Soc., Chem. Commun.*, 1981, 810.
- A. Vázquez de Miguel, K. Isobe, B. F. Taylor, A. Nutton, and P. M. Maitlis, *J. Chem. Soc., Chem. Commun.*, 1982, 758.
- K. Isobe, P. M. Bailey, P. Schofield, J. T. Gauntlett, A. Nutton, and P. M. Maitlis, *J. Chem. Soc., Chem. Commun.*, 1982, 425.
- R. C. Brady and R. Pettit, *J. Am. Chem. Soc.*, 1980, **102**, 6181; *ibid.*, 1981, **103**, 1287 and references therein.
- D. F. Shriver, *Am. Chem. Soc. Symp. Ser.*, 1981, **152**, 1.
- See, for example, M. A. Vannice, *J. Catal.*, 1982, **74**, 199 and references therein; P. R. Watson and G. A. Somorjai, *ibid.*, 1981, **72**, 347; F. Solymosi, I. Tombacz, and M. Kocsis, *ibid.*, 1982, **75**, 78.
- S. J. Tauster, S. C. Fung, and R. L. Garten, *J. Am. Chem. Soc.*, 1978, **100**, 170; S. J. Tauster, S. C. Fung, R. T. K. Baker, and J. A. Horsley, *Science*, 1981, **211**, 1121.

<sup>†</sup> Free ethylene does not appear to be intermediate; a reaction carried out in the presence of a four-fold excess of ethylene only showed a very small increase in propylene formed.

<sup>§</sup> One major organometallic product from such reactions of (1) in MeCN is [C<sub>5</sub>Me<sub>5</sub>Rh(MeCN)<sub>3</sub>]<sup>2+</sup> suggesting that a part of the total reaction may be given by [(C<sub>5</sub>Me<sub>5</sub>RhCH<sub>3</sub>)<sub>2</sub>(μ-CH<sub>2</sub>)<sub>2</sub>] + 6MeCN – 4e<sup>–</sup> → 2[C<sub>5</sub>Me<sub>5</sub>Rh(MeCN)<sub>3</sub>]<sup>2+</sup> + CH<sub>3</sub>CH=CH<sub>2</sub> + CH<sub>4</sub>.

contrast with the pyrolysis, which needs 350 °C, the reactions with IrCl<sub>6</sub><sup>2–</sup>, and those with Ag<sup>+</sup> or Ce<sup>4+</sup>, proceeded rapidly even under ambient conditions. Mass spectrometric analyses of the gases indicated no significant transfer of deuterium to the C<sub>1</sub>, C<sub>2</sub>, or C<sub>3</sub> products when complex (1) labelled only on the C<sub>5</sub>Me<sub>5</sub> ring decomposed under these conditions.