LETTERS

Exploiting a Difference in Leaving Group Ability: An Approach to β -Substituted Monofluoroalkenes Using *gem*-Chlorofluoropropenes

Jean-Denys Hamel, Mélissa Cloutier, and Jean-François Paquin*

CCVC, PROTEO, Département de chimie, Université Laval, 1045 Avenue de la Médecine, Québec, Québec, Canada G1V 0A6

(5) Supporting Information

ABSTRACT: The superior nucleofuge character of chlorine over fluorine was taken advantage of in the selective $S_N 2'$ substitution reaction of *gem*-chlorofluoropropenes, allowing for the clean formation of β -substituted monofluoroalkenes under metal-free conditions. Numerous *N*-, *S*-, *O*-, and *C*-nucleophiles behaved nicely in this system. Further synthetic transformations of selected monofluoroalkenes were also accomplished.



O rganofluorine compounds occupy a significant place in medicinal chemistry, agrochemistry, and material sciences due to the unique properties of the fluorine atom.¹ It is, therefore, not surprising that considerable effort has been dedicated to the discovery and improvement of methods for the preparation of fluorinated compounds.² Over the past years, we have been particularly interested in the synthesis of monofluoroalkenes,³ a useful enol or amide mimic, from 3,3-difluoropropenes⁴ using either transition-metal-catalyzed (Scheme 1, eq 1)^{5,6} or metal-free (Scheme 1, eq 2)^{7,8} reactions. All of these approaches relied on using one of the fluorine atoms as a leaving group, even though it is generally considered a poor one.^{1,9} While these methods provide a convenient access

Scheme 1. Previous Work and Current Work

Previous work from 3,3-difluoropropenes

Metal-catalyzed amination^{5,6}

$$F = Pd \text{ or } Pt$$

$$R = alkyl$$

$$R = alkyl$$

$$F = NR_2$$

$$R = alkyl$$

$$R = alky + a$$

Metal-free S_N2' substitution reactions using highly basic nucleophiles^{7,8}



This work

Metal-free S_N2' substitution reactions with various milder nucleophiles

gem-chlorofluoropropenes monofluoroalkenes

- Nu = Aliphatic and aromatic amines
 - Phenolates
 - Aliphatic and aromatic thiols
 - CyanideDimethyl malonate
 - Carboxylates and thiocarboxylates
 - Alkyl- and arylcuprates

to this important fluorinated motif, they suffer from drawbacks. For instance, the Pd-catalyzed reaction only allowed the use of aliphatic amines (Scheme 1, eq 1, [M] = Pd).⁵ Likewise, the Pt-catalyzed reaction only tolerated secondary aliphatic amines, and other nucleophiles behaved poorly (Scheme 1, eq 1, [M] = Pt).⁶ Concerning the metal-free approach (Scheme 1, eq 2), only organolithium reagents, lithium thiolates, and lithium anilides performed well. In this case, the high basicity of those reagents considerably limits the functional group tolerance. Considering those issues, we decided to explore a milder and more versatile alternative that would not only solve the concerns raised but also extend the scope in terms of nucleophiles, thus allowing the possibility to prepare more functionalized β -substituted monofluoroalkenes.

To this end, we sought to exploit the difference in leaving group ability between a fluorine and a chlorine in geminal relationship, ^{1b,10} a phenomenon that has been rarely taken advantage of.^{11–19} Herein, we report a convenient approach to β -substituted monofluoroalkenes from *gem*-chlorofluoropropenes exploiting this difference in nucleofuge character (Scheme 1, eq 3).²⁰ Notably, this transformation provides structurally diverse monofluoroalkenes and is compatible with a wide range of milder nucleophiles, some of which could not be employed using previous methods.

The reactivity was initially investigated using chlorofluoropropene **1**, which was easily obtained from 1-tetralone in three steps (α -fluorination using Selectfluor, α -chlorination using *N*chlorosuccinimide, and Peterson olefination),²¹ and morpholine as the nucleophile. After some optimization, it was found that the desired β -aminomonofluoroalkene **2a** could be obtained in 96% isolated yield (Scheme 2). This result compared favorably with the 72% and 68% yields achieved in the Pd- and Pt-catalyzed transformations, respectively (i.e., Scheme 1, eq 1).^{5a,6}

Delighted with this initial result, we began exploring the use of other nucleophiles and found that a variety could be employed successfully upon slight modifications of reaction

Received: March 1, 2016

Scheme 2. Synthesis and Reactivity of Chlorofluoropropene 1 with Morpholine



conditions (Table 1). For instance, the use of sodium phenolate provided the monofluoroalkene **2b** in 80% yield (Table 1, entry

Table 1. Reactivity of Chlorofluoropropene 1 with Various Nucleophiles^a





1). Notably, a phenolic nucleophile did not work with 3.3difluoropropenes (i.e., Scheme 1, eq 2),⁸ where an elimination side reaction was predominant. Sulfur-based nucleophiles, such as an alkyl- and an arylthiolate, as well as a thiocarboxylate, all behaved nicely (Table 1, entries 2-4), and the desired products 2c-e could be obtained in very good to excellent yields.²² The use of cyanide (Table 1, entry 5) as the nucleophile provided the β -cyanomonofluoroalkene 2f in 96% yield. Interestingly, the nitrile functionality opens the door to further transformations (vide infra). Dimethyl malonate reacted well, and the monofluoroalkene 2g was produced in 82% yield (Table 1, entry 6). Phenyl cuprate (generated in situ from the corresponding Grignard reagent and copper(I) bromide dimethyl sulfide complex)²³ was a suitable nucleophile, and the monofluoroalkene 2h was obtained in 95% yield, which compares favorably with the 80% obtained using highly basic phenyllithium with a 3,3-difluoropropene (i.e., Scheme 1, eq 2).7 At that point, despite extensive optimization, the use of either sodium methoxide (an alkoxide), sodium acetate (a carboxylate), or N-methylaniline (an aromatic amine) did not provide the desired product in satisfactory yields (not shown in Table 1). Finally, iodide could be used successfully as a

nucleophile to provide the β -iodomonofluoroalkene **2i** in good NMR yield (Table 1, entry 8). While the isolation of the product did not prove possible, its clean formation, as shown by NMR, provided an opportunity to use it as an intermediate to promote substitution with reluctant nucleophiles (Scheme 3).

Scheme 3. Reaction of 1 with (a) *N*-Methylaniline and (b) Sodium Acetate in the Presence of a Catalytic Amount of Iodide



Indeed, using either *N*-methylaniline or sodium acetate in the presence of a catalytic amount of tetrabutylammonium iodide (TBAI) provided the desired monofluoroalkenes **2j** and **2k** in 91% and 86% yield, respectively, thus further expanding the scope to carboxylates and aromatic amines as suitable nucleophiles for this transformation.

The reactivity of analogues of successful N-, S-, O-, and Cnucleophiles was next assessed (Scheme 4). Functionalized secondary amines such as L-proline methyl ester and 2-(methylamino)ethanol reacted well with 1 and provided the desired products 2l and 2m in 85% and 64% yield, respectively. Notably, in the latter case, protection of the alcohol was not necessary. An excellent yield of 2n was obtained with nbutylamine (a primary amine) from 1 as long as an excess of the nucleophile was used to avoid a mixture of mono- and dialkylated products. Interestingly, L-cysteine methyl ester reacted with 1 and provided the S-alkylation product 20 in 70% yield without any need for the protection of the nitrogen. Using estrone as a source of a functionalized phenolate provided product 2p in 78% yield. Other chlorofluoropropenes could also be utilized. For instance, the chlorofluoropropene derived from 1-indanone 4 reacted with an aliphatic amine, Nmethylbenzylamine, to provide monofluoroalkene 9 in 78% yield. Similarly, when the chlorofluoropropene obtained from 4-chromanone 6 was used, reaction with indoline, an aromatic amine, afforded monofluoroalkene 10 in 61% yield. In addition, monofluoroalkene 11 was isolated in 61% yield after exposure of the chlorofluoropropene prepared from 4,4-dimethyl-2cyclohexen-1-one (6) to sodium thiophenolate. Finally, acyclic chlorofluoropropenes could also be used, as exemplified by the transformations of chlorofluoropropenes 7 and 8 into monofluoroalkenes 12a,b and 13a,b, respectively, which were obtained in good to excellent yields as a mixture of diastereoisomers. Interestingly, when 7 was used with either morpholine 12a or sodium thiophenolate 12b as the nucleophile, the E/Z selectivity could be modestly modulated by changing the solvent. In both cases, the use of DMF, a more polar solvent, favored the E-isomer as the major one, while the use of a less polar solvent (i.e., toluene or THF) produced the opposite Z-isomer as the major isomer.^{24,25} The reason behind the change in selectivity is not clear at the moment, and further studies to understand this are in progress.



Scheme 4. Reaction of Cyclic and Acylic Chlorofluoropropenes with Various N-, S-, O-, and C-Nucleophiles^{a,b}

^{*a*}See the Supporting Information for the detailed experimental procedures. ^{*b*}Isolated yield. ^{*c*}5 equiv of the nucleophile was used. ^{*d*}10 equiv of the nucleophile was used. ^{*e*}Isolated as a mixture of stereoisomers; E/Z ratio estimated by ¹H or ¹⁹F NMR.

Finally, to further extend the utility of this new approach, we investigated potential transformations of the monofluoroalkenes generated (Scheme 5). For instance, reduction of the thioester moiety of 2c with LiAlH₄ provided an excellent yield of β -mercaptomonofluoroalkene 14.²² Hydrolysis of nitrile 2f under basic conditions gave acid 15 in 74% yield. As monofluoroalkenes can be seen as amide mimics,³ this particular product resembles an amide bond between a glycine and an N-substituted amino acid. Likewise, a two-step decarboalkoxylation²⁵/hydrolysis sequence of malonate 2g provided acid 16, which in this case, corresponds to an amide bond between a β -glycine and an N-substituted amino acid. Finally, deprotection of the acetate of 2k through methanolysis led to the alcohol $17^{22c,26}_{,,26}$ which could be methylated to provide methyl ether 18. Thus, even if, at this time, preparation of 18 directly from 1 using methoxide as the nucleophile is not possible, this short and high-yielding three-step procedure provides an alternative route to β -alkyloxymonofluoroalkenes.

In conclusion, we have reported the preparation of β substituted monofluorolalkenes through allylic substitution of *gem*-chlorofluoropropenes. This strategy relies on the difference in leaving group ability between chlorine and fluorine atoms bonded to the same carbon atom. As neither a transition metal nor a highly basic nucleophile is required in this reaction, a broader range of nucleophiles was shown to be compatible as compared to our previous work. Overall, this new approach allows the preparation of structurally diverse monofluoroalScheme 5. Synthetic Transformation of Some Monofluoroalkenes Obtained



kenes, potentially useful fluorinated building blocks for medicinal chemistry efforts, in good to excellent yields.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.6b00590.

Detailed experimental procedures and full spectroscopic data for all new compounds (PDF)

Crystallographic data of (E)-12a·HCl (CIF)

AUTHOR INFORMATION

Corresponding Author

* E-mail: jean-francois.paquin@chm.ulaval.ca.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the Natural Sciences and Engineering Research Council of Canada (NSERC), the FRQNT Centre in Green Chemistry and Catalysis (CCVC), the FRQNT Network for Research on Protein Function, Engineering, and Applications (PROTEO), and the Université Laval. J.D.H. thanks NSERC for a Vanier Canada Graduate Scholarship.

REFERENCES

(1) (a) Hiyama, T. In Organofluorine Compounds; Chemistry and Applications; Yamamoto, H., Ed.; Springer: New York, 2000 and references cited therein. (b) O'Hagan, D. Chem. Soc. Rev. 2008, 37, 308.

(2) For selected recent reviews on the subject, see (a) Liang, T.; Neumann, C.; Ritter, T. Angew. Chem., Int. Ed. 2013, 52, 8214.
(b) Campbell, M. G.; Ritter, T. Chem. Rev. 2015, 115, 612.
(c) Charpentier, J.; Früh, N.; Togni, A. Chem. Rev. 2015, 115, 650.
(d) Xu, X.-H.; Matsuzaki, K.; Shibata, N. Chem. Rev. 2015, 115, 731.
(e) Ni, C.; Hu, M.; Hu, J. Chem. Rev. 2015, 115, 765. (f) Yang, X.; Wu, T.; Phipps, R. J.; Toste, F. D. Chem. Rev. 2015, 115, 826. (g) Savoie, P.-R.; Welch, J. T. Chem. Rev. 2015, 115, 1130. (h) Champagne, P. A.; Desroches, J.; Hamel, J.-D.; Vandamme, M.; Paquin, J.-F. Chem. Rev. 2015, 115, 9073.

(3) Reviews: (a) Landelle, G.; Bergeron, M.; Turcotte-Savard, M.-O.; Paquin, J.-F. Chem. Soc. Rev. 2011, 40, 2867. (b) Yanai, H.; Taguchi, T. Eur. J. Org. Chem. 2011, 2011, 5939. (c) Hara, S. Top. Curr. Chem. 2012, 327, 59.

(4) Drouin, M.; Hamel, J.-D.; Paquin, J.-F. Synlett 2016, 27, 821.

(5) (a) Pigeon, X.; Bergeron, M.; Barabé, F.; Dubé, P.; Frost, H. N.; Paquin, J.-F. Angew. Chem., Int. Ed. **2010**, 49, 1123. (b) Paquin, J.-F. Synlett **2011**, 2011, 289.

(6) Hamel, J.-D.; Drouin, M.; Paquin, J.-F. J. Fluorine Chem. 2015, 174, 81.

(7) Bergeron, M.; Johnson, T.; Paquin, J.-F. Angew. Chem., Int. Ed. 2011, 50, 11112.

(8) Bergeron, M.; Guyader, D.; Paquin, J.-F. Org. Lett. 2012, 14, 5888.

(9) For a review on C-F bond activation, see: Amii, H.; Uneyama, K. *Chem. Rev.* **2009**, *109*, 2119.

(10) Dorwald, F. Z. In Side Reactions in Organic Synthesis: A Guide to Successful Synthesis Design; Wiley-VCH: Weinheim, 2005.

(11) Reactions of polyhalogenated compounds have not been included here.

(12) For the preparation of fluoroallenes from a fluorochloropropargyl moiety, see: Castelhano, A. L.; Krantz, A. J. Am. Chem. Soc. 1987, 109, 3491.

(13) For a Finkelstein reaction on a α -chloro- α -fluoroamide, see: Bailey, P. D.; Boa, A. N.; Crofts, G. A. *Tetrahedron Lett.* **1989**, *30*, 7457.

(14) For the zinc-mediated Barbier-type reaction of fluoropropargyl chloride, see: Xu, B.; Hammond, G. B. J. Org. Chem. 2006, 71, 3518.

(15) For $S_N 2$ reactions of α -chloro- α -fluoroketones and derivatives, see: (a) Shibatomi, K.; Yamamoto, H. *Angew. Chem., Int. Ed.* **2008**, 47, 5796. (b) Shibatomi, K.; Narayama, A.; Soga, Y.; Muto, T.; Iwasa, S. *Org. Lett.* **2011**, *13*, 2944.

(16) For the nickel-catalyzed Negishi cross-coupling using a α -chloro- α -fluoroketone, see: Liang, Y.; Fu, G. C. J. Am. Chem. Soc. **2014**, 136, 5520.

(17) For the nickel-catalyzed Suzuki cross-coupling using 1-chloro-1-fluoroalkanes, see: (a) Jiang, X.; Sakthivel, S.; Kulbitski, K.; Nisnevich, G.; Gandelman, M. J. Am. Chem. Soc. **2014**, 136, 9548. (b) Jiang, X.; Gandelman, M. J. Am. Chem. Soc. **2015**, 137, 2542.

(18) For the preparation of 2-fluorophenols from in situ generated α -chloro- α -fluoroketones using a selective elimination, see: Yasui, N.; Mayne, C. G.; Katzenellenbogen, J. A. Org. Lett. **2015**, 17, 5540.

(19) For an application in ¹⁸F-labeling, see: Verhoog, S.; Pfeifer, L.; Khotavivattana, T.; Calderwood, S.; Collier, T. L.; Wheelhouse, K.; Tredwell, M.; Gouverneur, V. Synlett **2015**, *27*, 25.

(20) To the best of our knowledge, only one example of such a reaction with highly basic nucleophiles has been reported before; however, a mixture of substitution products resulting from the departure of either the fluoride or the chloride was observed; see ref 7. for details.

(21) All of the chlorofluoropropenes were prepared following the same sequence. See the Supporting Information for more details.

(22) (a) McDonald, I. A.; Bey, P. Tetrahedron Lett. 1985, 26, 3807.
(b) Pirrung, M. C.; Rowley, E. G.; Holmes, C. P. J. Org. Chem. 1993, 58, 5683. (c) Song, Y.; Clizbe, L.; Bhakta, C.; Teng, W.; Li, W.; Wu, Y.; Jia, Z. J.; Zhang, P.; Wang, L.; Doughan, B.; Su, T.; Kanter, J.; Woolfrey, J.; Wong, P.; Huang, B.; Tran, K.; Sinha, U.; Park, G.; Reed, A.; Malinowski, J.; Hollenbach, S.; Scarborough, R. M.; Zhu, B.-Y. Bioorg. Med. Chem. Lett. 2002, 12, 1511.

(23) A stoichiometric amount of copper salt was used since 1 decomposed rapidly in the presence of Grignard reagents.

(24) Stereochemistry of **12a** was confirmed by XRD analysis of (*E*)-**12a**·HCl (CCDC 1451457). Stereochemistry of compounds **12b**, **13a**, and **13b** was attributed by examination of the ¹H, ¹⁹F, and ¹³C NMR spectra.

(25) Curran, D. P.; Zhang, Q. Adv. Synth. Catal. 2003, 345, 329.

(26) Shen, Q.; Hong, J. H. Nucleosides, Nucleotides Nucleic Acids 2008, 27, 213.