

Chiral Bisphosphazides as Dual Basic Enantioselective Catalysts

Hiroshi Naka,* Nobuhiko Kanase, Masahiro Ueno, and Yoshinori Kondo*[a]


Abstract: Chiral bisphosphazides complexed with lithium salts efficiently catalyze the direct enantioselective 1,4-addition of dialkyl malonates to acyclic enones. Spectroscopic studies on the stoichiometry of the bisphosphazide and lithium salt have indicated the formation of a 1:1 species as the active enantioselective catalyst. It is suggested that the catalyst generates a complex of the protonated phosphazide and the chiral nucleophile as the key intermediate. The phosphazide moiety appears to be a promising dual basic functionality for stereo- and chemoselective catalytic transformations.

Keywords: asymmetric catalysis • basicity • lithium • Michael addition • phosphazene base

Introduction

Phosphazene compounds are employed as strong neutral organic bases,^[1] functional materials,^[2] and chemoselective synthetic reagents.^[3] The utilization of Schwesinger's phosphazene bases^[1] as strongly basic reagents for the chemoselective deprotonation of acidic and even less acidic C–H protons has been extensively investigated.^[4] More recently, applications in the catalytic regulation of organometallic^[5] and organosilicon^[6] species have been reported. Most of these studies have dealt exclusively with simple iminophosphoranes ($R^1N=PR^2_3$), and several examples of chiral phosphazene-catalyzed enantioselective reactions have been reported.^[4i, 5b, c, 7]

However, little attention has been paid to phosphazides ($R^1N=N=N-PR^2_3$), which are generally considered to be unstable intermediates in the Staudinger reaction,^[8] although they are also potentially attractive as Lewis and/or Brønsted basic functionalities [Eq. (1)].



Verkade et al. recently reported that several azidoproazaphosphatranes resist thermal denitrogenation and may be used as catalytic bases in base-mediated reactions.^[9] It was noted that appropriate steric bulkiness and electronic properties of the substituents were required for sufficient stabilization of the phosphazide intermediates.^[9]

These findings motivated us to investigate the possible reactivity of the phosphazide moiety in base-catalyzed asymmetric reactions. Preparation of chiral phosphazides through the Staudinger reaction is attractive, because simply by mixing chiral azides with bulky and electron-donating tris(dialkylamino)phosphines, chiral phosphazides are generated, which may be used directly for asymmetric reactions without further purification.^[10]

We report here the first enantioselective catalysis using a chiral bisphosphazide (**1a**) complexed with lithium salts, by which direct, enantioselective 1,4-addition of dialkyl malonates to acyclic enones can be efficiently accomplished (Scheme 1). The asymmetric direct Michael addition of malonates to enones is a representative reaction in modern asymmetric catalysis,^[11] and known metallic^[12] and non-metallic^[13, 14] catalysts developed for this purpose are also effective for other base-catalyzed asymmetric processes.^[11] We also present a possible application of the phosphazide

[a] Prof. H. Naka, N. Kanase, Dr. M. Ueno, Prof. Y. Kondo
Graduate School of Pharmaceutical Sciences
Tohoku University
Aobayama, Aoba-ku, Sendai 980-8578 (Japan)
Fax: (+81) 22-795-5917
E-mail: naka@mail.pharm.tohoku.ac.jp
ykondo@mail.pharm.tohoku.ac.jp

Supporting information for this article is available on the WWW under <http://www.chemeurj.org/> or from the author: Preparation of bisazide, chiral HPLC charts for products, and NMR spectra for the titration experiments.

Table 3. Catalysis using other bases.^[a]

Entry	Base	Yield ^[b]	ee ^[c]
1	1a	95	88
2	1b	82	88
3	1c	99	34
4	1d	82	89
5	2	3	1
6	3a	68	12
7	3a ^[d]	92	2
8	–	0	–

[a] Unless otherwise noted, reactions were carried out at -40°C for 48 h in the presence of *trans*-chalcone (1.0 equiv), dimethyl malonate (2.0 equiv), base (10 mol %), and LiClO_4 (10 mol %). [b] Isolated yield of **4a**. [c] Determined by chiral HPLC. [d] Without LiClO_4 .

Next, we tested the conjugate addition of various kinds of nucleophiles to chalcone derivatives (Table 4). The reactions of various dialkyl malonates proceeded in excellent yields with good selectivities (entries 1–3). Di-*tert*-butyl malonate could also be employed when LiOtBu was used in place of LiClO_4 (entry 4). Interestingly, the reaction did not proceed with more acidic 1,3-dicarbonyl compounds such as ketoesters, diketones, or cyclic esters (data not shown). Bisnitrile gave the addition product but with no enantioselectivity (entry 5). Using dimethyl malonate as the nucleophile, the scope of the reaction was investigated in terms of the electrophiles that could be applied (entries 6–14). Chalcone derivatives with both electron-donating (Me, OMe) and electron-withdrawing (Cl, F, NO_2) substituents on their aryl moieties could be successfully used as substrates without altering the enantioselectivity (entries 6–11). Reaction with coor-

dinative substrates, such as those bearing 2-furyl and 2-pyridyl substituents, proceeded without any decrease in selectivity (entries 12 and 13). Substitution of the aryl ring at the α -position was also tolerated (entry 14).

To shed light on the possible structure of the catalyst and the catalytic mechanism, we first examined the ^1H and ^{31}P NMR spectra of complexes of **1a** and LiClO_4 in various stoichiometric ratios (Figure 2). The chemical shift of the benzylic proton changed when 0.5 equiv of LiClO_4 was added to **1a** (Figure 2a), implying the formation of a **1a**:Li chemical species with a molar ratio of 2:1 when less than 1.0 equiv of lithium salt was added. In both the ^1H and ^{31}P NMR spectra, saturation was observed when 1.0 equiv of LiClO_4 was added to **1a**, indicating 1:1 stoichiometric complexation of LiClO_4 and **1a**.




Figure 2. a) ^1H NMR titration of **1a** with LiClO_4 in CD_2Cl_2 (benzylic region) at 25°C . b) ^{31}P NMR titration of **1a** with LiClO_4 in CD_2Cl_2 at 25°C .

Table 4. Bisphosphazide-catalyzed conjugate additions of nucleophilic enols to enones.^[a]

Entry	Substrate	Product	Yield ^[b]	ee ^[c]
1				
2		EWG = COOEt (4b)	> 99	79 (S)
3		= COOiPr (4c)	96	84 (S)
4 ^[d]		= COOBn (4d)	85	80 (S)
5		= COOtBu (4e)	94	71
		= CN (4f)	> 99	1
6				
7	Ar = 4-MeC ₆ H ₄	Ar = 4-MeC ₆ H ₄ (4g)	66	87
8	= 4-MeOC ₆ H ₄	Ar = 4-MeOC ₆ H ₄ (4h)	84	88
9	= 4-ClC ₆ H ₄	Ar = 4-ClC ₆ H ₄ (4i)	86	84
10	= 4-FC ₆ H ₄	Ar = 4-FC ₆ H ₄ (4j)	54	85
11	= 4-NO ₂ C ₆ H ₄	Ar = 4-NO ₂ C ₆ H ₄ (4k)	> 99	88
12	= 2-ClC ₆ H ₄	Ar = 2-ClC ₆ H ₄ (4l)	> 99	89
13	= 2-furyl	Ar = 2-furyl (4m)	41	88
	= 2-pyridyl	Ar = 2-pyridyl (4n)	98	88
14				
	Ar = 4-ClC ₆ H ₄	Ar = 4-ClC ₆ H ₄ (4o)	92	83

[a] Unless otherwise noted, reactions were carried out at -40°C for 48 h in the presence of *trans*-chalcone (1.0 equiv), dimethyl malonate (2.0 equiv), **1a** (10 mol %), and LiClO_4 (10 mol %). [b] Isolated yield. [c] Determined by chiral HPLC. [d] LiOtBu (10 mol %) was used instead of LiClO_4 .

We next investigated the dependence of the yield and ee on the **1a**/ LiClO_4 ratio (Figure 3). In accordance with the ^{31}P and ^1H NMR findings, the selectivity was highest when a 1:1 ratio of **1a** and LiClO_4 was used. A loading of LiClO_4 of less than 1.0 equiv with respect to **1a** resulted in a significant decrease in the yield.

Combined with the above NMR findings, this result could be interpreted as follows. In the presence of 0.5 equiv of Li salt, **1a** forms an inactive 2:1 **1a**/ LiClO_4 species, whereas when 1.0 equiv of Li salt is added to **1a**, an active and stereoselective 1:1 **1a**/ LiClO_4 complex is generated (Scheme 3).

Because the bisphosphazide had to be added to achieve product formation, it presumably deprotonates the dialkyl






Figure 3. Dependence of yield (○) and *ee* (■) on the LiClO₄/1a ratio. The reaction was carried out at −40 °C for 48 h in the presence of 1.0 equiv of chalcone, 2.0 equiv of dimethyl malonate, 10 mol % of 1a, and 0, 3, 5, 7, 9, 10, 12, 15, or 20 mol % of LiClO₄.

malonate to form an enolate nucleophile. The reaction with the bisphosphazide catalyst alone gave poor enantioselectivity; equimolar addition of the lithium salt to the bisphosphazide was necessary for high enantioselectivity. This result suggests that the 1:1 1a/LiClO₄ complex (Scheme 3) is involved in the catalytic cycle. The fact that relatively acidic lithium salts gave better enantioselectivities further implies that the lithium cation complexed with the bisphosphazide plays a Lewis acidic role in the efficient stereoselective reaction. Although details of the reaction



Scheme 4. Proposed catalytic cycle.



Scheme 3. Possible species formed from 1a and LiClO₄ (0.5 or 1.0 equiv).


mechanism of this enantioselective conjugate addition remain unclear, a possible catalytic cycle is shown in Scheme 4.

In the reaction mixture, the 1:1 1a/LiClO₄ complex readily reacts with the dialkyl malonate to form a chiral enolate species and protonated phosphazide. The chiral enolate species reacts with the chalcone, possibly through acid-base dual activation,^[11b] whereby the protonated phosphazene center serves as an acid for the electrophilic activation of the enone, and the bisphosphazide serves as a base for the lithium enolate (Scheme 5a). A putative transition state is presented in Scheme 5b.

Overall, the bisphosphazide serves as a dual base, as a Lewis base for the lithium enolate, and as a Brønsted base for the deprotonation of the dialkyl malonate to form a chiral nucleophile. The resulting protonated bisphosphazide may thus serve as an acid-base dual activator. Further crystallographic, spectroscopic, and computational studies aimed at elucidating the origin of the selectivity are underway.

Conclusion

A novel chiral bisphosphazide (1a) has been developed for the enantioselective catalytic 1,4-addition of malonates to acyclic enones using controlled complexation with lithium salts. Spectroscopic studies on the stoichiometry of the complex formed between the bisphosphazide and the lithium salt suggest the formation of a 1:1 species as the active and selective catalyst. Both the bisphosphazide and the lithium salt proved necessary for obtaining high catalytic activity and enantioselectivity, suggesting that the bisphosphazide serves as a dual base, that is, both as a Lewis base and as a Brønsted base. We suggest that the phosphazide moiety represents a promising dual basic functionality for stereoselec-



Scheme 5. a) Acid-base dual activation in a putative transition state. b) 3D representation of the putative transition state.

tive catalytic transformations. Further studies to extend the scope of the reaction and to improve the enantioselectivity, employing phosphazides with other chiral frameworks, are underway.

Experimental Section

Reactions were carried out under Ar atmosphere using dry solvents. Melting points (m.p.) were determined with a Yazawa micro melting point apparatus and are uncorrected. Infrared (IR) data were recorded on a SensIR ATR (attenuated total reflectance) FT-IR spectrometer. The spectra were each acquired in 32 scans at a resolution of four using the ReactIR 2.20 software system. Absorbance frequencies are reported in cm^{-1} . NMR data were recorded on either a JEOL AL400 spectrometer (395.75 MHz for ^1H , 99.50 MHz for ^{13}C) or a JEOL ECA600 spectrometer (600.172 MHz for ^1H , 150.907 for ^{13}C , 242.956 MHz for ^{31}P). Chemical shifts are expressed in δ values (parts per million, ppm), and coupling constants are expressed in Hertz (Hz). ^1H NMR spectra were referenced to tetramethylsilane as an internal standard or to a solvent signal (CHCl_3 in CDCl_3 : 7.26 ppm, CHDCl_2 in CD_2Cl_2 : 5.32 ppm, or C_6HD_5 in C_6D_6 : 7.15 ppm). ^{13}C NMR spectra were referenced to tetramethylsilane as an internal standard or to a solvent signal (CDCl_3 : 77.0 ppm, CD_2Cl_2 : 53.8 ppm, or C_6D_6 : 128.0 ppm). ^{31}P NMR spectra were referenced to external 85% H_3PO_4 in D_2O . The following abbreviations are used: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet, dd=double doublet, dt=double triplet, td=triple doublet, dq=double quartet, brs=broad singlet. Low- and high-resolution mass spectra (LRMS and HRMS) were obtained on JEOL JMS-DX303 and JMS-700 spectrometers, respectively, at the Mass Spectrometry Resource, Graduate School of Pharmaceutical Sciences, Tohoku University. Analytical HPLC was performed using Daicel Chiralpak AD. The samples of **4** subjected to HPLC were the purified products from column chromatography on silica gel.

Materials: Unless otherwise noted, materials were purchased from Tokyo Kasei Co., Aldrich Inc., or other commercial suppliers, and were used after appropriate purification (distillation or recrystallization). (*R,R*)-(+)-

Hydrobenzoin was obtained from Kanto Chemical Co. Ltd. Flash column chromatographies were performed on Kanto silica gel 60N (spherical, neutral, 70–230 mesh).

Representative procedure for the preparation of catalysts: (1*S*,2*S*)-Bis[tris(dimethylamino)phosphazido]-1,2-diphenylethane (1a**):** For use as a reagent: Under Ar atmosphere, tris(dimethylamino)phosphine (325.9 mg, 2.0 mmol) was added to a solution of (+)-(1*S*,2*S*)-1,2-diphenylethane-1,2-diazide (264.3 mg, 1.00 mmol) in benzene (1.5 mL) at room temperature. After stirring the mixture for 20 h at the same temperature, further dry benzene (1.0 mL) was added to give a 0.20 M solution of **1a**. Quantitative conversion was confirmed by ^{31}P NMR analysis.

For analysis: Under Ar atmosphere, tris(dimethylamino)phosphine (325.9 mg, 2.00 mmol) was added to a solution of (+)-(1*S*,2*S*)-1,2-diphenylethane-1,2-diazide (263.0 mg, 0.995 mmol) in benzene (1.5 mL) in a 20 mL Schlenk tube at room temperature. The mixture was stirred for 21 h at the same temperature. It was then concentrated under reduced pressure and **1a** was isolated in a glove box (517.0 mg, 88 %). The product was recrystallized from toluene/hexane as colorless needles. M.p. 96–97 °C; ^1H NMR (400 MHz, CD_2Cl_2): δ =2.61 (d, J =9.2 Hz, 36H), 5.13 (s, 2H), 6.95–7.12 (m, 6H), 7.15–7.25 ppm (m, 4H); $^{13}\text{C}\{^1\text{H}\}$ NMR (150 MHz, CD_2Cl_2): δ =36.8, 78.7, 125.4, 127.0, 128.7, 141.9 ppm; $^{31}\text{P}\{^1\text{H}\}$ NMR (243 MHz, CD_2Cl_2): δ =41.5 ppm (s); ^{31}P NMR (243 MHz, CD_2Cl_2): δ =41.5 ppm (septet, J =9.2 Hz); IR (neat): $\tilde{\nu}$ =2917, 2890, 2846, 2806, 1451, 1293, 1109, 974, 737, 699 cm^{-1} ; LRMS (FAB): m/z : 591 [$M+1$] $^+$; HRMS: m/z : calcd for $\text{C}_{26}\text{H}_{49}\text{N}_{12}\text{P}_2$: 591.3678; found: 591.3683.

(1*S*,2*S*)-Bis[tris(diethylamino)phosphazido]-1,2-diphenylethane (1b**):** Yellow oil; ^1H NMR (400 MHz, CDCl_3/TMS): δ =1.00 (t, J =7.1 Hz, 36H), 2.98–3.09 (m, 24H), 5.30 (s, 2H), 6.92–7.05 (m, 6H), 7.15–7.22 ppm (m, 4H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{C}_6\text{D}_6/\text{TMS}$): δ =13.8 (d, J =2.5 Hz), 39.7 (d, J =3.3 Hz), 80.8, 125.8, 127.6, 129.8, 143.7 ppm; $^{31}\text{P}\{^1\text{H}\}$ NMR (243 MHz, CD_2Cl_2): δ =42.4 ppm (s); ^{31}P NMR (243 MHz, CD_2Cl_2): δ =42.5 ppm (nonet, J =9.7 Hz); IR (neat): $\tilde{\nu}$ =2917, 2890, 2846, 2806, 1451, 1293, 1109, 974, 737, 699 cm^{-1} ; LRMS (FAB): m/z : 760 [$M+1$] $^+$; HRMS: m/z : calcd for $\text{C}_{38}\text{H}_{73}\text{N}_{12}\text{P}_2$: 759.5556; found: 759.5552.

(1*S*,2*S*)-Bis[tris(dibutylamino)phosphazido]-1,2-diphenylethane (1c**):** Yellow oil; ^1H NMR (400 MHz, CDCl_3/TMS): δ =0.95 (t, J =7.2 Hz, 36H), 1.16–1.34 (m, 24H), 1.54–1.67 (m, 24H), 3.00–3.14 (m, 24H), 5.61 (s, 2H), 6.94–7.01 (m, 2H), 7.06–7.27 (m, 6H), 7.52–7.56 ppm (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, $\text{C}_6\text{D}_6/\text{TMS}$): δ =14.3, 20.9, 31.1 (d, J =2.5 Hz), 46.5 (d, J =2.5 Hz), 81.0, 125.8, 127.6, 129.8, 144.1 ppm; $^{31}\text{P}\{^1\text{H}\}$ NMR (243 MHz, CD_2Cl_2): δ =41.9 ppm (s); ^{31}P NMR (243 MHz, CD_2Cl_2): δ =41.9 ppm (nonet, J =9.9 Hz); IR (neat): $\tilde{\nu}$ =2958, 2933, 2873, 1617, 1455, 1329, 1160, 1034, 926, 812, 698, 677 cm^{-1} ; LRMS (FAB): m/z : 1096 [$M+1$] $^+$; HRMS: m/z : calcd for $\text{C}_{62}\text{H}_{121}\text{N}_{12}\text{P}_2$: 1095.9312; found: 1095.9342.

(1*S*,2*S*)-Bis[tris(pyrrolidino)phosphazido]-1,2-diphenylethane (1d**):** Yellow oil; ^1H NMR (400 MHz, CDCl_3/TMS): δ =1.67–1.75 (m, 24H), 3.10–3.20 (m, 24H), 5.26 (s, 2H), 6.93–7.06 (m, 6H), 7.15–7.21 ppm (m, 4H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3/TMS): δ =26.5 (d, J =6.4 Hz), 47.2 (d, J =2.5 Hz), 81.6, 125.9, 127.7, 129.7, 143.6 ppm; $^{31}\text{P}\{^1\text{H}\}$ NMR (243 MHz, CDCl_3): δ =31.5 ppm (s); ^{31}P NMR (243 MHz, CDCl_3): δ =31.3–31.6 ppm (m); IR (neat): $\tilde{\nu}$ =2958, 2865, 2850, 1445, 1204, 1111, 1069, 1009, 754, 697 cm^{-1} ; LRMS (FAB): m/z : 748 [$M+1$] $^+$; HRMS: m/z : calcd for $\text{C}_{38}\text{H}_{61}\text{N}_{12}\text{P}_2$: 747.4617; found: 747.4630.

Representative procedure for the conjugate addition of dialkyl malonates to chalcones: 2-(3-Oxo-1,3-diphenyl-propyl)-malonic acid dimethyl ester (4a**):**^[12a,13b,14] A solution of *trans*-chalcone (104.1 mg, 0.50 mmol) in CH_2Cl_2 (0.5 mL) was added dropwise to a mixture of LiClO_4 (5.3 mg, 0.05 mmol), **1a** (0.25 mL, 0.20 M in benzene, 0.05 mmol), and dimethyl malonate (132.1 mg, 1.00 mmol) in CH_2Cl_2 (0.5 mL) at -40°C under an Ar atmosphere. The reaction mixture was stirred for 48 h, then quenched with saturated NH_4Cl solution and extracted with AcOEt . The organic layer was washed with brine, dried over MgSO_4 , and concentrated using a rotary evaporator to afford the crude product. Purification by column chromatography on silica gel (0–15% AcOEt in hexane) afforded **4a**. The product was recrystallized from Et_2O /hexane as colorless needles. M.p. 84–85 °C; $[\alpha]_{\text{D}}^{25}$ =+24.20 (c =1.01 in CHCl_3) (lit.:^[14] $[\alpha]_{\text{D}}^{25}$ =+17.7 (c =1.2 in CHCl_3)); ^1H NMR (400 MHz, CDCl_3/TMS): δ =3.48 (dd, J =

16.8, 8.8 Hz, 1H), 3.51 (s, 3H), 3.55 (dd, $J=16.8$, 5.2 Hz, 1H), 3.73 (s, 3H), 3.86 (d, $J=9.6$ Hz, 1H), 4.19 (dt, $J=9.2$, 5.2 Hz, 1H), 7.14–7.29 (m, 5H), 7.42 (t, $J=7.6$ Hz, 2H), 7.53 (tt, $J=7.6$, 1.6 Hz, 1H), 7.87–7.92 ppm (m, 2H); IR (neat): $\tilde{\nu}=2954$, 2844, 1727, 1681, 1432, 1233, 1158, 1025, 747, 687 cm^{-1} ; LRMS (EI): m/z : 340 [M^+]; HRMS: m/z : calcd for $\text{C}_{20}\text{H}_{20}\text{O}_5$: 340.1311; found: 340.1307; elemental analysis calcd (%) for $\text{C}_{20}\text{H}_{20}\text{O}_5$: C 70.57, H 5.92; found: C 70.45, H 5.97; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=28.4$ min (minor enantiomer), $t_R=35.2$ min (major enantiomer).

2-(3-Oxo-1,3-diphenyl-propyl)-malonic acid diethyl ester (4b):^[13b,14] Recrystallized from Et_2O /hexane, colorless needles; m.p. 73–74 °C; $[\alpha]_D^{25}=+22.50$ ($c=1.01$ in CHCl_3) (lit.:^[14] $[\alpha]_D^{25}=+17.5$ ($c=0.8$ in CHCl_3)); ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=1.01$ (t, $J=7.1$ Hz, 3H), 1.24 (t, $J=7.1$ Hz, 3H), 3.45 (dd, $J=16.6$, 9.0 Hz, 1H), 3.54 (dd, $J=16.6$, 4.6 Hz, 1H), 3.82 (d, $J=9.8$ Hz, 1H), 3.95 (q, $J=7.1$ Hz, 2H), 4.13–4.24 (m, 3H), 7.13–7.30 (m, 5H), 7.38–7.55 (m, 3H), 7.86–7.92 ppm (m, 2H); IR (neat): $\tilde{\nu}=2983$, 1721, 1679, 1293, 1239, 1167, 1030, 745, 700, 687 cm^{-1} ; LRMS (EI): m/z : 368 [M^+]; HRMS: m/z : calcd for $\text{C}_{22}\text{H}_{24}\text{O}_5$: 368.1624; found: 368.1612; elemental analysis calcd (%) for $\text{C}_{22}\text{H}_{24}\text{O}_5$: C 71.72, H 6.57; found: C 71.72, H 6.42; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=22.4$ min (minor enantiomer), $t_R=39.1$ min (major enantiomer).

2-(3-Oxo-1,3-diphenyl-propyl)-malonic acid diisopropyl ester (4c):^[13b,14] Recrystallized from Et_2O /hexane, colorless needles; m.p. 88–89 °C; $[\alpha]_D^{25}=+15.80$ ($c=1.01$ in CHCl_3) (lit.:^[14] $[\alpha]_D^{25}=+21.3$ ($c=0.7$ in CHCl_3)); ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=0.97$ (d, $J=6.3$ Hz, 3H), 1.04 (d, $J=6.3$ Hz, 3H), 1.24 (d, $J=6.3$ Hz, 6H), 3.43 (dd, $J=16.5$, 9.5 Hz, 1H), 3.53 (dd, $J=16.5$, 4.1 Hz, 1H), 4.15 (dt, $J=9.7$, 4.1 Hz, 1H), 4.79 (sept, $J=6.3$ Hz, 1H), 5.07 (sept, $J=6.3$ Hz, 1H), 7.12–7.28 (m, 5H), 7.38–7.44 (m, 2H), 7.49–7.54 (m, 1H), 7.86–7.91 ppm (m, 2H); IR (neat): $\tilde{\nu}=2983$, 2935, 1721, 1683, 1281, 1235, 1104, 750, 702, 691 cm^{-1} ; LRMS (EI): m/z : 396 [M^+]; HRMS: m/z : calcd for $\text{C}_{24}\text{H}_{28}\text{O}_5$: 396.1937; found: 396.1928; elemental analysis calcd (%) for $\text{C}_{24}\text{H}_{28}\text{O}_5$: C 72.70, H 7.12; found: C 72.58, H 7.09; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=12.88$ min (minor enantiomer), $t_R=24.82$ min (major enantiomer).

2-(3-Oxo-1,3-diphenyl-propyl)-malonic acid dibenzyl ester (4d):^[13b,14] Recrystallized from Et_2O /hexane, colorless needles; m.p. 106–107 °C; $[\alpha]_D^{25}=+16.20$ ($c=1.01$ in CHCl_3) (lit.:^[14] $[\alpha]_D^{25}=+18.4$ ($c=1.0$ in CHCl_3)); ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.49$ (d, $J=6.4$ Hz, 2H), 3.94 (d, $J=9.5$ Hz, 1H), 4.18–4.25 (m, 1H), 4.90 (s, 2H), 5.11 (d, $J=12.2$ Hz, 1H), 5.16 (d, $J=12.2$ Hz, 1H), 7.04–7.42 (m, 17H), 7.48–7.53 (m, 1H), 7.77–7.82 ppm (m, 2H); IR (neat): $\tilde{\nu}=3346$, 3031, 1733, 1681, 1229, 1152, 1011, 965, 741, 699 cm^{-1} ; LRMS (EI): m/z : 492 [M^+]; HRMS: m/z : calcd for $\text{C}_{32}\text{H}_{28}\text{O}_5$: 492.1937; found: 492.1957; elemental analysis calcd (%) for $\text{C}_{32}\text{H}_{28}\text{O}_5$: C 78.03, H 5.73; found: C 77.99, H 5.86; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=38.86$ min (minor enantiomer), $t_R=73.36$ min (major enantiomer).

2-(3-Oxo-1,3-diphenyl-propyl)-malonic acid di-*tert*-butyl ester (4e):^[16] Recrystallized from Et_2O /hexane, colorless needles; m.p. 117–118 °C; $[\alpha]_D^{25}=+8.57$ ($c=0.75$ in CHCl_3); ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=1.19$ (s, 9H), 1.46 (s, 9H), 3.39 (dd, $J=16.4$, 9.9 Hz, 1H), 3.51 (dd, $J=16.4$, 3.9 Hz, 1H), 3.64 (d, $J=10.1$ Hz, 1H), 4.07 (dt, $J=10.0$, 3.9 Hz, 1H), 7.11–7.17 (m, 1H), 7.19–7.27 (m, 4H), 7.38–7.44 (m, 2H), 7.49–7.54 (m, 1H), 7.87–7.91 ppm (m, 2H); IR (neat): $\tilde{\nu}=2979$, 1735, 1719, 1683, 1368, 1246, 1138, 741, 700, 687 cm^{-1} ; LRMS (EI): m/z : 368 [$M-56$] $^+$; HRMS: m/z : calcd for $\text{C}_{26}\text{H}_{32}\text{O}_5$: 368.1624; found: 368.1635; elemental analysis calcd (%) for $\text{C}_{26}\text{H}_{32}\text{O}_5$: C 73.56, H 7.60; found: C 73.57, H 7.57; HPLC (Daicel Chiralpak AD, isopropanol/hexane 5:95, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=12.15$ min (minor enantiomer), $t_R=41.15$ min (major enantiomer).

2-(3-Oxo-1,3-diphenyl-propyl)-malononitrile (4f):^[13b,14] Recrystallized from CH_2Cl_2 /hexane, colorless needles; m.p. 125–126 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.65$ (dd, $J=18.6$, 5.3 Hz, 1H), 3.72 (d, $J=18.6$, 8.5 Hz, 1H), 3.96 (dt, $J=8.5$, 5.1 Hz, 1H), 4.66 (d, $J=5.1$ Hz, 1H), 7.35–7.54 (m, 7H), 7.60–7.66 (m, 1H), 7.96–8.00 ppm (m, 2H); IR (neat): $\tilde{\nu}=2919$, 2900, 2256, 1681, 1449, 1233, 762, 748, 700, 690 cm^{-1} ; LRMS (EI): m/z : 274 [M^+]; HRMS: m/z : calcd for $\text{C}_{18}\text{H}_{14}\text{N}_2\text{O}$: 274.1106; found:

274.1099; elemental analysis calcd (%) for $\text{C}_{18}\text{H}_{14}\text{N}_2\text{O}$: C 78.81, H 5.14, N 10.21; found: C 78.81, H 5.34, N 10.07; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=20.10$ min (major enantiomer), $t_R=27.67$ min (minor enantiomer).

2-[3-Oxo-3-phenyl-1-(4-tolyl)-propyl]-malonic acid dimethyl ester (4g):^[17] Recrystallized from Et_2O /hexane, colorless needles; m.p. 75–76 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=2.26$ (s, 3H), 3.40–3.57 (m, 5H), 3.72 (s, 3H), 3.83 (d, $J=9.5$ Hz, 1H), 4.15 (dt, $J=9.1$, 5.1 Hz, 1H), 7.04 (d, $J=8.3$ Hz, 2H), 7.13 (d, $J=8.3$ Hz, 2H), 7.39–7.46 (m, 2H), 7.50–7.56 (m, 1H), 7.88–7.93 ppm (m, 2H); IR (neat): $\tilde{\nu}=2954$, 1729, 1681, 1233, 1208, 1194, 1160, 1025, 820, 688 cm^{-1} ; LRMS (EI): m/z : 354 [M^+]; HRMS: m/z : calcd for $\text{C}_{21}\text{H}_{22}\text{O}_5$: 354.1467; found: 354.1478; elemental analysis calcd (%) for $\text{C}_{21}\text{H}_{22}\text{O}_5$: C 71.17, H 6.26; found: C 70.71, H 6.10; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=24.0$ min (minor enantiomer), $t_R=31.6$ min (major enantiomer).

2-[1-(4-Methoxyphenyl)-3-oxo-3-phenylpropyl]-malonic acid dimethyl ester (4h):^[17,18] Recrystallized from Et_2O /hexane, colorless needles; m.p. 81–82 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.43$ (dd, $J=16.6$, 8.9 Hz, 1H), 3.52 (dd, $J=16.6$, 4.8 Hz, 1H), 3.52 (s, 3H), 3.73 (s, 3H), 3.74 (s, 3H), 3.82 (d, $J=9.4$ Hz, 1H), 4.10–4.18 (m, 1H), 6.75–6.81 (m, 2H), 7.14–7.20 (m, 2H), 7.40–7.46 (m, 2H), 7.50–7.56 (m, 1H), 7.88–7.92 ppm (m, 2H); IR (neat): $\tilde{\nu}=2954$, 1729, 1681, 1515, 1231, 1179, 1158, 1025, 754, 729, 686 cm^{-1} ; LRMS (EI): m/z : 370 [M^+]; HRMS: m/z : calcd for $\text{C}_{21}\text{H}_{22}\text{O}_6$: 370.1416; found: 370.1413; elemental analysis calcd (%) for $\text{C}_{21}\text{H}_{22}\text{O}_6$: C 68.10, H 5.99; found: C 67.63, H 6.00; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=51.76$ min (minor enantiomer), $t_R=75.66$ min (major enantiomer).

2-[1-(4-Chlorophenyl)-3-oxo-3-phenylpropyl]-malonic acid dimethyl ester (4i): Recrystallized from Et_2O /hexane, colorless needles; m.p. 87–88 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.45$ (dd, $J=17.0$, 8.9 Hz, 1H), 3.53 (dd, $J=17.0$, 4.8 Hz, 1H), 3.53 (s, 3H), 3.74 (s, 3H), 3.82 (d, $J=9.2$ Hz, 1H), 4.17 (dt, $J=9.2$, 4.8 Hz, 1H), 7.18–7.26 (m, 4H), 7.40–7.47 (m, 2H), 7.52–7.58 (m, 1H), 7.87–7.92 ppm (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3/TMS): $\delta=40.1$, 42.1, 52.5, 52.7, 57.0, 128.0, 128.6, 128.6, 129.5, 133.0, 133.2, 136.6, 139.0, 168.0, 168.5, 197.2 ppm; IR (neat): $\tilde{\nu}=2956$, 1729, 1681, 1235, 1162, 1023, 827, 752, 685, 665 cm^{-1} ; LRMS (EI): m/z : 374 [M^+]; HRMS: m/z : calcd for $\text{C}_{20}\text{H}_{19}\text{O}_5\text{Cl}$: 374.0921; found: 374.0904; elemental analysis calcd (%) for $\text{C}_{20}\text{H}_{19}\text{O}_5\text{Cl}$: C 64.09, H 5.11; found: C 63.97, H 5.11; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=32.86$ min (minor enantiomer), $t_R=52.03$ min (major enantiomer).

2-[1-(4-Fluorophenyl)-3-oxo-3-phenylpropyl]-malonic acid dimethyl ester (4j): Recrystallized from Et_2O /hexane, colorless needles; m.p. 85–86 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.44$ (dd, $J=16.9$, 8.9 Hz, 1H), 3.40–3.57 (m, 5H), 3.72 (s, 3H), 3.83 (d, $J=9.5$ Hz, 1H), 4.15 (dt, $J=9.1$, 5.1 Hz, 1H), 7.04 (d, $J=8.3$ Hz, 2H), 7.13 (d, $J=8.3$ Hz, 2H), 7.39–7.46 (m, 2H), 7.50–7.56 (m, 1H), 7.88–7.93 ppm (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (100 MHz, CDCl_3/TMS): $\delta=40.1$, 42.3, 52.4, 52.7, 57.2, 115.3 (d, $J=21.3$ Hz), 128.1, 128.6, 129.7 (d, $J=7.8$ Hz), 129.8, 133.1 (d, $J=3.3$ Hz), 136.7, 161.8 (d, $J=245.8$ Hz), 168.0, 168.6, 197.3 ppm; IR (neat): $\tilde{\nu}=2954$, 1729, 1681, 1233, 1208, 1194, 1160, 1025, 820, 688 cm^{-1} ; LRMS (EI): m/z : 354 [M^+]; HRMS: m/z : calcd for $\text{C}_{21}\text{H}_{22}\text{O}_5$: 354.1467; found: 354.1478; elemental analysis calcd (%) for $\text{C}_{21}\text{H}_{22}\text{O}_5$: C 71.17, H 6.26; found: C 70.71, H 6.10; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=24.0$ min (minor enantiomer), $t_R=31.6$ min (major enantiomer).

2-[1-(4-Nitrophenyl)-3-oxo-3-phenylpropyl]-malonic acid dimethyl ester (4k):^[18] Recrystallized from Et_2O /hexane, colorless needles; m.p. 77–78 °C; ^1H NMR (400 MHz, CDCl_3/TMS): $\delta=3.50$ –3.63 (m, 5H), 3.75 (s, 3H), 3.89 (d, $J=9.1$ Hz, 1H), 4.31 (dt, $J=8.9$, 5.1 Hz, 1H), 7.41–7.50 (m, 4H), 7.53–7.79 (m, 1H), 7.86–7.92 (m, 2H), 8.10–8.16 ppm (m, 2H); IR (neat): $\tilde{\nu}=2954$, 1731, 1684, 1517, 1345, 1256, 1152, 856, 748, 688 cm^{-1} ; LRMS (EI): m/z : 385 [M^+]; HRMS: m/z : calcd for $\text{C}_{20}\text{H}_{19}\text{NO}_5$: 385.1162; found: 385.1154; elemental analysis calcd (%) for $\text{C}_{20}\text{H}_{19}\text{NO}_5$: C 62.33, H 4.97, N 3.63; found: C 62.08, H 5.05, N 3.40; HPLC (Daicel Chiralpak AD, isopropanol/hexane 30:70, flow rate 1.0 mL min^{-1} , $\lambda=254$ nm): $t_R=29.3$ min (minor enantiomer), $t_R=59.7$ min (major enantiomer).

2-[1-(2-Chlorophenyl)-3-oxo-3-phenylpropyl]-malonic acid dimethyl ester (4l):^[18] Pale-yellow oil; ¹H NMR (400 MHz, CDCl₃/TMS): δ = 3.57–3.76 (m, 8H), 4.10 (d, J = 8.4 Hz, 1H), 4.67 (dt, J = 8.6, 4.8 Hz, 1H), 7.10–7.19 (m, 2H), 7.27–7.37 (m, 2H), 7.40–7.46 (m, 2H), 7.51–7.57 (m, 1H), 7.90–7.95 ppm (m, 2H); IR (neat): $\tilde{\nu}$ = 2954, 1733, 1684, 1436, 1227, 1154, 1021, 750, 690 cm⁻¹; LRMS (EI): m/z : 374 [M⁺]; HRMS: m/z : calcd for C₂₀H₁₉ClO₅: 374.0921; found: 374.0929; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min⁻¹, λ = 254 nm): t_R = 17.5 min (minor enantiomer), t_R = 19.6 min (major enantiomer).

2-(1-Furan-2-yl-3-oxo-3-phenylpropyl)-malonic acid dimethyl ester (4m): Recrystallized from Et₂O/hexane, colorless crystals; m.p. 40–41 °C; ¹H NMR (400 MHz, CDCl₃/TMS): δ = 3.47 (dd, J = 17.1, 4.8 Hz, 1H), 3.58 (dd, J = 17.1, 8.5 Hz, 1H), 3.64 (s, 3H), 3.73 (s, 3H), 4.30–4.37 (m, 1H), 6.09–6.13 (m, 1H), 6.21–6.24 (m, 1H), 7.25–7.28 (m, 1H), 7.42–7.48 (m, 2H), 7.51–7.58 (m, 1H), 7.92–7.98 ppm (m, 2H); ¹³C{¹H} NMR (100 MHz, CDCl₃/TMS): δ = 10.9, 34.3, 39.6, 52.6, 54.8, 107.0, 110.3, 128.1, 128.6, 133.2, 136.6, 141.7, 153.4, 168.2, 168.4, 197.2 ppm; IR (neat): $\tilde{\nu}$ = 2960, 2921, 1744, 1735, 1683, 1256, 1158, 1146, 1007, 760, 727, 686 cm⁻¹; LRMS (EI): m/z : 330 [M⁺]; HRMS: m/z : calcd for C₁₈H₁₈O₆: 330.1103; found: 330.1103; elemental analysis calcd (%) for C₁₈H₁₈O₆: C 65.45, H 5.49; found: C 65.56, H 5.62; HPLC (Daicel Chiralcel OD, isopropanol/hexane 5:95, flow rate 1.0 mL min⁻¹, λ = 254 nm): t_R = 15.80 min (minor enantiomer), t_R = 16.92 min (major enantiomer).

2-(3-Oxo-3-phenyl-1-(pyridin-2-yl)-propyl)-malonic acid dimethyl ester (4n): Recrystallized from Et₂O/hexane, colorless crystals; m.p. 74–75 °C; ¹H NMR (400 MHz, CDCl₃/TMS): δ = 3.42 (dd, J = 17.6, 3.9 Hz, 1H), 3.54 (s, 3H), 3.74 (s, 3H), 3.79 (dd, J = 17.6, 9.5 Hz, 1H), 4.11 (d, J = 9.6 Hz, 1H), 4.32 (dt, J = 9.5, 3.9 Hz, 1H), 7.05–7.09 (m, 1H), 7.34–7.45 (m, 3H), 7.49–7.60 (m, 2H), 7.88–7.93 (m, 2H), 8.44–8.47 ppm (m, 1H); ¹³C{¹H} NMR (100 MHz, CDCl₃/TMS): δ = 41.5, 41.7, 52.3, 52.6, 55.8, 121.9, 124.8, 128.1, 128.5, 133.0, 136.2, 136.7, 149.0, 160.3, 168.4, 169.0, 197.7 ppm; LRMS (EI): m/z : 341 [M⁺]; HRMS: m/z : calcd for C₁₉H₁₉NO₅: 341.1263; found: 341.1254; elemental analysis calcd (%) for C₁₉H₁₉NO₅: C 66.85, H 5.61, N 4.10; found: C 66.82, H 5.84, N 3.85; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min⁻¹, λ = 254 nm): t_R = 30.6 min (minor enantiomer), t_R = 52.2 min (major enantiomer).

2-[3-(4-Chlorophenyl)-3-oxo-1-phenylpropyl]-malonic acid dimethyl ester (4o): Recrystallized from Et₂O/hexane, colorless needles; m.p. 71–72 °C; ¹H NMR (400 MHz, CDCl₃/TMS): δ = 3.42 (dd, J = 16.9, 8.9 Hz, 1H), 3.51 (s, 3H), 3.53 (dd, J = 16.9, 4.8 Hz, 1H), 3.73 (s, 3H), 3.84 (d, J = 9.2 Hz, 1H), 4.16 (dt, J = 9.2, 4.8 Hz, 1H), 7.15–7.28 (m, 5H), 7.36–7.43 (m, 2H), 7.81–7.86 ppm (m, 2H); ¹³C{¹H} NMR (100 MHz, CDCl₃/TMS): δ = 40.8, 42.3, 52.4, 52.6, 57.2, 127.3, 128.0, 128.5, 128.9, 129.5, 135.1, 139.5, 140.2, 168.1, 168.7, 196.4 ppm; IR (neat): $\tilde{\nu}$ = 2954, 1727, 1681, 1235, 1210, 1154, 1090, 1025, 825, 702 cm⁻¹; LRMS (EI): m/z : 374 [M⁺]; HRMS: m/z : calcd for C₂₀H₁₉O₅Cl: 374.0921; found: 374.0931; elemental analysis calcd (%) for C₂₀H₁₉O₅Cl: C 64.09, H 5.11; found: C 63.92, H 5.07; HPLC (Daicel Chiralpak AD, isopropanol/hexane 10:90, flow rate 1.0 mL min⁻¹, λ = 254 nm): t_R = 36.86 min (minor enantiomer), t_R = 51.84 min (major enantiomer).

NMR spectroscopic analyses for the titration of 1a with LiClO₄: ¹H and ³¹P NMR spectra were obtained on a JEOL ECA600 spectrometer (600.172 MHz for ¹H, 242.956 MHz for ³¹P). Unless otherwise noted, spectra were recorded from approximately 0.10 M solutions of the samples in CD₂Cl₂ at 25 °C. ¹H NMR spectra were referenced to a solvent signal (CHDCl₂, 5.32 ppm). ³¹P NMR spectra were referenced to external 85 % H₃PO₄ in D₂O.

Acknowledgements

This research was partly supported by Grants-in-Aid for Scientific Research on Priority Areas “Advanced Molecular Transformation of Carbon Resources” and “Synergistic Effects for Creation of Functional Molecules” (to Y.K.), and Grants-in-Aid for Young Scientists (B) (to H.N.).

- a) R. Schwesinger, H. Schlemper, *Angew. Chem.* **1987**, *99*, 1212–1214; *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 1167–1169; b) R. Schwesinger, H. Schlemper, C. Hasenfratz, F. Willaredt, T. Dambacher, T. Breuer, C. Ottaway, M. Fletschinger, F. Boele, J. Fritz, D. Putzas, J. W. Rotter, F. G. Bordwell, A. V. Satish, G.-Z. Ji, E.-M. Peters, K. Peters, H. G. von Schnering, L. Walz, *Liebigs Ann.* **1996**, 1055–1081.
- a) H. R. Allcock, “*Chemistry and Applications of Polyphosphazenes*”, Wiley, Hoboken, **2003**; b) J. Paulsdorf, N. Kaskhedikar, M. Burjanadze, S. Obeidi, N. A. Stolwijk, D. Wilmer, H.-D. Wiemhofer, *Chem. Mater.* **2006**, *18*, 1281–1288; c) J. X. Zhang, L. Y. Qiu, Y. Jin, K. J. Zhu, *Macromolecules* **2006**, *39*, 451–455; d) K. Huynh, A. J. Lough, I. Manners, *J. Am. Chem. Soc.* **2006**, *128*, 14002–14003.
- a) F. Palacios, E. Herran, C. Alonso, G. Rubiales, B. Lecea, M. Ayerbe, F. P. Cossio, *J. Org. Chem.* **2006**, *71*, 6020–6030; b) A. Solladié-Cavallo, M. Roje, R. Welter, V. Sunjic, *J. Org. Chem.* **2004**, *69*, 1409–1412.
- a) M. O'Donnell, F. Delgado, C. Hostettler, R. Schwesinger, *Tetrahedron Lett.* **1998**, *39*, 8775–8778; b) M. Caldarelli, J. Habermann, S. V. Ley, *J. Chem. Soc. Perkin Trans. 1* **1999**, 107–110; c) G. A. Kraus, N. Zhang, J. G. Verkade, M. Nagarajan, P. B. Kisanga, *Org. Lett.* **2000**, *2*, 2409–2410; d) A. Clark, D. Patel, M. J. Broadhurst, *Tetrahedron Lett.* **2003**, *44*, 7763–7765; e) T. Imahori, Y. Kondo, *J. Am. Chem. Soc.* **2003**, *125*, 8082–8083; f) T. Imahori, C. Hori, Y. Kondo, *Adv. Synth. Catal.* **2004**, *346*, 1090–1092; g) D. Bensa, T. Constantieux, J. Rodriguez, *Synthesis* **2004**, 923–927; h) V. Raab, E. Gauchenova, A. Merkoulou, K. Harms, J. Sundermeyer, B. Kovacevic, Z. B. Maksic, *J. Am. Chem. Soc.* **2005**, *127*, 15738–15743; i) D. Uraguchi, S. Sakaki, T. Ooi, *J. Am. Chem. Soc.* **2007**, *129*, 12392–12393.
- a) M. T. Reetz, E. Bohres, *Chem. Commun.* **1998**, 935–936; b) J. M. Brunel, O. Légrand, S. Reymond, G. Buono, *J. Am. Chem. Soc.* **1999**, *121*, 5807–5808; c) M. Sauthier, J. Fornies-Camer, L. Toupet, R. Reau, *Organometallics* **2000**, *19*, 553–562; d) U. Köhn, M. Schulz, A. Schramm, W. Günther, H. Görls, S. Schenk, E. Anders, *Eur. J. Org. Chem.* **2006**, 4128–4134; e) M. Ueno, A. E. H. Wheatley, Y. Kondo, *Chem. Commun.* **2006**, 3549–3550.
- a) M. Ueno, C. Hori, K. Suzawa, M. Ebisawa, Y. Kondo, *Eur. J. Org. Chem.* **2005**, 1965–1968; b) K. Kobayashi, M. Ueno, Y. Kondo, *Chem. Commun.* **2006**, 3128–3130; c) K. Suzawa, M. Ueno, A. E. H. Wheatley, Y. Kondo, *Chem. Commun.* **2006**, 4850–4852; d) M. Ueno, M. Yonemoto, M. Hashimoto, A. E. H. Wheatley, H. Naka, Y. Kondo, *Chem. Commun.* **2007**, 2264–2266.
- Synthesis of chiral phosphazenes: a) X. Liu, P. Ilankumaran, I. A. Guzei, J. G. Verkade, *J. Org. Chem.* **2000**, *65*, 701–706; b) P. Ilankumaran, G. Zhang, J. G. Verkade, *Heteroat. Chem.* **2000**, *11*, 251–253; c) U. Köhn, M. Schulz, A. Schramm, W. Günther, S. Schenk, E. Anders, *Eur. J. Org. Chem.* **2006**, 4128–4134.
- a) H. Staudinger, *Helv. Chim. Acta* **1919**, *2*, 635–646; reviews: b) Y. G. Gololobov, I. N. Zhmurova, L. F. Nasukhin, *Tetrahedron* **1981**, *37*, 437–472; c) Yu. G. Gololobov, I. N. Zhmurova, L. F. Kasukhin, *Tetrahedron* **1992**, *48*, 1353–1406; recent theoretical studies: d) W. Q. Tian, Y. A. Wang, *J. Org. Chem.* **2004**, *69*, 4299–4308; e) W. Q. Tian, Y. A. Wang, *J. Chem. Theory Comput.* **2005**, *1*, 353–362.
- a) J. G. Verkade, P. B. Kisanga, *Tetrahedron* **2003**, *59*, 7819–7858, and references therein; b) X. Liu, N. Thirupathi, I. A. Guzei, J. G. Verkade, *Inorg. Chem.* **2004**, *43*, 7431–7440; c) Ch. Venkat Reddy, J. G. Verkade, *J. Org. Chem.* **2007**, *72*, 3093–3096; d) Ch. Venkat Reddy, B. M. Fetterly, J. G. Verkade, *Energy Fuels* **2007**, *21*, 2466–2472.
- Conventional preparation of phosphazene bases generally requires deprotonation of phosphonium salts and purification of the product. The use of an in situ deprotonated phosphonium salt as a catalyst has also recently been reported, see ref. [4i].
- a) *Comprehensive Asymmetric Catalysis* (Eds.: E. N. Jacobsen, A. Pfaltz, H. Yamamoto), Springer, Berlin, **1999**; b) J.-A. Ma, D. Cahard, *Angew. Chem.* **2004**, *116*, 4666–4683; *Angew. Chem. Int.*

- Ed.* **2004**, 43, 4566–4583; c) H. Li, Y. Wang, L. Tang, L. Deng, *J. Am. Chem. Soc.* **2004**, 126, 9906–9907; d) H. Huang, E. N. Jacobsen, *J. Am. Chem. Soc.* **2006**, 128, 7170–7171; e) S. B. Tsogoeva, *Eur. J. Org. Chem.* **2007**, 1701–1716.
- [12] a) H. Sasai, T. Arai, Y. Satow, K. N. Houk, M. Shibasaki, *J. Am. Chem. Soc.* **1995**, 117, 6194–6198; b) M. Shibasaki, N. Yoshikawa, *Chem. Rev.* **2002**, 102, 2187–2210.
- [13] a) T. Ooi, K. Maruoka, *Acc. Chem. Res.* **2004**, 37, 526–533; b) T. Ooi, D. Ohara, K. Fukumoto, K. Maruoka, *Org. Lett.* **2005**, 7, 3195–3197.
- [14] J. Wang, H. Li, L. Zu, W. Jiang, H. Xie, W. Duan, W. Wang, *J. Am. Chem. Soc.* **2006**, 128, 12652–12653.
- [15] D. Pini, A. Iuliano, C. Rosini, P. Salvadori, *Synthesis* **1990**, 1023–1024.
- [16] N. End, L. Macko, M. Zehnder, A. Pfaltz, *Chem. Eur. J.* **1998**, 4, 818–824.
- [17] B. M. Choudary, M. Lakshmi Kantam, B. Kavita, Ch. Venkat Reddy, F. Figueras, *Tetrahedron* **2000**, 56, 9357–9364.
- [18] W. Ye, J. Xu, C.-T. Tan, C.-H. Tan, *Tetrahedron Lett.* **2005**, 46, 6875–6878.

Received: February 5, 2008
Published online: April 22, 2008