This article was downloaded by: [Florida State University] On: 19 October 2014, At: 13:03 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Deselenation of Overcrowded Triselenane Derivatives Containing Bismuth: Synthesis and Structure of the First Stable Dibismuthene

Norihiro Tokitoh , Yoshimitsu Arai & Renji Okazaki

^a Department of Chemistry, Graduate School of Science , The University of Tokyo , Hongo, Bunkyo-ku, Tokyo 113, Japan

^b Department of Chemistry, Graduate School of Science , The University of Tokyo , Hongo, Bunkyo-ku, Tokyo 113, Japan

^c Department of Chemistry, Graduate School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan Published online: 02 Apr 2008.

To cite this article: Norihiro Tokitoh , Yoshimitsu Arai & Renji Okazaki (1997) Deselenation of Overcrowded Triselenane Derivatives Containing Bismuth: Synthesis and Structure of the First Stable Dibismuthene, Phosphorus, Sulfur, and Silicon and the Related Elements, 124:1, 371-378, DOI: 10.1080/10426509708545643

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

DESELENATION OF OVERCROWDED TRISELENANE DERIVATIVES CONTAINING BISMUTH: SYNTHESIS AND STRUCTURE OF THE FIRST STABLE DIBISMUTHENE

NORIHIRO TOKITOH, YOSHIMITSU ARAI, and RENJI OKAZAKI Department of Chemistry, Graduate School of Science, The University of Tokyo, Hongo, Bunkyo-ku, Tokyo 113, Japan.

Deselenation of an overcrowded triselenatribismane, 2,4,6-{2,4,6-tris[bis(trimethylsilyl)methyl]phenyl}-1,3,5-triselenatribismane, with hexamethylphosphorous triamide resulted in the formation of the first stable dibismuthene {TbtBi=BiTbt; Tbt = 2,4,6-tris[bis(trimethyl-silyl)methyl]phenyl}, *i. e.* the doubly bonded compound consisting of the heaviest stable element in the periodic table, as deep purple crystals. X-ray crystallographic structural analysis, UV-vis and Raman spectra, and the theoretical calculations revealed the unique bonding character of the unprecedented Bi-Bi double bond.

<u>Keywords:</u> deselenation; steric protection; triselenatribismane; dibismuthene; X-ray structural analysis

INTRODUCTION

The synthesis of doubly bonded compounds containing heavier main group elements has stimulated wide interest^[1-5] because of their unusual structure and properties. Although there have been several examples of stable doubly bonded compounds of heavier group 14– 16 elements such as $R_2E=ER_2$ (E = Si,^[6] Ge,^[7] Sn^[8,9]), RE=ER (E = P,^[10] As^[11]), and $R_2E=X$ (E = Si, Ge; X = S, Se, Te)^[12–15], no stable examples containing 6th-raw element(s) have been reported so far. Here we report the first synthesis of a stable dibismuthene, TbtBi=BiTbt (1), *i. e.* the long-sought doubly bonded compound consisting of the heaviest stable element in the periodic table, by utilizing a novel synthetic methodology and an efficient steric protection group, 2,4,6-tris[bis(trimethylsilyl)methyl]phenyl (denoted as Tbt hereafter) developed by us.^[16]



RESULTS AND DISCUSSION

For the preparation of 1 we have developed a new synthetic method, *i. e.* deselenation reaction of triselenatribismane 2 with a phosphine reagent. As shown in Scheme 1, the precursor 2 was readily synthesized by the nucleophilic substitution of bismuth trichloride with TbtLi giving the corresponding bismuth dichloride, TbtBiCl₂ (3), followed by treatment of 3 with Li₂Se in tetrahydrofuran.



Triselenatribismane 2 thus isolated as a stable crystalline compound was then treated with an excess amount of hexamethylphosphorous triamide in toluene at 100 °C in a sealed tube. After heating for 12 h the solution turned purple and the expected dibismuthene 1, which precipitated from the mixture on cooling, was isolated by filtration in a glovebox filled with argon as deep purple single crystals (Scheme 2). The dibismuthene 1 has a very low solubility probably due to its high symmetry in the molecular structure, thus making its isolation quite simple.



Scheme 2

Dibismuthene 1 is the first compound containing a bismuthbismuth double bond which is the heaviest among those which consist of stable elements in the periodic table and the molecular geometry of 1 was finally determined by X-ray crystallographic structural analysis as shown in Figure 1. Dibismuthene 1 is in the trans-form, and has a center of symmetry, the two bismuth atoms and the two ipso carbons of the Tbt groups attached to bismuth atoms all lying in the same plane. The benzene ring of the Tbt groups are placed almost perpendicular to this plane with the dihedral angle of 90.70°. Two important parameters in the structure of 1 are the Bi-Bi bond length [2.8206(8) Å] and the Bi-Bi-C angle [100.5(2)°].



FIGURE 1. ORTEP drawing of TbtBi=BiTbt (1) with thermal ellipsoid plot (30% probability).

The Bi–Bi bond in 1 is 6% (0.169 Å) shorter than the Bi–Bi single bond length of 2.990(2) Å in Ph₂Bi–BiPh₂.^[17] This agrees reasonably well with the calculated bond shortening of 7% (0.214 Å) from H₂Bi–BiH₂ (3.009 Å) to HBi=BiH (2.795 Å).^[18] In addition, it is interesting that the bond shortenings are comparable with those reported for diphosphenes; the experimental value is 8% (0.183 Å) from (PhP)₅ [2.217(6) Å]^[19] to ArP=PAr [2.034(2) Å; Ar = 2,4,6-tri-*t*-butylphenyl]^[10], while the calculated value is 9% (0.200 Å) from H₂P–PH₂ (2.247 Å) to HP=PH (2.047 Å).^[18] Moreover, the bond shortening observed for 1 is in sharp contrast to the considerably

elongated Sn–Sn bond length in the isolated distance $R_2Sn=SnR_2$ [2.768(1) Å for R = CH(SiMe₃)₂^[8] and 2.910(1)Å for R = 2-*t*-butyl-4,5,6-trimethylphenyl^[20]], the bond lengths of which are almost equal to or longer than the typical Sn–Sn single bond length [*e. g.*, 2.780(4) Å for Ph₃Sn–SnPh₃].^[21]

The observed Bi-Bi-C angle of 100.5 ° deviates greatly from the ideal sp^2 hybridized bond angle (120 °). This is due to the fact that the heavy Bi atom has the lowest tendency to form a hybrid orbital because the size-difference of the valence s and p orbitals increases upon going from N to Bi (the significant 6s orbital contraction originates mostly from the relativistic effect) and prefers to maintain the $(6s)^2(6p)^3$ valence electron configuration. The use of these three orthogonal 6p orbitals without significant hybridization leads to a bond angle of *ca.* 90 ° at Bi.^[22] The Bi-Bi-C bond angle in 1, being close to 90 °, is exactly the experimental evidence for the core-like nature of the 6s electrons, *i. e.* so called "inert s-pair effect" or "non-hybridization effect".^[23]

Dibismuthene 1 is purple in hexane and shows two absorption maxima at $\lambda_1 = 660$ nm (sh, $\varepsilon 100$) and $\lambda_2 = 525$ nm ($\varepsilon 4000$), which correspond to the $n \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$ transitions of the Bi=Bi chromophore, respectively. These red-shifts relative to the values reported for diphosphenes^[10,24,25] and diarsenes^[11,26] agree with the changes in the n, π , and π^* orbital levels calculated for HE=EH (E = P, As, Sb, and Bi).^[22] Meanwhile, in the FT-Raman spectrum a strong band attributable to the Bi–Bi stretching was observed at 134 cm⁻¹ for the dibismuthene 1 (solid; excitation, He-Ne laser 632.8 nm). This is 31 cm⁻¹ higher than the Bi–Bi stretching frequency of Ph₂Bi–BiPh₂ (103 cm⁻¹),^[27] agreeing with the frequency shift of 34 cm⁻¹ calculated for HBi=BiH (153 cm⁻¹) and H₂Bi–BiH₂ (119 cm⁻¹).^[18] Thus, the spectroscopic data here obtained indicate again that 1 features double bonding between the Bi atoms even in solution.



Scheme 3

Finally, it should be noted that dibismuthene 1 undergoes a unique reaction with oxygen (Scheme 3). Dibismuthene 1 is considerably stable in the solid state, retaining its purple color for several hours on exposure to the open air, but it undergoes slow reaction with oxygen to give quantitatively the corresponding colorless 1,3,2,4-dioxadibismuthene derivative, (TbtBiO)₂ 4, the structure of which was determined by X-ray crystallographic analysis. In solution 1 reacts with oxygen quite rapidly to give 4.

The successful isolation of dibismuthene 1 is worthy of special note because the Bi–Bi double bond is the heaviest known and changes drastically our understanding of double-bond compounds in chemistry textbooks. Future work along this line may lead us to a whole new class of multiply bonded compounds.

ACKNOWLEDGMENTS This work was partially supported by a
Grant-in-Aid for Scientific Research on Priority Areas (No. 09239208) from the Ministry of Education, Science, Sports and Culture, Japan. We are grateful to Prof. Yukio Furukawa, the University of Tokyo, for the measurement of FT-Raman spectra. N. T. is thankful to Ciba-Geigy Foundation (Japan) for the Promotion of Science for the financial support. We are also grateful to Shin-etsu Chemical Co., Ltd. and Tosoh Akzo Co., Ltd. for the generous gift of chlorosilanes and alkyllithiums, respectively.

REFERENCES

- Multiply Bonded Main Group Metals and Metalloids in Advances in Organometallic Chemistry (Eds., F. G. A. Stone and R. West), 39 (1996).
- [2.] G. Raabe and J. Michl, in *The Chemistry of Organic Silicon Compounds Part 2* (Eds. S. Patai and Z. Rappoport), 1015-1142 (John Wiley & Sons, New York, 1989).
- [3.] J. Barrau, J. Escudié, and J. Satgé, Chem. Rev., 90, 283-319 (1990).
- [4.] T. Tsumuraya, S. A. Batcheller, and S. Masamune, Angew. Chem., Int. Ed. Engl., 30, 902-930 (1991).
- [5.] M. Yoshifuji, in Multiple Bonds and Low Coordination in Phosphorus Chemistry (Eds, M. Regitz and O. J. Scherer), 321-337 (Thieme, Stuttgart, 1990).
- [6.] R. West, M. J. Fink, and J. Michl, *Science*, **214**, 1343-1344 (1981).
- [7.] J. T. Snow, S. Murakami, S. Masamune and D. J. Williams, *Tetrahedron Lett.*, 25, 4191-4194 (1984).
- [8.] P. J. Davidson, D. H. Harris, and M. F. Lappert, J. Chem. Soc., Dalton Trans., 2268-2274 (1976).
- [9.] S. Masamune and L. Sita, J. Am. Chem. Soc., 107, 6390-6391 (1985).
- [10.] M. Yoshifuji, I. Shima, N. Inamoto, K. Hirotsu, and T. Higuchi, J. Am. Chem. Soc., 103, 4587-4589 (1981).
- [11.] A. H. Cowley, J. G. Lasch, N. C. Norman, and M. Pakulski, J. Am. Chem. Soc., 105, 5506-5507 (1983).
- [12.] H. Suzuki, N. Tokitoh, S. Nagase, and R. Okazaki, J. Am. Chem. Soc., 116, 11578-11579 (1994).
- [13.] N. Tokitoh, T. Matsumoto, K. Manmaru, and R. Okazaki, J. Am. Chem. Soc., 115, 8855-8856 (1993).
- [14.] T. Matsumoto, N. Tokitoh, and R. Okazaki, Angew. Chem., Int. Ed. Engl., 33, 2316-2317 (1994).
- [15.] N. Tokitoh, T. Matsumoto, and R. Okazaki, J. Am. Chem. Soc., 119, 2337 (1997).
- [16.] R. Okazaki, N. Tokitoh, and T. Matsumoto, in Synthetic Methods of Organometallic and Inorganic Chemistry (Ed by W. A. Herrmann), Vol. 2 (Vol. Eds; N. Auner and U. Klingebiel), Thieme, New York, 260-269 (1996).
- [17.] F. Calderazzo, R. Poli, and G. Pelizzi, J. Chem. Soc., Dalton Trans., 2365-2369 (1984).
- [18.] All the calculations here described were carried out at the QCISD/TZ(2d,p) level with Gaussian 94 program.

- [19.] J. J. Daly, J. Chem. Soc., 6147-6166 (1964).
- [20.] M. Weidenbruch, H. Kilian, K. Peters, H. G. Schnering, and H. Marsmann, H. Chem. Ber., 128, 983-985 (1995).
- [21.] v. H. Preut, H.-J. Haupt, and F. Huber, Z. Anorg. Allg. Chem., 396, 81-89 (1973).
- [22.] S. Nagase, S. Suzuki, and T. Kurakake, J. Chem. Soc., Chem. Commun., 1724-1726 (1990).
- [23.] S. Nagase, in The Chemistry of Organic Arsenic, Antimony and Bismuth Compounds, (Ed by Patai, S.) 1-24 (John Wiley & Sons, New York, 1994).
- [24.] M. Yoshifuji, K. Shibayama, and N. Inamoto, J. Am. Chem. Soc., 105, 2495-2497 (1983).
- [25.] A. H. Cowley, J. E. Kilduff, J. G. Lasch, S. K. Mehrotra, N. C. Norman, M. Pakulski, B. R. Whittlesey, J. L. Atwood, and W. E. Hunter, *Inorg. Chem.*, 23, 2582-2593 (1984).
- [26.] C. Couret, J. Escudié, Y. Madaule, H. Ranaivonjatovo, and J.-G. Wolf, *Tetrahedron Lett.* 24, 2769-2770 (1983).
- [27.] A. J. Ashe, III, E. G. Ludwig, Jr., and J. Oleksyszyn, Organometallics, 2, 1859-1866 (1983).