




that on irradiation **1** was quantitatively isomerized to 4-vinylgermacyclobutene **3**. Then, the irradiation of **3** formed in C₆D₆ containing excess amounts of methanol at 0 °C for 15 min gave a mixture composed of **1**, **3** and a methoxygermane in ca. 1:1:2 ratio. The structure for the methoxygermane was assigned to either **6** or **6'** on the basis of its NMR spectra.⁶⁾ Interestingly, on heating in benzene containing excess amounts of methanol at 45 °C for 5 min **3** reverted to **1** quantitatively without formation of **6** (**6'**). On the other hand, **1** in C₆D₆ containing methanol was irradiated for 1 min to give **6** (**6'**) in 70% yield accompanied by a trace amount of **3**. Since the germanium-carbon double bonded species are known to be highly reactive toward alcohol,⁷⁾ the formation of methoxygermane **6** (**6'**) could be rationalized by intermediary of **4**.^{2,8)} However, the results indicate that thermal isomerization of **3** to **1** may not proceed via **4** in contrast to the case of the silicon analog.²⁾

Diphenyl derivative **2** in degassed C₆D₆ in an NMR tube was irradiated under similar conditions for 45 h, but any noticeable spectral change was not recognized. However, irradiation of **2** for 10 min in C₆D₆ containing methanol readily gave **7** and **7'** in 10 and 39%, respectively.⁹⁾ These results indicate that 1-germahexatriene **5** should be generated and trapped efficiently by methanol. In the absence of methanol, **5** should revert thermally quite efficiently to the starting diene **2**. In contrast to the silicon analog,²⁾ the corresponding 2-germabicyclo[3.1.0]hexa-3-ene **8** was not detected in the photoproducts, the reason for which is not certain at this moment. Direct photochemical methanolysis of **1** and **2** leading to the corresponding methoxygermanes may be less probable but conceivable as the other possible reaction paths (Scheme).

1-Germa-2,4-cyclohexadienes **1** and **2** show their UV maxima at the same wavelength (330 nm), but the intensity of the maxima is much stronger for **2**. This implies that the germacyclohexadiene ring of **1** may be more sterically strained, and this may cause **1** to be more photochemically labile in benzene.

This research was supported in part by the Ministry of Education, Science, and Culture (Grant-in-Aid for Scientific Research on Priority Area of Reactive Organometallics).

References

- 1) W. G. Dauben, M. S. Kellog, J. I. Seeman, N. D. Vietmeyer, and P. H. Wendschuh, *Pure Appl. Chem.*, **33**, 197 (1973); A. Padwa, L. Brodsky, and S. Clough, *J. Am. Chem. Soc.*, **94**, 6767 (1972); S. W. Spangler and R. P. Hennis, *J. Chem. Soc., Chem. Commun.*, **1972**, 24.
- 2) Y. Nakadaira, S. Kanouchi, and H. Sakurai, *J. Am. Chem. Soc.*, **96**, 5621 (1974).
- 3) **1**: mp 136-142 °C; MS m/z 474 (M⁺, ⁷⁴Ge); ¹HNMR (C₆D₆) δ 0.36 (s, 6H), 2.33 (s, 2H), 6.95 (m, 20H); UV (cyclohexane) λ_{max} 330 nm (ε 7760).
- 4) **2**: an oil; MS m/z 322 (M⁺, ⁷⁴Ge); ¹HNMR (C₆D₆) δ 0.40 (s, 6H), 2.02 (d, J=1.8 Hz, 2H), 6.46 (dt, J=6.7 and 1.8 Hz, 1H), 6.92 (d, J=6.7 Hz, 1H), 7.3 (m, 10H); UV (cyclohexane) λ_{max} 330 nm (ε 21000).
- 5) **3**: MS m/z 474 (M⁺, ⁷⁴Ge); ¹HNMR (C₆D₆) δ 0.38 (s, 3H), 0.62 (s, 3H), 5.53 (d, J=0.8 Hz, 1H), 5.71 (d, J=0.8 Hz, 1H), 7.1 (m, 20H).
- 6) **6** (**6'**): MS m/z 475 (M⁺-OMe, ⁷⁴Ge), 371 (M⁺-GeMe₂OMe, 100%); ¹HNMR (C₆D₆) δ 0.28 (s, 3H), 0.37 (s, 3H), 3.36 (s, 3H), 4.66 (s, 1H), 5.34 (d, J<0.1 Hz, 1H), 5.77 (d, J>0.1 Hz, 1H). The signal at δ 4.66 was missing on irradiation in the presence of methanol-d₁.
- 7) J. Satgé, *Adv. Organomet. Chem.*, **21**, 241 (1982).
- 8) The structure **6** having a hydrogen at C₃ suggests that the direct photo-induced methanolysis of **3** may be another possible route to **6** as indicated in Scheme.
- 9) **7**: MS m/z 354 (M⁺, ⁷⁴Ge); ¹HNMR (C₆D₆) δ 0.29 (s, 6H), 3.39 (s, 3H), 3.28 (d, J=7.0 Hz, 2H), 5.13 (q, J=1.6 Hz, 1H), 5.37 (dt, J=1.6 and 0.8 Hz), 6.18 (t, J=7.0 Hz, 1H), 7.19 (m, 10H). **7'**: MS m/z 354 (M⁺, ⁷⁴Ge); ¹HNMR (C₆D₆) δ 0.18 (s, 3H), 0.21 (s, 3H), 3.33 (s, 3H), 3.88 (d, J=12.3 Hz, 1H), 5.31 (dd, J=2.0 and 2.1 Hz, 1H), 5.50 (dd, J=2.0 and -1 Hz, 1H), 6.14 (dd, J=12.3 and 11.3 Hz, 1H), 6.26 (ddd, J=11.6, 2.0, and -1 Hz, 1H), 7.19 (m, 10H). On the NMR spectra of **7** and **7'** obtained on irradiation in the presence of methanol-d₁, the signal at δ 3.28 was reduced to one-half, and that at δ 3.88 was almost missing.

(Received July 21, 1994)