

Acta Cryst. (1999). C55, 1814–1815

The first amidozirconium(IV)–porphyrin complex: *cis*-[bis(trimethylsilyl)amido]chloro(tetraphenylporphyrinato)zirconium(IV)

HEE-JOON KIM AND KIMOON KIM

*Department of Chemistry, Pohang University of Science and Technology, San 31, Hyojadong, Pohang 790-784, Korea.
E-mail: kkim@postech.ac.kr*


(Received 3 June 1999; accepted 20 July 1999)

Abstract

The title compound, [ZrCl(C₄₄H₂₈N₄)(C₆H₁₈NSi₂)], is the first amidozirconium(IV)–porphyrin complex and it adopts an out-of-plane *cis* structure with the Zr atom in a distorted trigonal prismatic coordination geometry.

Comment

Organometallic early transition metal porphyrin complexes have attracted much attention in recent years due to their unusual structures and reactivities (Brand & Arnold, 1995). In the course of our ongoing research on organometallic zirconium–porphyrin complexes (Kim *et al.*, 1993, 1997, 1999), we have synthesized the title complex, *cis*-[bis(trimethylsilyl)amido]chloro(tetraphenylporphyrinato)zirconium(IV), (I), and determined its crystal structure. To our knowledge, complex (I) is the first example of an amidozirconium(IV)–porphyrin complex.



Complex (I) adopts an out-of-plane *cis* structure in which the Zr atom is displaced by 1.0264 (7) Å from the mean N₄ plane of the porphyrin ligand and the Cl—Zr—N5 angle is 81.33 (4)°. The coordination geometry of the Zr centre is best described as a distorted trigonal prism. The plane defined by atoms Cl, N5 and Zr bisects two of the N—Zr—N angles (N1—Zr—N4 and N2—Zr—N4). The distance between the Zr and amide N atoms [Zr—N5 2.1009 (15) Å] is similar to the average value of 2.112 Å found in *cis*-[(NMe₂)₂Zr(Me₄taen)] (Me₄taen is the 5,7,12,14-tetramethyl-1,4,8,11-tetraazacyclotetradeca-4,6,11,13-tetraenate dianion; Black *et al.*, 1995). As observed in *cis*-[(NMe₂)₂Zr(Me₄taen)], the amide N

atom is nearly flat (the sum of the angles around N5 is 356.28°), which indicates the presence of π -bonding between the Zr and amide N atoms. The steric bulk around N5 may also contribute to its planarity. The Zr—Cl distance of 2.5112 (5) Å is significantly longer than those in *cis*-[Cl₂(thf)Zr(tp)] [2.473 (1) and 2.499 (2) Å; thf is tetrahydrofuran and tpp is the 5,10,15,20-tetraphenylporphyrinate dianion; Kim *et al.*, 1993] or those in *cis*-[Cl₂Zr(oep)] [2.473 (1) and 2.459 (1) Å; oep is the 2,3,7,8,12,13,17,18-octaethylporphyrinate dianion; Brand & Arnold, 1993]. The Zr—N_{porphyrin} distances range from 2.2465 (14) to 2.2948 (15) Å, with an average value of 2.266 (3) Å. Interestingly, at 2.2948 (15) Å the Zr—N2 bond is considerably longer than the others, which may be due in part to the *trans* influence of the amide ligand, although the Zr—N2 bond is not perfectly *trans* to the Zr—N5 bond [N2—Zr—N5 140.12 (6)°].




Fig. 1. The molecular structure of (I). Displacement ellipsoids are shown at the 50% probability level and H atoms have been omitted for clarity.

Experimental

For the preparation of (I), a toluene solution (*ca* 2 ml) of LiN(SiMe₃)₂ (75 mg, 0.43 mmol) was added slowly to a suspension of *cis*-[Cl₂Zr(tp)] (263 mg, 0.272 mmol) in toluene (*ca* 20 ml). The reaction mixture was stirred at room temperature for 12 h and then filtered. The filtrate was concentrated under reduced pressure and pentane was layered over the solution to give (I) as dark purple X-ray quality crystals in 95% yield. ¹H NMR (300 MHz, CDCl₃): δ 8.99 (*s*, 8H, H_{pyrrole}), 8.65 (*bs*, 4H, *o*-Ph), 7.83–7.71 (*br, m*, 16H, Ph), –1.51 (*s*, 18H, CH₃); ¹³C NMR (CDCl₃): δ 152.1, 141.9, 134.7, 134.1, 132.3, 128.7, 127.4, 126.8, 3.06.

Crystal data

[ZrCl(C₄₄H₂₈N₄)(C₆H₁₈NSi₂)] Mo K α radiation
M_r = 899.77 λ = 0.71073 Å

Triclinic

 $P\bar{1}$ $a = 11.9557(7) \text{ \AA}$ $b = 12.1110(7) \text{ \AA}$ $c = 17.931(1) \text{ \AA}$ $\alpha = 99.790(1)^\circ$ $\beta = 98.943(1)^\circ$ $\gamma = 114.365(1)^\circ$ $V = 2255.9(2) \text{ \AA}^3$ $Z = 2$ $D_x = 1.325 \text{ Mg m}^{-3}$ D_m not measured

Cell parameters from 5599 reflections

 $\theta = 2.6\text{--}28.3^\circ$ $\mu = 0.396 \text{ mm}^{-1}$ $T = 188(2) \text{ K}$

Block

 $0.4 \times 0.3 \times 0.3 \text{ mm}$

Purple

Science and Engineering Foundation through the Center for Biofunctional Molecules for support of this work. We also thank Dr Dongmok Whang for help with the X-ray work.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: BM1353). Services for accessing these data are described at the back of the journal.

Data collection

Siemens SMART CCD area-detector diffractometer

 φ and ω scans

Absorption correction:

multi-scan (SADABS;

Sheldrick, 1997a)

 $T_{\min} = 0.873$, $T_{\max} = 0.976$

14 640 measured reflections

10 520 independent

reflections

9504 reflections with

 $I > 2\sigma(I)$ $R_{\text{int}} = 0.016$ $\theta_{\max} = 28.33^\circ$ $h = -15 \rightarrow 15$ $k = -13 \rightarrow 16$ $l = -23 \rightarrow 23$

Refinement

Refinement on F^2 $R[F^2 > 2\sigma(F^2)] = 0.032$ $wR(F^2) = 0.081$ $S = 1.104$

10 520 reflections

532 parameters

H atoms constrained

 $w = 1/[\sigma^2(F_o^2) + (0.0227P)^2 + 1.9648P]$ where $P = (F_o^2 + 2F_c^2)/3$ $(\Delta/\sigma)_{\max} = 0.04$ $\Delta\rho_{\max} = 0.40 \text{ e \AA}^{-3}$ $\Delta\rho_{\min} = -0.47 \text{ e \AA}^{-3}$

Extinction correction: none

Scattering factors from

International Tables for

Crystallography (Vol. C)

Table 1. Selected geometric parameters ($\text{\AA}$, $^\circ$)

Zr—Cl	2.5112 (5)	Zr—N3	2.2631 (15)
Zr—N1	2.2465 (14)	Zr—N4	2.2608 (15)
Zr—N2	2.2948 (15)	Zr—N5	2.1009 (15)
Cl—Zr—N1	135.15 (4)	N2—Zr—N3	79.25 (5)
Cl—Zr—N2	79.27 (4)	N2—Zr—N4	125.85 (5)
Cl—Zr—N3	84.21 (4)	N2—Zr—N5	140.12 (6)
Cl—Zr—N4	143.71 (4)	N3—Zr—N4	76.87 (5)
Cl—Zr—N5	81.33 (4)	N3—Zr—N5	132.77 (6)
N1—Zr—N2	76.37 (5)	N4—Zr—N5	89.13 (6)
N1—Zr—N3	126.41 (5)	Si1—N5—Si2	114.59 (9)
N1—Zr—N4	80.15 (5)	Si1—N5—Zr	122.04 (8)
N1—Zr—N5	94.06 (6)	Si2—N5—Zr	119.65 (8)

The crystal showed no significant decay and thus no correction was applied for decay. The structure was solved by direct methods. H atoms were added at calculated positions and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$.

Data collection: SMART (Siemens, 1996). Cell refinement: SAINT (Siemens, 1996). Data reduction: SAINT. Program(s) used to solve structure: SHELXTL (Sheldrick, 1997b). Program(s) used to refine structure: SHELXTL. Molecular graphics: SHELXTL. Software used to prepare material for publication: SHELXTL.

The authors thank the Korea Research Foundation (Non-Directed Research Fund, 1996) and the Korea

References

- Black, D. G., Swenson, D. C., Jordan, R. F. & Rogers, R. D. (1995). *Organometallics*, **14**, 3539–3550.
- Brand, H. & Arnold, J. (1993). *Organometallics*, **12**, 3655–3665.
- Brand, H. & Arnold, J. (1995). *Coord. Chem. Rev.* **140**, 137–168.
- Kim, H.-J., Jung, S., Jeon, J.-M., Whang, D. & Kim, K. (1997). *Chem. Commun.* pp. 2201–2202.
- Kim, H.-J., Jung, S., Jeon, J.-M., Whang, D. & Kim, K. (1999). *Chem. Commun.* pp. 1033–1034.
- Kim, H.-J., Whang, D., Kim, K. & Do, Y. (1993). *Inorg. Chem.* **32**, 360–362.
- Sheldrick, G. M. (1997a). *SADABS. Program for Semi-Empirical Absorption Correction*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997b). *SHELXTL. Structure Determination Programs*. Version 5.03. Bruker AXS Inc., Madison, Wisconsin, USA.
- Siemens (1996). *SMART and SAINT. Area Detector Control and Integration Software*. Version 4.0. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1999). **C55**, 1815–1817

Tetraaqua(1,10-phenanthroline)zinc(II) sulfate dihydrate

CUNGEN ZHANG,^a KAIBEI YU,^b DAN WU^a AND CHENGXUE ZHAO^a

^aChemistry Department, Shanghai Jiaotong University, Shanghai 200240, People's Republic of China, and ^bAnalysis Center, Chengdu Branch of Chinese Academy of Science, Chengdu 610041, People's Republic of China. E-mail: zcungen@kali.com.cn

(Received 12 April 1999; accepted 6 July 1999)

Abstract

A novel monomeric phenanthroline (phen) zinc complex, $[\text{Zn}(\text{C}_{12}\text{H}_8\text{N}_2)(\text{H}_2\text{O})_4]\text{SO}_4 \cdot 2\text{H}_2\text{O}$, has been characterized by X-ray analysis. The Zn atom is coordinated in a slightly distorted octahedral geometry by two N atoms from 1,10-phenanthroline and four O atoms from water ligands. The Zn—N bond lengths are in the range 2.136(3)–2.156(2) $\text{\AA}$ and the Zn—O bond lengths are in the range 2.083(2)–2.130(2) $\text{\AA}$. Hydrogen bonds stabilize the structure.