ATMOSPHERE DEPENDENT PHOTOREACTION OF METHYL 4-PYRIDINECARBOXYLATE IN METHANOL. METHOXYLATION AT 3-POSITION UNDER 0, AND METHOXYLATION AND HYDROXYMETHYLATION AT 2-POSITION UNDER N2

Toru SUGIYAMA\*, Kazuya YAGI, Yoko ITO, and Akira SUGIMORI\* Department of Chemistry, Faculty of Science and Technology, Sophia University, Kioi-cho 7-1, Chiyoda-ku, Tokyo 102

The UV-Irradiation of methyl 4-pyridinecarboxylate in methanol under oxygen in the presence of  $H_2SO_4$  brings about methoxylation at the 3-position of the pyridine ring, while under nitrogen methoxylation and hydroxymethylation occur at the 2-position.

In previous papers  $^{1-4}$ ) we have reported pronounced photoreactivities of 2and 3-pyridinecarboxylic esters: parallel occurrence of radical reaction (alkylation) and ionic reaction (alkoxylation), different paths to give the same product, and the coexistence of several different excited states of the same multiplicity. The photoreactions of pyridinecarboxylic esters depend on the reaction conditions, especially, the acidity of the solutions and polarity of the solvent alcohol.

We report here a remarkable atmosphere dependence of photoreactions of methyl 4-pyridinecarboxylate(1): the presence of oxygen changes dramatically the position of methoxylation.

The irradiation of methyl 4-pyridinecarboxylate (nomally 50  ${\rm cm}^3$  solution containing 0.02 mol dm<sup>-3</sup> of 1) in the presence of  $H_2SO_4$  in methanol under oxygen atmosphere with a low pressure mercury lamp for 25 min at 30°C gave methyl 3methoxy-4-pyridinecarboxylate(2)<sup>5)</sup>quantitatively (Table 1).





Table 1. Photoreactions of methyl 4-pyridinecarboxylate( 1 ) in acidic methanolic solutions. [1] = 0.02 mol dm<sup>-3</sup>; Irradiation time, 25 min; Temp., 30°C.

[H <sub>2</sub> SO <sub>4</sub> ]	Atmosphere	Conversion of $l$	(Yi	(Yield of product) <sup>a)</sup> /%			
$mol dm^{-3}$		e	2	3	<b>4</b> ~	5	
0.25	°2	14	88	trace	-	_	
0.25	N <sub>2</sub>	47.5	trace	14	5.5	trace	

a) Yields were determined by means of GLC and/or HPLC, and were calculated on the basis of 1 consumed.

On the other hand, in the irradiation under nitrogen atmosphere, methyl 2methoxy-4-pyridinecarboxylate(3),<sup>5</sup> methyl 2-hydroxymethyl-4-pyridinecarboxylate (4),<sup>5</sup> and a trace of 3,3'-dimer(5)<sup>5</sup> were obtained.

The photoreaction of 1 is characterized by the fact that the presence of oxygen causes a dramatic change of photoreaction. Such a remarkable oxygen effect was not observed in the photoreactions of 2- and 3-pyridinecarboxylic esters.

The effects of anthracene on the two methoxylations are different, as shown in the Stern-Volmer plots (Figs. 1 and 2).<sup>6)</sup> These figures indicate that methoxylation under  $N_2$  is inhibited by anthracene effectively, while that under  $O_2$  is not inhibited by anthracene at all.

Although the photo-methoxylation at the 2-position under  $N_2$  is inhibited by anthracene, benzene, and triphenylene, it is slightly affected by *trans*-1,3-pentadiene (Table 2). The inhibition of photo-methoxylation under  $N_2$  by aromatic hydrocarbons could be explained not by the triplet energy transfer, but by an electron transfer from the aromatic hydrocarbons to the excited 1. A similar result has been obtained in the photo-methoxylation of methyl 2-pyridinecarboxylate.<sup>3)</sup> Thus, methoxylation under  $N_2$  should originates from a singlet state.

For the photo-methoxylation at the 3-position under O<sub>2</sub>, the contribution of an comlex between 1 and oxygen can be postulated because the absorbance of the 1 - O<sub>2</sub> system in 235 - 255 nm region is greater than that of 1 under N<sub>2</sub> in CH<sub>3</sub>CN. A feature of the photoreaction is that oxygen promotes an ionic reaction. A few reports on the similar promotion of photochemical ionic reactions by oxygen have appeared except for alcoholyses of benzotrichloride<sup>7)</sup> and for alcoholyses of trihaloacetyl groups.<sup>8)</sup>



Fig. 1. Stern-Volmer plot for an effect of anthracene on the photoreaction of 1 in methanol under  $O_2$  atmosphere.



Fig. 2. Stern-Volmer plots for effects
of anthracene on the photoreaction of
l in methanol under N<sub>2</sub> atmosphere.
[ l ] = 0.01 mol dm<sup>-3</sup>, [H<sub>2</sub>SO<sub>4</sub>] = 0.05
mol dm<sup>-3</sup>; Irradiation time, 1 h;

Temp., 20°C.

-O-: 3, -<u>A</u>-: 4

Atmospher	Additive	[Additive]	Ratio of light	Relative Yield(=A) <sup>6)</sup>	
		mol dm <sup>-3</sup>	absorbed by $\frac{1}{2}$ (=L) <sup>6</sup> )	2	3
°2	<i>trans-</i> 1,3- Pentadiene	$4 \times 10^{-4}$	1.00	1.24	-
N <sub>2</sub>	<i>trans-</i> 1,3- Pentadiene	$4 \times 10^{-4}$	1.00	-	0.90
N <sub>2</sub>	Benzene	$1 \times 10^{-1}$	0.69	-	0.96
<sup>N</sup> 2	Triphenylene	$3 \times 10^{-4}$	0.69	-	0.32

Table 2. Effects of additives on the photo-methoxylation of 1

 $[1] = 1 \times 10^{-2} \text{mol dm}^{-3}; [H_2SO_4] = 5 \times 10^{-2} \text{mol dm}^{-3};$ Irradiation time, 1 h; Temp., 20°C. References

- 1) A.Sugimori, Yuki Gosei Kagaku Kyokai Shi, 39, 712 (1981).
- T.Sugiyama, E.Tobita, K.Takagi, M.Sato, Y.Kumagai, G.P.Satô, and A.Sugimori, Chem. Lett., 1980, 131.
- T.Sugiyama, T.Furihata, K.Takagi, M.Sato, S.Akiyama, G.P.Satô, and A.Sugimori, Bull. Chem. Soc. Jpn., 54, 3785 (1981).
- A.Sugimori, E.Tobita, Y.Kumagai, and G.P.Satô, Bull. Chem. Soc. Jpn., <u>54</u>, 1761 (1981).
- 5) 2: mp, 55-56.5°C; IR(KBr), 1720(ester C=O) and  $1220 \text{ cm}^{-1}(\text{C-O-C})$ ; <sup>1</sup>H NMR(CDCl<sub>3</sub>)  $\delta = 8.44(1\text{H},\text{s}), 8.29(1\text{H},\text{d},J=4\text{Hz}), 7.55(1\text{H},\text{d},J=4\text{Hz}), 4.02(3\text{H},\text{s}) \text{ and } 3.91(3\text{H},\text{s})$ ; MS(70eV), m/z (rel intensity) 167(M<sup>+</sup>;77), 152(14), 138(18), 136(100), 135(18), 133(31), Found: m/z 167.0585. Calcd for C<sub>8</sub>H<sub>9</sub>NO<sub>3</sub>:M, 167.0583.

3: mp, 33-34.5°C;<sup>1</sup>H NMR spectra are in accord with those obtained by Deady et al. (L.W.Deady, P.M.Harrison, and R.D.Topsom, *Org. Magn. Reson.*, <u>7</u>, 41 (1975); L.W.Deady, private communication).

4: mp, 67.5-68°C (lit,68-69°C); The spectral data are in accord with those obtained by Ninomiya (I.Ninomiya, private communication).

5: mp, 184-187°C(sub); IR(KBr),  $1720 \text{ cm}^{-1}$ (ester C=O); <sup>1</sup>H NMR(CDCl<sub>3</sub>)  $\delta$  = 8.95 (1H,d,J=1.6Hz), 8.85(1H,d,J=5Hz), 7.87(1H,dd,J=1.6 and 5Hz), and 4.00(3H,s); Found: C, 61.7; H, 4.7; N, 10.2 %; M<sup>+</sup>, 272. Calcd for C<sub>14</sub>H<sub>12</sub>N<sub>2</sub>O<sub>4</sub>: C, 61.76; H, 4.44; N, 10.29 %; M, 272.

- 6) In this case, the effects of additives were analyzed by means of an index A, the relative quantum yield calculated on the basis of the light absorbed by 1, because the additives absorb the 254 nm light competitively with 1. A<sup>-1</sup> corresponds to  $\phi_0/\phi$  in the Stern-Volmer plots for the quenching experiment. A = Y/L, Y = (Yield in the presence of additive(Q))/(Yield in the absence of additive(Q)), L =  $\varepsilon_1 [\frac{1}{2}]/(\varepsilon_1 [\frac{1}{2}] + \varepsilon_0 [Q])$ , where  $\varepsilon_1$  and  $\varepsilon_0$  are the molar absorption coefficients of 1 and Q at 254 nm, respectively.
- 7) T.Ishigami, Y.Kinoshita, and A.Sugimori, Chem. Lett., 1974, 149.
- Y.Izawa, H.Tomioka, M.Natsume, S.Beppu, and H.Tsuji, J. Org. Chem., <u>45</u>, 4835 (1980).

(Received March 24, 1982)

920