

S0040-4039(96)00081-0

Deprotection of mono and dimethoxy phenyl methyl ethers using catalytic amounts of DDQ

S.Chandrasekhar*, G.Sumithra and J.S. Yadav

Indian Institute of Chemical Technology, Hyderabad-500 007, India

Abstract: 4-Methoxy and 3,4 dimethoxy benzyl ethers have been deprotected with catalytic amounts of DDQ by oxidative recycling of the byproduct DDHQ with FeCl for the first time.

Protective groups play an essential role in synthetic strategies¹ and easy removal of a particular protective group in the presence of other labile protective groups and sensitive functional groups always determines its usefulness. Oxidative methods for removing protecting groups are relatively rare and 4-methoxy and 3,4 dimethoxy benzyl (MPM and DMPM) ethers are lately added to the repertoire of standard protective groups. These undergo easy electron transfer to 2,3 dichloro-3,6-dicyano-1,4-benzoquinone (DDQ) to liberate the free alcohols under essentially neutral conditions². However the DDHQ (dichloro dicyano hydroquinone) is an inevitable byproduct, removal of which is sometimes cumbersome, besides which stoichiometric DDQ is required which is expensive. To overcome these problems, herein we report an altogether novel DDQ 'regeneration technique' for the removal of said protective groups.



Thus treatment of n-decyl MPM ether (table 1, entry 1) and DMPM ether (entry 2) with 10 mole percent of DDQ and 3 equivalents of FeCl₉, a reagent generally used for oxidation of hydroquinones to quinones³ in DCM:H₂O (10:1) furnished cleanly the parent alcohols in excellent yields (Table 1). To test the generality and selectivity under the reaction conditions, 1,10-decanediol was differentially protected as MPM ether and acetate(entry 3) or THP ether (entry 4) or TBDPS ether (entry 5) or benzyl ether (entry 6) or methoxy methyl ether (entry 7) and subjected to described conditions produced consistently high yields of deprotected alcohols without effecting the other protective groups. The products were isolated easily from the other byproduct using a simple column chromatography.

IICT Communication No. 3610

Cholesteryl-OMPM (entry 8) and sugar derivative (entry 9) also cleanly underwent the desired deprotection. However with FeCL alone starting materials were recovered.



Table 1

* About 10% THP ether cleavage was noticed at the end of 3h

Thus an altogether new strategy for the removal of MPM and DMPM ethers using catalytic amounts of DDQ was developed which is simple, easy, economical and selective.

Acknowledgement : One of us (GS) is thankful to CSIR, New Delhi for financial assistance.

References:

- Greene, T.W.; Wuts, P.G.M. Protective Groups in Organic Synthesis, 2nd ed.; Wiley: New 1. (a) York, 1991.
 - Kocienski, P.J. Protecting Groups. Georg Thiene Verlag Stuttgart. New York, 1994. (b)
- Oikawa, Y.; Yoshioka, T.; Yomenitsu, O. Tetrahedron Lett. 1982, 23, 885-888. 2. (a)
 - Srikrishna, A.; Viswajanani, R.; Sattigeri, J.A.; Vijaykumar, D. J. Org. Chem. 1995, 60, 5961-(b) 5962
 - Yan, L.; Kahne, D.Synlett. 1995, 523. (c)
- Alcalay, W. Helv. Chim. Acta. 1947, 30, 578. 3.

(Received in UK 8 November 1995; revised 11 January 1996; accepted 12 January 1996)