

Pergamon

0957-4166(95)00087-9

An Asymmetric Route to the 5,6-Dihydro-4H-1,3-Thiazine Skeleton via an Asymmetric Hetero Diels-Alder Reaction

Anne Marchand, Dominique Mauger, André Guingant* and Jean-Paul Pradère

Laboratoire de Synthèse Organique, Faculté des Sciences et des Techniques, U.R.A au C.N.R.S N°475, 2 rue de la Houssinière, 44072 NANTES Cedex 03, France

Abstract: Magnesium bromide mediated hetero Diels-Alder cycloaddition of diene 1 ($\mathbb{R}^1 = \mathbb{P}h$) to chiral N-acryloyl-2-oxazolidinone 4b leads to 5b as the sole adduct formed. Chiral auxiliary removal leads to enantiomerically pure 5,6-dihydro-4H-1,3-thiazine 3 ($\mathbb{R}^2 = \mathbb{B}n$). The selectivity of the cycloaddition is reversed under thermal or high pressure activation. The predominant diastereomeric adduct 6b is isolated in pure form after flash chromatography.

For some years, studies in our laboratory have focused on the construction of the dihydro-1,3-thiazine ring, a subunit found in biologically active natural products (eg. cephamycin C) as well as in many unnatural compounds displaying a broad range of activities (medicinal drugs, herbicids, pesticids, etc). In particular it has been shown that the hetero Diels-Alder reaction of 3-aza-4-dimethylamino-1-thia-butadienes 1 with acrylates 2 induced by heating¹ or by application of high pressure² provides a convenient and efficient route to substituted 5,6-dihydro-4H-1,3-thiazines 3, allowing for the simultaneous construction of carbon-carbon and sulfur-carbon bonds and installation of two contiguous stereogenic centers (Scheme 1).



Scheme 1

In continuation of this work we have now considered the possibility that acrylates bearing a chiral substituent might provide ready access to optically active 3 and we report herein our preliminary results which have culminated with the first asymmetric synthesis of the substituted 5,6-dihydro-4H-1,3-thiazine 3 in the case where $R^1 = Ph$ and $R^2 = Bn$.

The chiral dienophiles selected for study were the known acrylate 4a and N-acryloyl-2-oxazolidinone 4b. Although both of them have been widely used in asymmetric Diels-Alder reactions, there is no report of their use in hetero Diels-Alder reactions involving an heterodiene³. The cycloaddition of acrylate 4a with cyclopentadiene has been previously reported by Helmchen et al⁴. These authors have observed a remarkable reversal of selectivity of the addition process depending on the Lewis acid utilized (TiCl₄ versus EtAlCl₂).

A. MARCHAND et al.

With this result in mind we first studied the reaction of heterodiene $1 (R^1 = Ph)$ with acrylate 4a in the presence of Lewis acid catalysts. As can be seen from the data in Table 1 such a dichotomy is no longer found in the present reaction.



 Table 1 : Hetero Diels-Alder reaction of diene 1 with acrylate 4a and N-acryloyl-2-oxazolidinone 4b.

 Entry
 Dienophile
 Functionantal Conditionantal
 Yield %

Entry		Experiment		ntal Conditions ^a		Yield %	Diastereoselectionb
		none		temp.(°C)	time (h)		5/6
1	4a			20	3	72	50/50
2	4a	TiCl ₄	(1)	-78 to -20	3	80	60/40
3	4 a	TiCl ₂ (OiPr) ₂	(2)	-78 to -20	3	58	65/35
4	4a	EtAlCl ₂	(1)	-78 to -30	3	86	65/35
5	4 a	Et ₂ AlCl	(1)	-20	3	90	70/30
6	4 a	Et ₂ AlCl	(1)	-60	3	90	70/30
7	4a	MgBr ₂	(1.5)	20	1.5	90	65/35
8	4 b	none		110	20	75	20/80
9	4 b	none (10Kbar)		20	40	95	16/84
10	4b	MgBr ₂	(1)	0	3	95	100/0

^aLewis acid catalyzed reactions were conducted in CH₂Cl₂ with molar ratio of diene:dienophile = 1:1.5; Reaction under thermal activation was conducted in toluene with molar ratio of diene:dienophile = 1:1.5; Reaction under high pressure activation was conducted in THF with molar ratio of diene:dienophile = 1:1. ^bRatios of diastereomers were derived by integration of the NMe₂ peaks in the¹H NMR spectra (400MHz) of the unpurified cycloaddition products.

In all cases examined the same adduct is predominant in the diastereomeric mixture. Also in sharp contrast with the Helmchen results, the diastereoselection is not very high (70/30 at best, absence of diastereoselection in the uncatalyzed reaction).

Such disappointing results prompted us next to turn our attention to the reaction of heterodiene 1 with Nacryloyl-2-oxazolidinone 4b. As can be seen from entry 10 *the reaction catalyzed by magnesium bromide in* CH_2Cl_2 as the solvent led to a single diastereomer 5b as judged by NMR analysis. It is worthy of note that other Lewis acid such as TiCl₄ and AlEt₂Cl, which are efficient in all carbon Diels-Alder reactions involving dienophile 4b⁵, failed to promote the present cycloaddition. Also of interest is the reversal of selectivity and the significant diastereoselection observed under thermal (entry 8) and high pressure activation (entry 9). The mixture of diastereomers formed in the latter case was easily separated by column chromatography to give a pure sample of 6b. Thus, depending upon the activation mode employed (Lewis acid or high pressure), either 5b or 6b could be obtained in optically pure form.

The NMR coupling constants of the protons at C-4 and C-5 (10.1Hz) provide a simple means for establishing the relative *trans* configuration of the two newly created stereogenic centers in adducts $5b^6$ and $6b^7$. The absolute configuration could be established as being 4R,5R in diastereomer 5b by a single crystal X-ray⁸ and is in accordance with the generally observed bias of N-acryloyl-2-oxazolidinone 4b to react on the less hindered si face⁵, here via a magnesium chelated form 7, and a cyclocondensation process occurring through an exo topology (Scheme 3). The exo addition mode is consistent with previous results in achiral series^{1,2,9} and seems to be a general property of dienes 1.



Finally chiral auxiliary removal was effected without epimerization^{10,11} by treatment of adduct **5b** with lithium benzyloxyde (1.5 eq, -78°c to 0°C, THF) to give the 5,6-dihydro-4*H*-1,3-thiazine **3** in 40% yield¹².

In summary, the magnesium bromide catalyzed hetero Diels-Alder addition of diene 1 ($R^1 = Ph$) to chiral N-acryloyl-2-oxazolidinone 4b can be carried out with a very high level of diastereoselection to give, after chiral auxiliary removal, the substituted (4R,5R)-5,6-dihydro-4H-1,3-thiazine 3 which may serve as useful template for stereoselective transformations. Additional studies in order to expand the scope of this reaction to other heterodienes are currently under investigations.

Acknowledgments

We thank the "Conseil Général de Loire Atlantique" for a pre-doctoral fellowship ("prix atlanthèse") to AM.

References and Notes

- (a) Meslin, J.C.; Quiniou, H. Bull. Soc. Chim. Fr. 1979, II, 347-349. (b) Chehna, M.; Pradère, J.P.;
 Quiniou, H.; Le Botlan, D.; Toupet, L. Phosphorus Sulfur and Silicon, 1989, 42, 15-19.
- (a) Chehna, M.; Pradère, J.P.; Guingant, A. Synthetic Comm. 1987, 17, 1971-1976. (b) Cellerin,C.;
 Gokou Tea, C.; Pradère, J.P.; Guingant, A.; Guénot, P. Sulfur Lett. 1988, 8, 205-209.
- 3 For a recent review on asymmetric hetero Diels-Alder reactions, see: Waldmann, H. Synthesis, **1994**, 535-551.
- Poll, T.; Metter, J.O.; Helmchen, G. Angew. Chem. Int. Ed. Engl. 1985, 24, 112-114. See also: Hamley, P.; Helmchen, G.; Holmes, A.B.; Marshall, D.R.; MacKinnon, J.W.M.; Smith, D.F.; Ziller, J.W. J. Chem.; Chem.Commun. 1992, 786-788.
- 5 Evans, D.A.; Chapman, K.T.; Bisaha, J. J. Am. Chem. Soc. 1988, 110, 1238-1256.
- 5b: white solid, mp 125-126°C (diethylether-petroleum ether); [α]_D = -65.4 (c = 0.52, CHCl₃);
 IR (KBr): 1762, 1692, 1616 cm⁻¹; ¹H NMR (400 Mz,CDCl₃, δ): 2.47 (s, 6H), 2.88 (dd, J = 8.6 Hz, 13.7 Hz, 1H), 3.10 (dd, J = 3.7 Hz, 12.2 Hz, 1H), 3.26 (dd, J = 3.2 Hz, 13.7 Hz, 1H), 3.55 (dd, J = 12.2 Hz, 12.2 Hz, 1H), 4.20 (m, 2H), 4.38 (ddd, J = 3.7 Hz, 10.1 Hz, 12.2 Hz, 1H), 4.69 (d, J = 10.1 Hz, 1H), 4.79 (m, 1H), 7.35, (m, 8H), 7.85 (m, 2H); ¹³C (100MHz, CDCl₃, δ): 28.3, 36.7, 37.4, 40.7, 55.3, 65.9, 80.5, 126.7, 127.4, 128.4, 129.0, 129.8, 130.8, 135.3, 138.6, 153.5, 157.5, 174.9; MS: m/e = 423 (M⁺), 192, 159, 121, 55, 44.
- 7 **6b**: white solid, mp 107-108°C (diethylether-petroleum ether); $[\alpha]_D = +196.4$ (c = 0.28, CHCl₃); ¹H NMR (400 Mz,CDCl₃, δ): 2.41 (s,6H), 2.82 (dd, J = 8.6 Hz, 13.7 Hz, 1H), 3.14 (dd, J = 3.7 Hz, 12.2 Hz, 1H), 3.30 (dd, J = 3.2 Hz, 13.7 Hz, 1H), 3.47 (dd, J = 12.2 Hz, 12.2 Hz, 1H), 4.24 (m, 2H), 4.49 (ddd, J = 3.7 Hz, 10.1 Hz, 12.2 Hz, 1H), 4.71 (d, J = 10.1 Hz, 1H), 4.79 (m, 1H), 7.32 (m, 8H), 7.85 (m, 2H).
- 8 Toupet, L. Université de Rennes. Details of the crystal structure of **5b** will be reported in a forthcoming full paper.
- 9 Additional results to be reported in a forthcoming full paper strongly suggest that 5b is a kinetic adduct. Studies are in progress in order to best understand the origin of the exo selectivity.
- 10 Absence of epimerization during the auxiliary removal process could be ascertained after NMR analysis of 3 (400MHz, CDCl₃) using mandelic acid (1 eq.) as chiral solvating agent.
- 11 We have experienced considerable difficulties in removing the 2-oxazolidinone chiral auxiliary from adduct **5b** probably due to steric shielding of the exocylic carbonyl group. Transesterification with lithium benzyloxide gave no more than 40% yield of the corresponding **3**. Other protocols employing lithium hydroxide, basic hydrogen peroxide or lithium borohydride proved to be ineffective in the case at hand.
- 12 3: white solid, mp 88-89°C (diethylether-petroleum ether); $[\alpha]_D = -123.1$ (c = 0.26, CHCl₃); IR (KBr): 1733, 1612 cm⁻¹; ¹H NMR (400 Mz, CDCl₃, δ): 2.41 (s, 6H), 2.76 (ddd, J = 3.9 Hz, 10.1 Hz, 12.2 Hz, 1H), 3.11 (dd, J = 3.9 Hz, 12.2 Hz, 1H), 3.54 (dd, J = 12.2 Hz, J = 12.2 Hz, 1H), 4.45 (d, J = 10 Hz, 1H), (.5.20 (d, J = 12 4 Hz, 1H), 5.25 (d, J = 12.4 Hz, 1H), 7.38 (m, 3H), 7.87 (m, 2H); ¹³C (100MHz, CDCl₃, δ): 28.3, 40.1, 40.3, 66.9, 79.2, 126.6, 128.2, 128.3, 128.4, 128.6, 130.8, 136.0, 138.5, 157.8, 173.4; MS: m/e =354 (M⁺), 192, 159, 121, 89, 77.

(Received in UK 20 February 1995)