

Evaluation of Enzyme Substrate Radiotracers as PET/MRS Hybrid Imaging Agents.

Allen F Brooks, Lindsey R Drake, Xia Shao, Austin Zhao, Peter J. H. Scott, and Michael R. Kilbourn

ACS Med. Chem. Lett., **Just Accepted Manuscript** • DOI: 10.1021/acsmchemlett.8b00402 • Publication Date (Web): 03 Oct 2018

Downloaded from <http://pubs.acs.org> on October 4, 2018

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

Evaluation of Enzyme Substrate Radiotracers as PET/MRS Hybrid Imaging Agents.

Allen F. Brooks,^{a,†} Lindsey R. Drake,^{b,†} Xia Shao,^a Austin Zhao,^a Peter J. H. Scott,^{a,b,*} Michael R. Kilbourn.^{a,*}

^a Department of Radiology, University of Michigan Medical School, 1301 Catherine Street, Ann Arbor, Michigan 48109, United States.

^b Department of Medicinal Chemistry, University of Michigan, 930 North University Avenue, Ann Arbor, Michigan 48109, United States.

KEYWORDS. Hybrid Imaging • Carbon-11 • Positron Emission Tomography • Magnetic Resonance Spectroscopy • Neuroimaging.

ABSTRACT: The development of a PET/MRS hybrid imaging agent allows for functional imaging by both methods with a single imaging agent. Enzyme substrates that are cleaved to form two metabolites present an interesting opportunity, as the unique metabolites generated might each be detected by a different modality. To be successful, such enzyme substrates would require administration of doses which (a) reach the *in vivo* target tissue at concentrations necessary for MRS imaging, (b) do not show substrate inhibition of tissue uptake or enzyme activity, and (c) provide PET images that still reflect the action of the enzyme. We report *in vitro* and *in vivo* proof-of-concept studies of a carbon-11 small molecule substrate for brain monoamine oxidases that upon enzyme-mediated cleavage produces two metabolites, one detectable by PET and the other by MRS.



Positron emission tomography (PET) imaging is a non-invasive molecular imaging method that has been utilized for decades in studies of physiology and pharmacology in living animals and human subjects.^{1,2} Most applications of PET target high affinity but low capacity sites on proteins (e.g. receptors, enzymes and transporters) and typically require high molar activities (radioactivity per mole), with clear decreases of specific binding as the mass of the injected radiotracer is increased.³ The molar activities commonly acceptable for PET imaging (>37 GBq/μmol; >1,000 Ci/mmol) are thus usually considered incompatible with the higher concentrations (mmol) of imaging agents required for alternate functional imaging methods like magnetic resonance spectroscopy (MRS).⁴

The recent development of PET/MR imaging instrumentation has made it possible to simultaneously acquire imaging data for both modalities.⁵⁻⁷ However, combining PET imaging with functional magnetic resonance spectroscopy (MRS) imaging has proven challenging. This would require a PET imaging agent that can function successfully at the high mass concentrations required for MRS imaging, despite the resulting low molar activity known to be problematic for PET radioligands. We hypothesized that the technique of metabolic trapping, whereby the product of an enzymatic reaction is trapped and retained in tissues at the site of the enzyme, might provide a suitable approach to combining PET and MRS: if the enzyme does not exhibit substrate inhibition *in vivo*, then a high mass/low molar activity PET radiotracer should still function well. To test this hypothesis, we have examined potential metabolically-trapped substrates

for monoamine oxidases (MAO-A and -B), enzymes of considerable interest in studies of brain biochemistry using PET, and report an example of a small molecule hybrid PET/MRS imaging agent.

Our prior radiotracer development efforts yielded several carbon-11 labeled N-methyl-4-aryloxy-1,2,3,6-tetrahydropyridines,⁸ based on the work of Castagnoli and coworkers,^{9,10} which are *in vitro* and *in vivo* substrates for MAO-A or MAO-B. These function *in vivo* as metabolic trapping PET agents, as the enzymatic action releases a carbon-11 labeled N-methylpyridinone that is retained in brain tissues. Inclusion of a coumarin group as the phenoxy substituent produced a radiotracer that released both radiolabeled and fluorescent metabolic products (Figure 1a), similar to a previous report of a MAO-B selective fluorogenic probe (but which was not radiolabeled).¹¹

To produce a combined PET/MRS radiotracer, we desired a 4-aryloxy substituent which would provide a metabolite with a unique signal visible by MRS following MAO oxidation. For this, we turned to the work of Yamaguchi et. al. who reported a MAO substrate described as a chemical shift-switching agent (Figure 1b), as the ¹⁹F NMR signal of the substrate (a 2-fluoro-4-nitrophenyl ether) could be distinguished from that of the metabolite (2-fluoro-4-nitrophenol).¹² We thus hypothesized that combining 2-fluoro-4-nitrophenol with the N-[¹¹C]methyl-1,2,3,6-tetrahydropyridine moiety should provide a molecule suitable for testing our hypothesis that a combined PET/MRS radiotracer is feasible using the metabolic trapping concept (Figure 1c).

a) PET/Fluorescence (Our Previous Work⁸):b) Chemical Shift-Switching (Yamaguchi¹²):

c) A PET/MR Hybrid Probe (This Work):





Figure 1. Design of a PET/MRS hybrid probe for monoamine oxidases, building on published examples of PET/fluorescence and chemical shift-switching *in vitro* MAO substrates.

The authentic standard 4-(2-fluoro-4-nitrophenoxy)-1-methyl-1,2,3,6-tetrahydropyridine (**1**) and corresponding N-desmethyl compound (**2**) were prepared using methods previously reported for syntheses of 4-aryloxy substituted tetrahydropyridines.^{8,13} To first demonstrate that the proposed radiotracer was a substrate for monoamine oxidases, kinetic studies were done with solutions of human MAO (hMAO)-A or -B and using the change in absorbance from substrate to metabolite in a spectrophotometric assay. As shown in Figure 2, **1** is a substrate for both hMAO-A and -B, with similar catalytic efficiencies (k_{cat}/K_M) for both isozymes.



	k_{cat} (min^{-1})	K_M (μM)	k_{cat}/K_M ($\text{min}^{-1}\mu\text{M}^{-1}$)
MAO-A	0.0416 ± 0.0016	19 ± 2.3	0.0022
MAO-B	0.240 ± 0.048	67 ± 4.5	0.0036

Figure 2. Michaelis-Menten kinetic analysis data for MAO-A and -B by absorbance assay.

To next demonstrate that the compound would be useful as a chemical shift-switching agent for MRS, enzymatic reactions with hMAO-A and **1** were run for 90 min and quenched by addition of an excess of the irreversible inhibitor clorgyline. ¹⁹F NMR analyses showed that over the time period the signal for substrate **1** at -130 ppm was replaced by that of the phenol metabolite **4** at -136 ppm, as well as a secondary metabolite at -129 ppm (Figure 3).




Figure 3. ¹⁹F NMR experiments (376 MHz) demonstrating time-dependent formation of 2-fluoro-4-nitrophenol metabolite (-136 ppm) & secondary metabolite (-129 ppm) via oxidation of substrate **1** (-130 ppm) by MAO-A.

With **1** confirmed *in vitro* as a MAO chemical shift-switching agent, the last step was to radiolabel with carbon-11 and evaluate whether [¹¹C]**1** could be utilized as a PET imaging agent at low molar activity. The procedure to radiolabel [¹¹C]**1** was similar to the method previously described.^{8,13} Briefly, [¹¹C]carbon dioxide was converted by standard procedure into [¹¹C]CH₃OTf and sparged through a solution of the pyridine precursor (1 mg) in ethanol (0.2mL). The [¹¹C]methyl pyridinium intermediate was transferred to a conical vial containing NaBH₄ (2 mg) in ethanol (0.3mL) and stirred for 5 min. [¹¹C]**1** was then purified by HPLC and reformulated into a 10% ethanol in saline dose (392 ± 207 MBq; non-decay corrected 1.2% radiochemical yield based on starting [¹¹C]CH₃OTf; 30 min from end of bombardment) for use in animal imaging studies. Final product was obtained in high molar activities (>259 GBq/ μmol), and to prepare doses of low molar activity [¹¹C]**1**, unlabeled standard was added to the saline used for reformulation. Final molar activities were confirmed using HPLC analysis of the UV-254nm absorbances.

MAO is known to be a highly conserved enzyme between species with, for example, high sequence homology between hMAO and rat MAO (rMAO).¹⁴ Therefore kinetic PET imaging studies were conducted in Sprague Dawley rats. Experiments conducted with high (259 GBq/ μmol) and low (0.0185 GBq/ μmol) molar activity [¹¹C]**1** gave very similar kinetic curves for whole brain radioactivity (Figure 4). These data were also used in conjunction with specific activity to generate curves estimating mass of tracer in the brain (inset, Figure 4). For the low specific activity PET studies, 1.29 ± 0.86 μmol of **1** was administered, similar to

the mass administered for prior rodent ^{19}F -MRI studies,¹⁵ and peak uptake of **1** in the rodent brain was ~ 0.02 mM. Given that the total concentration of MAO (A + B) in the rodent brain is ~ 11 pmol/mg protein,¹⁶ and that rat brain is $\sim 10\%$ protein,¹⁷ this corresponds to an estimated total rMAO concentration of ~ 1.1 pmol/mg tissue (0.0011 mM). Thus $[\mathbf{1}]/[\text{rMAO}] \sim 18$ and, given the high catalytic efficiency of MAO (Figure 2), we would not expect to overwhelm MAO at this concentration of **1** in the brain. Reflecting this, there was indeed no evidence that the MAO enzymes in the brain tissues were subject to *in vivo* substrate inhibition at this concentration.




Figure 4. Main panel: averaged ($n = 2$) rodent whole brain time-radioactivity curves (0 – 60 min post-iv-injection of high (259 GBq/ μmol , □) and low (0.0185 GBq/ μmol , ■) molar activity doses of [^{11}C]1). Inset: whole brain concentration curve (0 – 60 min post-iv-injection of low (●) molar activity doses of [^{11}C]1).

In summary, enzyme substrates that form trapped metabolites appear as a promising way forward to design PET/MRS hybrid imaging agents. In the system evaluated here, the enzyme reaction produces two metabolites that are separately visible by PET or NMR. When considering potential *in vivo* applications of such a hybrid PET-MRS imaging agent in the future, if both metabolites are trapped in the tissue they will yield a measure of the same parameter – MAO oxidation. Of more interest would be if the two methods gave different biochemical measures. How might that be achieved? Two possibilities come to mind. First, if the MRS-visible metabolite is freely diffusible across the BBB but is designed to have high affinity but reversible binding to a specific CNS site (e.g., receptor), then the retention and delayed clearance from the brain tissue might reflect the amount of specific binding (the pharmacokinetics of delivery are identical to the radiotracer trapping curve). As an alternative, if the metabolite is designed not to be BBB permeable but instead is a substrate for an efflux transporter (e.g., P-glycoprotein - Pgp), then the rate of clearance from

the tissues will reflect the transporter function, as previously described for the glutathione-conjugated metabolites of radiolabeled purines.¹⁸ Future work will address the interesting possibilities of designing bifunctional PET-MRS brain imaging tracers based on MAO, and the extension to other enzymes that can be studied using the metabolic trapping principle.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: XXX:

Experimental procedures for chemistry and radiochemistry as well as spectra / chromatograms for all novel compounds synthesized. Procedures for *in vitro* assays and *in vivo* rodent PET imaging.

AUTHOR INFORMATION

Corresponding Author

* Email: pjhscott@umich.edu

* Email: mkilbour@umich.edu

Author Contributions

The manuscript was written through contributions of all authors, and all authors have given approval to the final version of the manuscript. † These authors contributed equally.

Funding Sources

We thank NIH (R21NS075553 (MRK); T32GM007767 (PJHS and LRD)) and the University of Michigan (Energy Institute/Michigan Memorial Phoenix Project (AFB), Michigan Institute for Clinical and Health Research/TREC (LRD), College of Pharmacy (LRD and PJHS), Dept of Radiology (PJHS) and Rackham Graduate School (LRD)) for financial support.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Phillip Sherman, Janna Arteaga and Jenelle Stauff for conducting pre-clinical PET imaging.

ABBREVIATIONS

BBB, Blood-brain barrier; Bq, Becquerel; CNS, Central nervous system; MAO, Monoamine oxidase; PET, Positron emission tomography; Pgp, P-glycoprotein.

REFERENCES

- (1) For a general overview of PET imaging, see: Ametamey, S. M.; Honer, M.; Schubiger, P. A. Molecular Imaging with PET. *Chem. Rev.* **2008**, *108*, 1501-1516.
- (2) For general reviews covering PET radiochemistry, see: (a) Miller, P. W.; Long, N. J.; Vilar, R.; Gee, A. D. Synthesis of ^{11}C , ^{18}F , ^{15}O , and ^{13}N Radiolabels for Positron Emission Tomography. *Angew. Chemie Int. Ed.* **2008**, *47*, 8998-9033; (b) Brooks, A. F.; Topczewski, J. J.; Ichiishi, N.; Sanford, M. S.; Scott, P. J. H. Late-stage [^{18}F]Fluorination: New Solutions to Old Problems. *Chem. Sci.* **2014**, *5*, 4545-4553.
- (3) Noguchi, J.; Zhang, M. R.; Yanamoto, K.; Nakao, R.; Suzuki, K. *In Vitro* Binding of [^{11}C]Raclopride with Ultrahigh Specific Activity in Rat Brain Determined by Homogenate Assay and Autoradiography. *Nucl. Med. Biol.* **2008**, *35*, 19-27.

(4) Dietz, C.; Ehret, F.; Palmas, F. Vandergrift, L. A.; Jiang, Y.; Schmitt, V.; Dufner, V.; Habel, P.; Nowak, J.; Cheng, L. L. Applications of high-resolution magic angle spinning MRS in biomedical studies II-Human diseases. *NMR Biomed.* **2017**, *30*, e3784.

(5) For a recent review of PET-MRI in the context of brain imaging, see: Chen, Z.; Jamadar, S. D.; Li, S.; Sforazzini, F.; Baran, J.; Ferris, N.; Shah, N. J.; Egan, G. F. From Simultaneous to Synergistic MR-PET Brain Imaging: A Review of Hybrid MR-PET Imaging Methodologies. *Hum. Brain Mapp.* **2018**, <https://doi.org/10.1002/hbm.24314>.

(6) Judenhofer, M. S.; Wehrl, H. F.; Newport, D. F.; Catana, C.; Siegel, S. B.; Becker, M.; Thielscher, A.; Kneilling, M.; Lichy, M. P.; Eichner, M.; Klingel, K.; Reischl, G.; Widmaier, S.; Röcken, M.; Nutt, R. E.; Machulla, H. J.; Uludag, K.; Cherry, S. R.; Claussen, C. D.; Pichler, B. J. Simultaneous PET-MRI: A New Approach for Functional and Morphological Imaging. *Nat. Med.* **2008**, *14*, 459-465.

(7) Pichler, B. J.; Wehrl, H. F.; Kolb, A. Judenhofer, M. S. Positron Emission Tomography/Magnetic Resonance Imaging: the Next Generation of Multimodality Imaging? *Semin. Nucl. Med.* **2018**, *38*, 199-208.

(8) Brooks, A. F.; Shao, X.; Quesada, C. A.; Sherman, P. Scott, P. J. H.; Kilbourn, M. R. *In Vivo* Metabolic Trapping Radiotracers for Imaging Monoamine Oxidase-A and -B Enzymatic Activity. *ACS Chem. Neurosci.* **2015**, *6*, 1965-1971.

(9) Zhao, Z.; Dalvie, D.; Naiman, N.; Castagnoli, K.; Castagnoli, N. Jr. Design, Synthesis, and Biological Evaluation of Novel 4-Substituted 1-Methyl-1,2,3,6-tetrahydropyridine Analogs of MPTP. *J. Med. Chem.* **1992**, *35*, 4473-4478.

(10) Kalgutkar, A. S.; Castagnoli, K.; Hall, A.; Castagnoli, N. Jr. Novel 4-(Aryloxy)tetrahydropyridine Analogs of MPTP as Monoamine Oxidase A and B Substrates. *J. Med. Chem.* **1994**, *37*, 944-949.

(11) Long, S.; Chen, L.; Xiang, Y.; Song, M.; Zheng, Y.; Zhu, Q. An Activity-based Fluorogenic Probe for Sensitive and Selective Monoamine Oxidase-B Detection. *Chem. Commun.* **2012**, *48*, 7164-7166.

(12) Yamaguchi, K.; Ueki, R.; Nonaka, H.; Sugihara, F.; Matsuda, T.; Sando, S. Design of Chemical Shift-switching ¹⁹F Magnetic Resonance Imaging Probe for Specific Detection of Human Monoamine Oxidase A. *J. Am. Chem. Soc.* **2011**, *133*, 14208-14211.

(13) See Supporting Information for experimental details.

(14) Bai, C.; Meng, H.; Zhao, W.; Pan, Y. Molecular Cloning and Characterization of Monoamine Oxidase A and B Genes in Pig. *DNA Sequence*, **2005**, *16*, 161-165.

(15) Ko, I. O.; Jung, K-H.; Kim, M. H.; Kang, K. J.; Lee, K. C.; Kim, K. M.; Noh, I.; Lee, Y. J.; Lim, S. M.; Kim, J. Y.; Park, J-A. Preliminary ¹⁹F-MRS Study of Tumor Cell Proliferation with 3'-deoxy-3'-fluorothymidine and Its Metabolite (FLT-MP). *Contrast Media Mol. Imaging* **2017**, Article ID 3981358.

(16) Saura, J.; Kettler, R.; Da Prada, M.; Richards, J. G. Quantitative Enzyme Radioautography with ³H-Ro 41-1049 and ³H-Ro 19-6327 *in vitro*: Localization and Abundance of MAO-A and MAO-B in Rat CNS, Peripheral Organs, and Human Brain. *J. Neurosci.* **1992**, *12*, 1977-1999.

(17) Banay-Schwartz, M.; Kenessey, A.; DeGuzman, T.; Lajtha, A.; Palkovits, M. Protein Content of Various Regions of Rat Brain and Adult and Aging Human Brain. *AGE*, **1992**, *15*, 51-54.

(18) Galante, E.; Okamura, T.; Sander, K.; Kikuchi, T.; Okada, M.; Zhang, M. R.; Robson, M.; Badar, A.; Lythgoe, M.; Koeppe, M.; Årstad, E. Development of Purine-derived ¹⁸F-labeled Pro-drug Tracers for Imaging of MRP1 Activity with PET. *J. Med. Chem.* **2014**, *57*, 1023-1032.

Table of Contents artwork

