

Synthesis of Lysergic Acid Derivatives by Triple Radical Cyclisation

David E. Cladingboel^a and Philip J. Parsons^{*b}

^a Department of Chemistry, The University of Southampton, Highfield, Southampton SO9 5NH, UK

^b Department of Chemistry, The University of Reading, Whiteknights, Reading RG6 2AD, UK


The lysergic acid ring system, characteristic of the ergot alkaloids, is constructed from 2-bromoaniline derivatives by triple radical cyclisation, initiated with tri-*n*-butyltin hydride; formation of a 6- rather than a 5-membered D ring is controlled by a terminal phenylthio group.

Lysergic acid **1**¹ and its derivatives have been used over the years as psychoactive drugs, and reports as early as 1582 describe the use of ergot extracts to control uterine contractions.² The interesting biological profile of the ergot alkaloids coupled with the possibility of deriving new 5-HT bioactive molecules led us to design new ways of constructing lysergic acid and its analogues. Earlier synthetic approaches to **1** have relied on annulation reactions from an indolyl unit;³ we now report a method for the construction of lysergic acid which utilises the triple radical cyclisation reaction⁴ detailed in the analysis shown in Scheme 1.


Our synthetic approach to the framework of lysergic acid is shown in Scheme 2 and begins with the readily available amine **2** (Scheme 2a). Treatment of the substituted aniline **2** with one equivalent of lithium diisopropylamide (LDA) in tetrahydrofuran (THF) at -78°C followed by the sequential addition of allyl bromide and acetyl chloride gave, after work-up with aqueous HF in acetonitrile (2%), the alcohol **3** (83% overall). Conversion of alcohol **3** into aldehyde **4** was achieved in 95% yield using freshly prepared MnO_2 in dry THF. The aldehyde **4** reacted smoothly with methoxymethylenetriphenylphosphorane to give, after work-up with aqueous HCl in THF, the homologated aldehyde **5**[†] in 85% yield. Exposure of the aldehyde **5** to the allylic amine **6** in dry benzene containing Linde 13X molecular sieves gave the enamine **7** (90%).

Addition of Bu_3SnH (1 equiv.) in benzene containing azobutyronitrile (AIBN) (10 mol%) to a boiling solution of enamine **7** in benzene (18 h) gave the tetracyclic amine **8** (70%) as a mixture of diastereoisomers at C-8. Formation of the five-membered ring in the D ring of **8** was disappointing, but a ring expansion could be used to create the desired six-membered ring.


A direct approach to the desired six-membered D ring utilises the enamine **10** (Scheme 3) in that the phenylthio group would act as a leaving group in the radical cyclisation.




[†] Spectroscopic data: Compound **5**: ν_{max} (film) 1735 s, and 1660 s br cm^{-1} ; δ_{H} (270 MHz; CDCl_3) 1.82 (3 H, s), 3.63–3.71 (1 H, m), 4.00 (2 H, d, J 1.35 Hz), 4.78–4.85 (1 H, m), 5.03–5.13 (2 H, m), 5.82–5.97 (1 H, m), 7.03–7.44 (3 H, m) and 9.81 (1 H, t, J 1.35 Hz); δ_{C} (67.9 MHz; CDCl_3) 22.4 (q), 50.8 (t), 51.1 (t), 118.6 (t), 126.8 (s), 128.4 (d), 130.2 (d), 131.6 (d), 132.7 (d), 135.4 (s), 142.2 (s), 170.0 (s) and 197.4 (d).



Scheme 2 Reagents: i, LDA, $\text{BrCH}_2\text{CH}=\text{CH}_2$, THF; ii, AcCl , THF; iii, HF, MeCN, H_2O ; iv, MnO_2 , THF; v, $\text{Ph}_3\text{P}=\text{CH}(\text{OMe})$; vi, HCl, THF, H_2O ; vii, **6**, PhH, 13X mol. sieves, room temp.; viii, Bu_3SnH , AIBN, PhH, heat



Scheme 3



Scheme 2a Reagents: i, BH_3 , THF; ii, Fe, HOAc, EtOH; iii, $\text{Bu}^t\text{Me}_2\text{SiCl}$, DMF, imidazole

Reaction of the secondary amine **9** with **5** as before gave the enamine **10** (90%). Cyclisation of **10** mediated with Bu_3SnH gave the desired tetracyclic indoline **11** (20%) as the only isolable product.

With these results we are working on the ring expansion of the amine **8** and methods to establish the desired 9,10-double bond in lysergic acid using radical chemistry.

We thank the SERC for a research studentship to D. E. C.

Received, 20th June 1990; Com. 0/02759F

References

- 1 E. C. Kornfield, E. J. Fornefeld, G. B. Kline, M. J. Mann, D. E. Morrison, R. G. Jones and R. B. Woodward, *J. Am. Chem. Soc.*, 1956, **78**, 3087.
- 2 F. Nunes, *J. Chem. Educ.*, 1968, **45**, 688.
- 3 D. C. Horwell, *Tetrahedron*, 1980, **36**, 3123.
- 4 For other triple radical cyclisations see J. L. Lallemand, M. Julia and D. Mansuy, *Tetrahedron Lett.*, 1973, **24**, 4461; L. Stella, *Angew. Chem., Int. Ed. Engl.*, 1983, **22**, 337; A. L. Beckwith, D. H. Roberts, C. H. Schiesser and A. Wallner, *Tetrahedron Lett.*, 1985, **26**, 3349; M. A. Drombroski, S. A. Kates and B. B. Snider, *J. Am. Chem. Soc.*, 1990, **112**, 2759.