This article was downloaded by: [University of Auckland Library] On: 09 February 2015, At: 19:25 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn19

Ribosylation of Pyrimidine 2'-Deoxynucleosides

Sergey N. Mikhailov^a, Erik De Clercq^b & Piet Herdewijn^b ^a Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str. 32, Moscow, 117984, Russia

^b Regs Institute, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000, Leuven, Belgium Published online: 22 Aug 2006.

To cite this article: Sergey N. Mikhailov , Erik De Clercq & Piet Herdewijn (1996) Ribosylation of Pyrimidine 2'-Deoxynucleosides, Nucleosides and Nucleotides, 15:7-8, 1323-1334, DOI: 10.1080/07328319608002433

To link to this article: http://dx.doi.org/10.1080/07328319608002433

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

RIBOSYLATION OF PYRIMIDINE 2'-DEOXYNUCLEOSIDES

Sergey N.Mikhailov^{1*}, Erik De Clercq² and Piet Herdewijn²

¹Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Vavilov str. 32, Moscow, 117984, Russia; ²Rega Institute, Katholieke Universiteit Leuven, Minderbroedersstraat 10, B-3000 Leuven, Belgium.

ABSTRACT : The previously developed method for the preparation of 2'-O-Dribofuranosyl-nucleosides is extended to ribosylation of 2'-deoxynucleosides. The scope and limitations of this reaction are discussed.

Several antibiotics (amicetin, hikizimycin, ezomycins, thuringiensin, tunicamycins) have a disaccharide structure. Their structure, synthesis and properties were recently reviewed [1,2]. Disaccharide nucleosides are also found as modified nucleosides in t-RNA [3-5]. Nevertheless, little information is available about the biological activity of these nucleosides and even the most simple representative of this series (3'-O-ribosylated-2'-deoxy nucleosides) were hitherto unknown. Therefore we started a systematic study of the synthesis of these disaccharide nucleosides and evaluated their antiviral activity.

Disaccharide nucleosides may be prepared either by coupling of suitably blocked disaccharides with nucleic acid bases or by condensation of protected nucleosides with monosaccharides. Most of complex nucleoside antibiotics have been prepared using the first route [1]. This route is rather lengthy. The synthesis of disaccharide nucleosides may be shortened when using natural nucleosides as starting compounds. Several attempts are described to use classical methods of glycosidic bond formation with blocked nucleosides as an alcohol component. The yields of these reactions were usually very low (20-30%), due to formation of several by-products [1].

Recently a general method has been developed [6-10] for the preparation of 2'-O-ribofuranosylnucleosides which were isolated from the yeast methionine initiator tRNAs [3-5]. It consists in condensation of preactivated 1-O-acetyl-2,3,5-tri-Obenzoyl- β -D-ribofuranose with tin tetrachloride and N,3',5'-O-protected ribonucleosides under the conditions proposed by S.Hanessian and J. Banoub [11] for the preparation of alkyl β -D-ribofuranosides. To get further insight into the scope and limitations of this reaction we decided to examine glycosylation of 2'-deoxynucleosides. The starting material, N,5'-O-protected 1a-e were prepared by selective silylation [12,13] of the corresponding 2'-deoxynucleosides [14].

Condensation of 1a with 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose under the previously developed conditions [7,10] (dichloroethane, 0°C, under nitrogen) gave the desired 2a in 79% yield. The reaction was completed in 40 min. Compound 2b was prepared in the same yield (71%). The glycosylation reaction proceeded stereospecifically with β -glycosidic bond formation. NMR analysis of compounds 2a and 2b revealed that the coupling constant between H-1 and H-2 (J_{1',2'}) of the ribose residue is near zero. It should be mentioned that several byproducts with higher R_f were formed in small quantities. This was also observed during the previously reported synthesis of 2'-O-ribofuranosylnucleosides [7,10]. The amounts of these by-products depended on the purity of the starting material and on the reaction conditions. Performing the reaction under strictly anhydrous conditions reduced the formation of by-products. The silyl group was selectively deblocked to yield partially protected 3a,b which may be used for further modification. Deblocking with ammonia in methanol gave free disaccharide nucleosides 4a,b in high overall yield.

Glycosylation of 1c-e yielded an untractable mixture of products due to the known instability of glycosidic bond in purine 2'-deoxynucleosides [15]. This means that the present method [3-7] can not be applied for acid labile nucleosides without further modification.

The method has also been tested for the synthesis of β -D-ribopyranosyl derivatives. It is known that pyranoses are less reactive [16] than furanoses under similar reaction conditions. In this case the reaction was performed at 20°C for 2 hr and 5 was obtained in 38% yield. After deblocking respectively, first 6 and then 7 were obtained in a high overall yield. It should be mentioned that in this series only one disaccharide 7 was obtained in crystalline form. It is clear that in spite of much lower yields in condensation reaction this method may be used for the preparation of *O*-pyranosylnucleosides.

The glycosylation reaction was examined under different reaction conditions. When the condensation of 1a with 1.2 eq. of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose was carried out in acetonitrile at 0°C for 30 min only a minor quantity of 2a was formed. The main product isolated in 28% yield (44% based on starting

sugar) proved to be trisaccharide 8. The ¹H NMR and ¹³C NMR spectra clearly demonstrated the presence of two ribose residues (both with $J_{1',2'} = 0$ Hz). The same product 8 (38% based on the sugar) was obtained when the reaction was carried out in the presence of ZnBr₂. In this case the suspension was stirred at 20°C for 3 hr. It was shown previously that silvl blocking groups could be replaced by an acetyl in the reaction with acetic bromide in the presence of Lewis acid (SnBr₂) [17].



The compounds 4a,b and 7 were found to be inactive against a number of viruses, including herpes simplex virus (HSV-1, HSV-2), vaccinia virus, vesicular stomatitis virus, parainfluenza-3 virus, reovirus-1, Sindbis virus, Coxsackie B-4 virus, Semliki forest virus and polio-1 virus. Nor did they exhibit any toxicity to the host cells [human embryonic skin-muscle (ESM) Hela, or Vero cells].

Conclusions

The general method for the preparation of O-ribofuranosylnucleosides was found to be applicable for the synthesis of pyrimidine 2'-deoxynucleosides. In the case of purine 2'-deoxynucleosides a complex mixture was formed evidently due to the glycosidic bond instability. We have demonstrated that the method may be extended to the preparation of O-pyranosylnucleosides and may be used for simple and efficient preparation of different disaccharide nucleosides for systematic SAR studies.

EXPERIMENTAL

Melting points (uncorrected) were determined with a Buchi-Tottoli instrument. UV spectra were recorded on a Philips PU8740 UV/VIS spectrophotometer. Optical rotations were measured with a JASCO DIP-360 automatic polarimeter at 20°C. Liquid secondary ion mass spectra (LSIMS) were obtained using a Kratos Concept 1H mass spectrometer. Column chromatography was performed on silica gel (0.06-0.20 mm), TLC was carried out on Kieselgel 260 F (Merck) with detection by UV light. NMR spectra were recorded using a Gemini 200 NMR spectrometer at 20°C. Chemical shifts were measured relative to solvent signals. The signals were assigned by the double resonance techniques.

General method for the preparation of 1a-e.

To a solution of 2'-deoxynucleoside (N-benzoyl-2'-deoxynucleoside) (10 mmol) and imidazole (10 mmol) in dry pyridine (20 ml), tert-butyldiphenylsilyl chloride (11 mmol) was added and the solution was kept at 20°C for 3-5 hrs until the reaction was complete according to TLC. The solvent was evaporated to dryness, the residue was dissolved in methylene chloride (in ethyl acetate for 1e), the organic layer was washed successfully with water, 10% aqueous solution of sodium bicarbonate and again with water. The organic layer was dried with Na₂SO₄, filtered, the filtrates were evaporated to dryness and evaporated with toluene. The products were purified by column chromatography on silica gel (compounds 1a and 1e were directly recrystallized from a mixture of methylene chloride - hexane).

5'-O-tert-Butyldiphenylsilyl-2'-deoxythymidine (1a).

Yield 88%. M.p. 162-163°C. Lit. [13]: M.p. 163-164°C. ¹H NMR (CDCl₃): 9.37 brs (1H, NH), 7.70-7.34 m (11H, H-6, Ph), 6.43 dd (1H, $J_{1',2'a} = 5.6$ Hz, $J_{1',2'b} = 8.2$ Hz, H-1'), 4.57 m (1H, H-3'), 4.04 m (1H, H-4'), 3.97 dd (1H, $J_{5'a,4'} = 2.8$ Hz, $J_{5'a,5'b} = -11.5$ Hz, H-5'a), 3.85 dd (1H, $J_{5'b,4'} = 2.6$ Hz, H-5'b), 3.00 brs (1H,

OH), 2.43 m (1H, H-2'a), 2.19 m (1H, H-2'b), 1.62 d (3H, $J_{5,6} = 1.2$ Hz, Me-5), 1.08 s (9H, t-Bu). ¹³C NMR (CDCl₃): 163.88 (C-4), 150.58 (C-2), 135.53 (C-6), 135.29, 132.87, 132.32, 130.15, 130.04 and 127.97 (Ph), 111.25 (C-5), 87.11 (C-4'), 84.71 (C-1'), 72.22 (C-3'), 64.14 (C-5'), 40.98 (C-2'), 26.96 (Me₃), 19.33 (Si*C*Me₃), 12.07 (Me-5).

5'-O-tert-Butyldiphenylsilyl-N⁴-benzoyl-2'-deoxycytidine (1b).

Yield 80% (foam). ¹H NMR (CDCl₃): 8.86 brs (1H, NH), 8.34 d (1H, $J_{6,5} = 7.5$ Hz, H-6), 7.92-7.35 m (16H, Ph, Bz, H-5), 6.38 t (1H, $J_{1',2'a} = J_{1',2'b} = 6.0$ Hz, H-1'), 4.53 m (1H, H-3'), 4.13 m (1H, H-4'), 4.01 dd (1H, $J_{5'a,4'} = 2.8$ Hz, $J_{5'a,5'b} = -11.6$ Hz, H-5'a), 3.86 dd (1H, $J_{5'b,4'} = 2.9$ Hz, H-5'b), 2.79 m (1H, H-2'a), 2.21 m (1H, H-2'b), 1.09 s (9H, t-Bu). ¹³C NMR (CDCl₃): 166.66 (C-4), 162.17 (C=O), 155.03 (C-2), 144.51 (C-6), 135.57, 135.42, 133.07, 132.60, 132.35, 130.08, 129.94, 127.94 and 127.55 (Ph and Bz), 96.65 (C-5), 87.53 (C-4'), 87.14 (C-1'), 70.94 (C-3'), 63.44 (C-5'), 42.16 (C-2'), 26.91 (Me₃), 19.18 (SiCMe₃).

1-[5-O-tert-Butyldiphenylsily]-3-O-(2,3,5-tri-O-benzoy]-β-D-ribofuranosyl)-2-deoxy-β-D-ribofuranosyl]-thymine (2a).

To a cold solution (0°C) of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (630 mg, 1.25 mmol) in 1,2-dichloroethane (12 ml) under nitrogen tin tetrachloride (0.16 ml, 1.38 mmol) was added and the solution was kept at 0°C for 10 min. After addition of nucleoside 1a (480 mg, 1 mmol) the resulting solution was kept at 0° C for 40 min. Then 10% aqueous solution of sodium bicarbonate (5 ml) was added and the suspension was stirred at 0° C for 20 min. The suspension was diluted with methylene chloride (20 ml), filtered through Hyflo Super Cel, the organic layer was separated, washed with water (10 ml), dried and evaporated to dryness. The residue was purified by column chromatography on silica gel (50 g). The column was washed with methylene chloride (500 ml) and then eluted with 1% methanol in methylene chloride to give 2a as a foam. Yield 730 mg (79%). LSIMS: (M + H) 925. ¹H NMR (CDCl₃): 8.68 brs (1H, NH), 8.10-7.30 m (26H, Ph, Bz, H-6), 6.29 dd $(1H, J_{1',2'a} = 5.6 \text{ Hz}, J_{1',2'b} = 8.1 \text{ Hz}, \text{H-1' Thd}), 5.84 \text{ dd} (1H, J_{3',2'} = 5.0 \text{ Hz}, J_{3',4'})$ = 6.5 Hz, H-3' Rib), 5.66 d (1H, H-2' Rib), 5.32 s (1H, H-1' Rib), 4.82-4.56 m (4H, H-3',4',5'a,5'b Thd), 4.12 m (1H, H-4' Rib), 3.95 dd (1H, $J_{5'a,4'} = 3.1$ Hz, $J_{5'a,5'b} =$ - 11.5 Hz, H-5'a Rib), 3.83 dd (1H, $J_{5'b,4'} = 2.7$ Hz, H-5'b Rib), 2.52 m (1H, H-2'a Thd), 2.18 m (1H, H-2'b Thd), 1.61 d (3H, $J_{5,6} = 1.0$ Hz, Me-5), 1.06 s (9H, t-Bu). ¹³C NMR (CDCl₃): 166.09, 165.29 and 165.16 (C=O), 163.61 (C-4), 150.04 (C-2), 135.51 (C-6), 135.29, 133.57, 133.45, 133.25, 132.82, 132.28, 130.04, 129.74, 129.05,

128.84, 128.40 and 127.96 (Ph and Bz), 111.05 (C-5), 104.69 (C-1' Rib), 84.55 (C-1' Thd), 84.24 (C-4' Thd), 79.40 (C-3' Thd), 77.66 (C-4' Rib), 75.74 (C-2' Rib), 72.21 (C-3' Rib), 64.73 (C-5' Thd), 63.88 (C-5' Rib), 38.57 (C-2' Thd), 26.93 (Me₃), 19.30 (Si*C*Me₃), 12.07 (Me-5).

N⁴-Benzoyl-1-[5-*O*-tert-butyldiphenylsilyl-3-*O*-(2,3,5-tri-*O*-benzoyl-β-*D*-ribofuranosyl)-2-deoxy-β-*D*-ribofuranosyl]-cytosine (2b).

Analogous condensation of 1-O-acetyl-2,3,5-tri-O-benzoyl-β-D-ribofuranose (1.32 g, 2.61 mmole) with 1b (1.18 g, 2.07 mmol) in the presence of tin tetrachloride (0.33 ml, 2.83 mmol) in 1,2-dichloroethane (20 ml) at 0°C for 40 min gave 2b as a foam. Yield 1.5 g (71%). LSIMS: (M + H) 10164. ¹H NMR (CDCl₃): 8.72 brs (1H, NH), 8.26 d (1H, $J_{6,5} = 7.5$ Hz, H-6), 8.10-7.25 m (31H, Ph, Bz, H-5), 6.25 t (1H, $J_{1',2'a}$ $= J_{1',2'b} = 5.6$ Hz, H-1' dCyd), 5.81 dd (1H, $J_{3',2'} = 4.7$ Hz, $J_{3',4'} = 6.7$ Hz, H-3' Rib), 5.63 d (1H, H-2' Rib), 5.28 s (1H, H-1' Rib), 4.81-4.53 m (4H, H-3' dCyd, H-4',5'a,5'b Rib), 4.15 m (1H, H-4' dCyd), 4.02 dd (1H, $J_{5'a} 4' = 3.1$ Hz, $J_{5'a} 5'b = -$ 11.4 Hz, H-5'a dCyd), 3.85 dd (1H, $J_{5'b,4'} = 2.5$ Hz, H-5'b dCyd), 2.79 ddd (1H, $J_{2'a,3'} = 5.1 \text{ Hz}, J_{2'a,2'b} = -14.0 \text{ Hz}, \text{ H-2'a dCyd}, 2.30 \text{ ddd} (1\text{ H}, J_{2'b,3'} = 6.0 \text{ Hz}, \text{ H-}$ 2'b dCyd), 1.08 s (9H, t-Bu). ¹³C NMR (CDCl₃): 165.98 (C-4), 165.18 and 165.04 (C=O), 161.93 (C-2), 144.19 (C-6), 135.51, 135.36, 133.50, 133.35, 133.16, 133.03, 132.45, 132.14, 130.06, 129.67, 128.92, 128.76, 128.36, 128.29, 127.91 and 127.47 (Ph and Bz), 104.81 (C-1' Rib), 96.33 (C-5), 86.69 (C-1' dCyd), 84.69 (C-4' dCyd), 79.23 (C-3' dCyd), 76.01 (C-4' Rib), 75.59 (C-2' Rib), 72.12 (C-3' Rib), 64.57 (C-5' dCyd), 62.82 (C-5' Rib), 39.88 (C-2' dCyd), 26.85 (Me₃), 19.12 (SiCMe₃).

$1-[3-O-(2,3,5-tri-O-Benzoyl-\beta-D-ribofuranosyl)-2-deoxy-\beta-D-ribofuranosyl]-thymine (3a).$

Nucleoside 2a (670 mg, 0.72 mmol) was dissolved in 0.5 M tetrabutylammonium fluoride trihydrate in tetrahydrofuran (2 ml). The solution was kept for 45 min at 20°C, evaporated to dryness, coevaporated with chloroform (10 ml) and applied onto a silica gel column (20 g). The column was washed with methylene chloride (300 ml) and with 1% methanol in methylene chloride (200 ml) and then eluted with 2% methanol in methylene chloride to give 3a as a foam. Yield 0.43 g (86%). LSIMS: (M + H) 687. ¹H NMR (CDCl₃): 9.01 brs (1H, NH), 8.10-7.25 m (16H, Bz, H-6), 5.95 t (1H, J₁', 2'a = J₁', 2'b = 6.8 Hz, H-1' Thd), 5.84 dd (1H, J₃', 2' = 4.9 Hz, J₃', 4' = 6.4 Hz, H-3' Rib), 5.66 d (1H, H-2' Rib), 5.40 s (1H, H-1' Rib), 4.80-4.52 m (4H, H-3', 5'a, 5'b Thd, H-4' Rib), 4.10 m (1H, H-4' Thd), 3,91 dd (1H, J₅'a, 4' = 3.0 Hz, J₅'a, 5'b = -12.3 Hz, H-5'a Rib), 3.77 dd (1H, J₅'b, 4' = 1.5 Hz, H-5'b

Rib), 3.21 brs (1H, HO-5'), 2.54-2.40 m (2H, H-2'a,2'b Thd), 1.88 d (3H, $J_{5,6} = 1.2$ Hz, Me-5). ¹³C NMR (CDCl₃): 166.15, 165.39 and 165.29 (C=O), 163.72 (C-4), 150.23 (C-2), 137.63 (C-6), 133.62, 133.51, 133.36, 129.71, 129.49, 128.93, 128.75, 128.50 and 128.40 (Ph), 110.89 (C-5), 105.31 (C-1' Rib), 88.01 (C-1' Thd), 84.82 (C-4' Thd), 79.35 (C-3' Thd), 78.24 (C-4' Rib), 75.72 (C-2' Rib), 72.03 (C-3' Rib), 64.63 (C-5' Thd), 62.35 (C-5' Rib), 37.60 (C-2' Thd), 12.41 (Me-5).

N^4 -Benzoyl-1-[3-O-(2,3,5-tri-O-benzoyl- β -D-ribofuranosyl)-2-deoxy- β -D-ribofuranosyl]-cytosine (3b).

Analogous desilylation of 2b yielded 3b as a foam (89%). LSIMS: (M + H) 776. ¹H NMR (CDCl₃): 8.82 brs (1H, NH), 8.14-7.27 m (22H, Bz, H-5,6), 6.02 t (1H, J₁',2'a = J₁',2'b = 6.2 Hz, H-1' dCyd), 5.83 dd (1H, J₃',2' = 4.8 Hz, J₃',4' = 6.5 Hz, H-3' Rib), 5.67 d (1H, H-2' Rib), 5.41 s (1H, H-1' Rib), 4.78-4.52 m (4H, H-3' dCyd, H-4',5'a,5'b Rib), 4.19 m (1H, H-4' dCyd), 4.00 dd (1H, J₅'a,4' = 2.0 Hz, J₅'a,5'b = -11.5 Hz, H-5'a dCyd), 3.85 dd (1H, J₅'b,4' = 2.5 Hz, H-5'b dCyd), 2.72-2.51 m (2H, H-2'a,2'b dCyd). ¹³C NMR (CDCl₃): 166.08 (C-4), 165.28 and 162.23 (C=O), 154.83 (C-2), 146.16 (C-6), 133.52, 133.40, 133.25, 133.06, 129.68, 128.90, 128.44, 128.33 and 127.55 (Bz), 105.27 (C-1' Rib), 96.61 (C-5), 89.54 (C-1' dCyd), 85.49 (C-4' dCyd), 79.20 (C-3' dCyd), 77.64 (C-4' Rib), 75.69 (C-2' Rib), 72.11 (C-3' Rib), 64.61 (C-5' dCyd), 61.83 (C-5' Rib), 38.75 (C-2' dCyd).

1-(3-O-β-D-Ribofuranosyl-2-deoxy-β-D-ribofuranosyl)-thymine (4a).

A solution of nucleoside 3a (350 mg, 0.51 mmol) in 5 M ammonia in methanol (10 ml) was kept for 5 days at 20° C and then concentrated in vacuo to dryness. The residue was partitioned between methylene chloride (10 ml) and water (20 ml), and the aqueous layer was washed with methylene chloride (2 x 10 ml). The aqueous layer was evaporated to dryness, the residue was coevaporated with methanol (2 x 5 ml) to give a hygroscopic foam. Yield 190 mg (99%). $[\alpha]_{D}$ -28° (c 1.0, water). UV (pH 1-7): λ max 267 nm (ϵ 9600); (pH 13): λ max 267 nm (ϵ 7300). LSIMS: $(C_{15}H_{22}N_{2}O_{9} + H)$. Calc. 375.1403. Found 375.1419. ¹H NMR (D₂O): 7.50 d (1H, $J_{6,5} = 1.2$ Hz, H-6), 6.11 t (1H, $J_{1',2'a} = J_{1',2'b} = 6.8$ Hz, H-1' Thd), 4.99 s (1H, H-1' Rib), 4.33 m (1H, H-3' Thd), 4.10 dd (1H, $J_{3',2'} = 4.8$ Hz, $J_{3',4'} = 6.3$ Hz, H-3' Rib), 4.01-3.87 m (3H, H-4',5'a,5'b Thd), 3.76-3.65 m (3H, H-2',4',5'a Rib), 3.57 dd (1H, $J_{5'b,4'} = 6.6$ Hz, $J_{5'b,5'a} = -12.4$ Hz, H-5'b Rib), 2.50-2.12 m (2H, H-2'a,2'b Thd), 1.76 d (3H, Me-5). ¹³C NMR (D₂O): 167.48 (C-4), 152.69 (C-2). 138.41 (C-6), 112.44 (C-5), 107.64 (C-1' Rib), 86.22 (C-1' Thd), 85.73 (C-4' Thd), 83.96 (C-3' Thd), 78.55 (C-4' Rib), 75.63 (C-2' Rib), 71.82 (C-3' Rib), 63.88 (C-5' Thd), 62.33 (C-5' Rib), 38.33 (C-2' Thd), 12.60 (Me-5).

1-(3-O-β-D-Ribofuranosyl-2-deoxy-β-D-ribofuranosyl)-cytosine (4b).

Analogous debenzoylation of 3b after evaporation with methanol yielded 4b as a hygroscopic foam (95%). $[\alpha]_{D}$ -2.2° (c 0.55, water). UV (pH 1): λ max 279 nm (ϵ 12100); (pH 7-13): λ max 270 nm (ϵ 8600). LSIMS: (C₁₄H₂₁N₃O₈ + H). Calc. 360.1407. Found 360.1431. ¹H NMR (D₂O): 7.68 d (1H, J₆, 5 = 7.7 Hz, H-6), 6.10 t (1H, J₁', 2'a = J₁', 2'b = 6.6 Hz, H-1' dCyd), 5.91 d (1H, H-5), 5.00 s (1H, H-1' Rib), 4.30 m (1H, H-3' dCyd), 4.11 dd (1H, J₃', 2' = 4.7 Hz, J₃', 4' = 6.7 Hz, H-3' Rib), 4.05-3.88 m (3H, H-4', 5'a, 5'b dCyd), 3.77-3.65 m (3H, H-2', 4', 5'a Rib), 3.55 dd (1H, J_{5'b}, 4' = 6.6 Hz, J_{5'b}, 5'a = -12.3 Hz, H-5'b Rib), 2.60-2.08 m (2H, H-2'a, 2'b dCyd). ¹³C NMR (D₂O): 167.12 (C-4), 158.40 (C-2), 142.45 (C-6), 107.59 (C-1' Rib), 97.16 (C-5), 87.29 (C-1' dCyd), 85.85 (C-4' dCyd), 83.97 (C-3' dCyd), 78.62 (C-4' Rib), 75.64 (C-2' Rib), 71.85 (C-3' Rib), 63.93 (C-5' dCyd). 62.40 (C-5' Rib), 39.11 (C-2' dCyd).

$1-[5-O-tert-Butyldiphenylsilyl-3-O-(2,3,4-tri-O-benzoyl-\beta-D-ribopyranosyl)-2-deoxy-\beta-D-ribofuranosyl]-thymine (5).$

To a solution of 1,2,3,4-tetra-O-O-acetyl-β-D-ribopyranose (716 mg, 2.25 mmol) in 1,2-dichloroethane (20 ml) under nitrogen, tin tetrachloride (0.29 ml, 2.5 mmol) was added and the solution was kept at 20°C for 10 min. Nucleoside 1a (680 mg, 1,415 mmol) was added in three portions over 40 min under stirring and the resulting yellow solution was kept further for 1.3 hr at 20°C. Then 10% aqueous solution of sodium bicarbonate (5 ml) was added and the suspension was stirred at 20° C for 20 min. The suspension was diluted with methylene chloride (20 ml), filtered through Hyflo Super Cel, the organic layer was separated, washed with water (10 ml), dried and evaporated to dryness. The residue was purified by column chromatography on silica gel (50 g). The column was washed with methylene chloride (500 ml) and eluted with 1% methanol in methylene chloride to give 5 as a foam. Yield 400 mg (38%). LSIMS: (M + H) 739. ¹H NMR (CDCl₃): 8.95 brs (1H, NH), 7.70-7.38 m (11H, Ph, H-6), 6.30 dd (1H, $J_{1',2'a} = 5.6$ Hz, $J_{1',2'b} = 8.2$ Hz, H-1' Thd), 5.41 dd (1H, $J_{3',2'} = 3.2$ Hz, $J_{3',4'} = 3.6$ Hz, H-3' Rib), 5.16 m (1H, H-4' Rib), 4.98 dd (1H, $J_{2',1'}$ = 4.0 Hz, H-2' Rib), 4.85 d (1H, H-1' Rib), 4.48 m (1H, H-3' Thd), 4.09 m (1H, H-4' Thd), 4.05-3.76 m (4H, H-5'a,5'b Thd, H-5'a,5'b Rib), 2.54 ddd (1H, $J_{2'a,2'b} = -13.8$ Hz, $J_{2'a,3'} = 2.1$ Hz, H-2'a Thd), 2.18 m (1H, H-2'b Thd), 2.11 s (3H, Ac), 2.10 s (3H, Ac), 2.06 s (3H, Ac), 1.63 d (3H, $J_{5.6} = 1.0$ Hz, Me-5), 1.10 s (9H, t-Bu). ¹³C NMR (CDCl₃): 169.91 and 169.62 (C=O), 163.68 (C-4), 150.16 (C-2), 135.04 (C-6), 135.46, 135.28, 132.70, 132.29, 130.11 and 127.98 (Ph), 111.15 (C-5), 97.88 (C-1' Rib), 84.73 (C-1' Thd), 84.52 (C-4' Thd), 78.18 (C-3' Thd), 68.27 (C-2' Rib), 66.48 (C-3' Rib), 65.98 (C-4' Rib), 63.93 (C-5' Thd), 61.63 (C-5' Rib), 38.65 (C-2' Thd), 26.96 (Me₃), 20.76 and 20.63 (Me, Ac), 19.31 (Si*C*Me₃), 12.06 (Me-5).

1-[3-O-(2,3,4-tri-O-Acetyl-β-D-ribopyranosyl)-2-deoxy-β-D-ribofuranosyl]thymine(6).

Nucleoside 5 (750 mg, 1.015 mmol) was dissolved in 0.5 M tetrabutylammonium fluoride trihydrate in tetrahydrofuran (3 ml), the solution was kept for 20 min at 20°C, evaporated to dryness, coevaporated with chloroform (10 ml) and applied onto a silica gel column (20 g). The column was washed with methylene chloride (300 ml) and with 1% methanol in methylene chloride (200 ml) and then eluted with 2% methanol in methylene chloride to give 6 as a foam. Yield 0.45 g (89%). LSIMS: (M + H) 501, (M + Na) 523. ¹H NMR (CDCl₃): 9.03 brs (1H, NH), 7.35 q (1H, $J_{6,5} = 1.0$ Hz, H-6), 6.05 t (1H, $J_{1',2'a} = J_{1',2'b} = 6.9$ Hz, H-1' Thd), 5.41 dd (1H, $J_{3',2'} = 3.5 \text{ Hz}, J_{3',4'} = 3.1 \text{ Hz}, \text{ H-3' Rib}, 5.13 \text{ m}$ (1H, H-4' Rib), 4.98 dd (1H, $J_{2',1'}$ = 4.1 Hz, H-2' Rib), 4.91 d (1H, H-1' Rib), 4.53 m (1H, H-3' Thd), 4.08 m (1H, H-4' Thd), 4.02-3.71 m (4H, H-5'a,5'b Thd, H-5'a,5'b Rib), 3.02 brs (1H, OH), 2.47 m (2H, H-2'a, 2'b Thd), 2.12 s (3H, Ac), 2.10 s (3H, Ac), 2.08 s (3H, Ac), 1.90 d (3H, Me-5). ¹³C NMR (CDCl₃): 169.96 and 169.77 (C=O), 163.72 (C-4), 150.28 (C-2), 137.28 (C-6), 111.05 (C-5), 98.14 (C-1' Rib), 87.69 (C-1' Thd), 84.78 (C-4' Thd), 77.98 (C-3' Thd), 68.34 (C-2' Rib), 66.44 (C-3' Rib), 66.09 (C-4' Rib), 62.31 (C-5' Thd), 61.61 (C-5' Rib), 37.96 (C-2' Thd), 20.79 and 20.67 (Me, Ac), 12.44 (Me-5).

$1-(3-O-\beta-D-Ribopyranosyl-2-deoxy-\beta-D-ribofuranosyl)-thymine (7).$

A solution of nucleoside 6 (300 mg, 0.60 mmol) in 5 M ammonia in methanol (10 ml) was kept for 2 days at 20°C and then concentrated in vacuo to dryness. The residue was partitioned between methylene chloride (10 ml) and water (20 ml), and the aqueous layer was washed with methylene chloride (2 x 10 ml). The aqueous layer was concentrated to dryness and acetone was added. After 2 days at 0°C hygroscopic crystals were filtered off, washed with acetone and dried. Yield 180 mg (80%). M.p. 126-128°C. $[\alpha]_D$ -40.7° (c 0.73, water). UV (pH 1-7): λ max 267 nm (ϵ 9400); (pH 13): λ max 267 nm (ϵ 7100). LSIMS: (C₁₅H₂₂N₂O₉ + H). Calc. 375.1403. Found 375.1380.

Anal. Calc. for C₁₅H₂₂N₂O₉.0.25H₂O: C 47.56; H 5.99; N 7.39. Found: C 47.60; H 6.01; N 7.31.

¹H NMR (D₂O): 7.55 q (1H, J_{6,5} = 1.0 Hz, H-6), 6.18 dd (1H, J_{1',2'a} = 6.2 Hz, J_{1',2'b} = 7.3 Hz, H-1' Thd), 4.77 d (1H, J_{1',2'} = 5.5 Hz, H-1' Rib), 4.41 ddd (1H,

 $J_{3',2a'} = 3.6$ Hz, $J_{3',2'b'} = 6.8$ Hz, $J_{3',4'} = 2.5$ Hz, H-3' Thd), 4.09 m (1H, H-4' Thd), 4.00 t (1H, $J_{3',2'} = J_{3',4'} = 2.9$ Hz, H-3', Rib), 3.85-3.53 m (6H, H-5'a,5'b Thd, H-2',4',5'a,5'b Rib), 2.44 ddd (1H, $J_{2'a,2'b} = -14.1$ Hz, H-2'a Thd), 2.32 ddd (1H, H-2'b Thd), 1.80 d (3H, Me-5). ¹³C NMR (D₂O): 166.90 (C-4), 152.10 (C-2), 137.84 (C-6), 111.91 (C-5), 100.55 (C-1' Rib), 85.70 (C-1' Thd), 85.22 (C-4' Thd), 78.69 (C-3' Thd), 70.61 (C-2' Rib), 68.40 (C-3' Rib), 68.03 (C-4' Rib), 63.87 (C-5' Thd), 61.76 (C-5' Rib), 37.59 (C-2' Thd), 12.00 (Me-5).

1-[Bis-3,5-O-(2,3,5-tri-O-Benzoyl- β -D-ribofuranosyl)-2-deoxy- β -D-ribofuranosyl]-thymine (8).

A. To a cold solution (0°C) of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (1.21 g, 2.4 mmol, 1.27 eq.) in dry acetonitile (20 ml) under nitrogen tin tetrachloride (0.304 ml, 2.6 mmol, 1.38 eq.) was added and the solution was kept at 0°C for 10 min. After addition of nucleoside 1a (905 mg, 1.883 mmol, 1 eq.) the resulting solution was kept at 0°C for 30 min. After usual work-up the product 8 was purified by column chromatography on silica gel (50 g). The column was washed with methylene chloride (500 ml) and then eluted with 0.8% methanol in methylene chloride. Yield 600 mg (foam) (28% based on starting 1a; 44% based on sugar).

B. A mixture of 1-O-acetyl-2,3,5-tri-O-benzoyl- β -D-ribofuranose (353 mg, 0.7 mmol), nucleoside 1a (226 mg, 0.47 mmol) and zinc bromide (450 mg, 2 mmol) in dry 1,2-dichloroethane (10 ml) was stirred under nitrogen for 3 hr at 20°C. After usual work-up the product 8 was obtained (yield 28% on nucleoside, 38% on sugar).

LSIMS: (M + H) 1131. ¹H NMR (CDCl₃): 8.38 brs (1H, NH), 8.07-7.26 m (31H, Bz, H-6), 6.16 t (1H, $J_{1',2'a} = J_{1',2'b} = 6.6$ Hz, H-1' Thd), 5.83 dd (1H, $J_{3',2'} = 5.1$ Hz, $J_{3',4'} = 6.3$ Hz, H-3' Rib), 5.82 dd (1H, $J_{3',2'} = 5.4$ Hz, $J_{3',4'} = 6.6$ Hz, H-3' Rib), 5.70 d (1H, $J_{2',3'} = 5.1$ Hz, H-2' Rib), 5.65 d (1H, $J_{2',3'} = 5.4$ Hz, H-2' Rib), 5.40 s (1H, H-1' Rib), 5.34 s (1H, H-1' Rib), 4.80-3.68 m (10H6, H-3',4',5'a,5'b Thd, 2xH-4',5'a,5'b Rib), 2.42 m (1H, H-2'a Thd), 2.23 m (1H, H-2'b Thd), 1.90 d (3H, $J_{5,6} = 1.0$ Hz, Me-5). ¹³C NMR (CDCl₃): 166.02 and 165.51 (C=O), 163.12 (C-4), 150.11 (C-2), 135.02 (C-6), 133.39, 133.20, 129.68, 128.94, 128.75, 128.33 and 127.84 (Bz), 111.14 (C-5), 106.08 (C-1' Rib), 104.96 (C-1' Rib), 84.95 (C-1' Thd), 82.39 (C-4 Thd), 79.32 (2xC-4' Rib), 77.44 (C-3' Thd), 75.72 (C-2' Rib), 75.23 (C-2' Rib), 72.00 (2xC-3' Rib), 67.29 (C-5' Thd), 64.58 (C-5' Rib), 64.41 (C-5' Rib), 37.92 (C-2' Thd), 12.48 (Me-5).

1-(Bis-3,5-O- β -D-ribofuranosyl-2-deoxy- β -D-ribofuranosyl)-thymine (9).

Analogous debenzoylation of 8 with 5 M ammonia in methanol for 4 days yielded after evaporation with methanol 9 as a hygroscopic foam (93%). $[\alpha]_D$ -46.4° (c 0.91, water). UV (pH 1-7): λ max 267 nm (ϵ 9300); (pH 13): λ max 267 nm (ϵ 6900). LSIMS: (C₂₀H₃₀N₂O₁₃ + H). Calc. 507.1826. Found 507.1812. ¹H NMR (D₂O): 7.41 d (1H, J_{6,5} = 1.0 Hz, H-6), 6.12 t (1H, J_{1',2'a} = J_{1',2'b} = 6.6 Hz, H-1' Thd), 5.00 s (1H, H-1' Rib), 4.95 s (1H, H-1' Rib), 4.38 m (1H, H-3' Thd), 4.11-3.43 m (13H, H-4',5'a,5'b Thd, 2xH-2',3',4',5'a,5'b Rib), 2.52-2.21 m (2H, H-2'a,2'b Thd), 1.80 d (3H, Me-5). ¹³C NMR (D₂O): 166.85 (C-4), 152.08 (C-2), 137.74 (C-6), 111.80 (C-5), 107.72 (C-1' Rib), 107.07 (C-1' Rib), 85.73 (C-1' and C-4' Thd), 83.36 (2xC-4' Rib), 78.04 (C-3' Thd), 74.99 (C-2' Rib), 74.74 (C-2' Rib), 71.31 (C-3' Rib), 71.20 (C-3' Rib), 67.82 (C-5' Thd), 63.22 (2xC-5' Rib), 37.48 (C-2' Thd), 12.07 (Me-5).

Financial support from INTAS (project 93-1500) and NATO (grant HTECH.CRG 940362) is gratefully acknowledged.

REFERENCES

- Lerner, L.M. Synthesis and Properties of Various Disaccharide Nucleosides in: *Chemistry of Nucleosides and Nucleotides* (Townsend L.B. ed.), Plenum Press, New York, 1991, vol. 2, 27-79.
- 2. Isono, K. J. Antibiotics, 1988, 41, 1711.
- 3. Keith, G., Glasser, Desgres, J., Keith, G., Kuo, K.C., Gehrke, C.W. Nucl. Acids Res., 1990, 18, 5989.
- Glasser, A.-L., Desgres, J., Heitzler, J., Gehrke, C.W., Keith, G. Nucl. Acids Res. 1991, 19, 5199.
- 5. Limbach, P.A., Crain, P.F., McCloskey, J.A. Nucl. Acids Res. 1994, 22, 2183.
- 6. Niewczyk, A.; Krzyzaniak, A.; Barciszewski, J.; Markiewicz, W.T. Nucleosides and Nucleotides, 1991, 10, 635.
- Mikhailov, S.N., De Bruyn, A., Herdewijn, P. Nucleosides and Nucleotides, 1995, 14, 481.
- 8. Markiewicz, W.T. Abstracts of Nucleic Acids Symposium, August 6-11, Leiden the Netherlands, L12.
- Mikhailov, S.N., Gurskaya, G.V., Efimtseva, E.V., Ermolinsky, B.S., V.E., De Bruyn, A., Rozenski, J., Herdewijn, P. Abstracts of Nucleic Acids Symposium. August 6-11. Leiden the Netherlands, O15.

- Mikhailov, S.N., Efimtseva, E.V., Gurskaya, G.V., Zavodnik, V.E., De Bruyn, A., Janssen, G., Rozenski, J., Herdewijn, P. J. Carbohydr. Chem., 1995, in press.
- Hanessian, S.; Banoub, J. in: *Methods in Carbohydrate Chemistry*, Academic Press, New York, 1980, 243.
- 12. Hanessian, S., Lavallee, P. Can. J. Chem. 1977, 53, 2975.
- 13. Larsen, E., Jorgensen, P.T., Sofan, M.A., Pedersen E.B. Synthesis 1994, 1037.
- 14. Ti, G.S., Gaffney, B.L., Jones, R.A. J. Am. Chem. Soc. 1982, 104, 1316.
- 15. Sonveaux, E. in *Protocols for Oligonucleotide Conjugates (Ed. Agrawal, S.)* Humana Press. **1994**, *26*, 1.
- 16. Vorbruggen, H., Bennua, B. Chem. Ber. 1981, 114, 1279.
- 17. Oriyama, T., Oda, M., Gono, J., Koga, G. Tetrahedron Lett. 1994, 35, 2027.

Received November 7, 1995 Accepted April 10, 1996