

BUILDING BLOCKS FOR OLIGONUCLEOTIDE ANALOGS WITH DIMETHYLENE-SULFIDE, -SULFOXIDE, AND -SULFONE GROUPS REPLACING PHOSPHODIESTER LINKAGES

K. Christian Schneider and Steven A. Benner*


Laboratory for Organic Chemistry
E.T.H. Zurich, CH-8092 Switzerland

Summary


Routes are presented for the synthesis of 3',5'-bishomo-deoxyribonucleosides, building blocks needed to synthesize oligonucleotide analogs where the -O-PO₂-O- groups are replaced by -CH₂-SO₂-CH₂-, -CH₂-S-CH₂-, and -CH₂-SO-CH₂- units. As isosteric, diastereomerically pure (in the first two cases), non-ionic analogs of natural oligonucleotides, such molecules have potential application as "anti-sense" oligonucleotide analogs.

Non-ionic oligonucleotide analogs are important synthetic targets, as they are likely to be stable to biological degradation, cross biological barriers, bind to natural oligonucleotides with a complementary sequence, and disrupt their biological function.¹ It is well-recognized that if such compounds could be found and applied *in vivo*, they could be used to treat virtually every disease involving the unwanted expression of genetic information.² As all viral diseases, many cancers, most bacterial and parasitic maladies, and many other diseases are of this type, the excitement has been understandable.


Some time ago we reported that sulfones (such as dimethylsulfoxide and sulfolane) as cosolvents can assist the penetration of natural oligonucleotides into cells.³ This observation suggested to us that incorporating the sulfone unit directly into an oligonucleotide by replacing the phosphodiester (-O-PO₂-O-) groups by sulfone (-CH₂-SO₂-CH₂-) groups might yield an especially attractive oligonucleotide analog capable of penetrating cell membranes without co-solvents. Further, sulfones are non-ionic, achiral, isosteric analogs of phosphate diesters, and are stable to both chemical and biochemical degradation, making them ideal analogs for phosphate esters on other grounds as well.



We report here a route for the synthesis of deoxyribonucleoside analogs 1-4, analogs bearing functionalization appropriate for their use as building blocks in the synthesis of oligonucleotide analogs having the phosphodiester groups replaced by dimethylene-sulfide, -sulfoxide, or -sulfone units (5). The building blocks are prepared in their protected forms (1b-4b), as these derivatives are more suitable for long term storage, and are readily converted to their unprotected forms immediately prior to coupling by standard procedures.⁴



The synthesis started from 6,⁵ which was deacetylated with NaOEt in EtOH, and then converted to cis-benzoate 7 by treatment with PPh₃, diisopropylazodicarboxylate (DIAD) and benzoic acid in THF (72 % overall). Benzoate 7 was resolved into its antipodes by enantioselective hydrolysis with pig liver esterase⁶ in H₂O/*t*-BuOH 9:1 at pH 7 (maintained by automatic titration with dilute NaOH) to give, after addition of 0.45 mol eq. of hydroxide and extractive workup, acid (-)8 ([α]_D = -89.5°, c=6.35, acetone) with >99 % ee (determined, after conversion to the corresponding methylester with diazomethane, by NMR in the presence of Eu(hfc)₃). The absolute configuration of (-)8 was assigned by direct correlation to glucose.⁷ (-)8 was reduced with LiAlH₄ in THF to diol 9 (mp: 65-66°C), which was selectively protected with pivaloyl chloride (PvCl) in pyridine at -15°C to furnish 10 in an 85% overall yield. 10 was converted to 12 in 63 % yield by the following one pot procedure: (1) ozonization in methanol at -78°C, (2) reductive work-up with dimethylsulfide, (3) selective acetalization to the cyclic acetal 11 by stirring the reaction mixture in methanol for 14 days, and (4) reduction of 11 with excess NaBH₄. Presumably, traces of acid which formed during ozonolysis catalyse the selective acetalization in step 3. The 6-hydroxyl group of 12 was internally protected by acid catalysed cyclization (Dowex[®] H⁺) in refluxing toluene (85%) to give 13. Since the pivaloate group in 13 is too stable to be removed in the presence of amide protected bases, 13 was converted to benzoate 14 by hydrolysis with 1 M NaOH (3 h, 25°C), followed by benzylation (BzCl, pyridine, 83% overall).



The appropriately protected bases were (after bis-silylation with trimethylsilyl chloride/hexamethyldisilazane (TMSCl/HMDS) or, preferentially, *in situ* using *N*-methyl-*N*-TMS-trifluoroacetamide, MSTFA) introduced into 14 via Lewis-acid catalysis using TMSOTf in acetonitrile.⁸ The thioacetate moiety was introduced in all cases by a Mitsunobu reaction involving the slow addition of a mixture

of the nucleoside analog and AcSH (2 eq.) to a pre-formed complex of PPh₃ (2 eq.) and DIAD in THF at 0°C.⁹ Thus, **14** was reacted with bis-TMS-benzoyladenine in the presence of TMS-triflate (TMSOTf, 0.5 eq.) at 40°C for 30 min. to furnish **15** as a 1:1 mixture of anomers in 64% yield. The anomers were separated by HPLC (silica gel, CH₂Cl₂/THF 65:35) and the β-anomer was then converted to thioacetate **1b** in 69% yield. Reaction of **14** with bisTMS-*i*-butyrylguanidine in the presence of TMSOTf (0.5 eq.) at 40°C for 1h gave the guanosine analog **16** as a 2:1 mixture of N⁹- and N⁷-isomers, each as a 1:1 mixture of anomers. The regioisomers were separated by silica gel chromatography (CH₂Cl₂/MeOH 9:1) to furnish the N⁹-isomer **16** in 56% yield and the N⁷-isomer **22** (30%). After conversion of the N⁹-isomer (**16**) to thioacetate **17** (73%) the anomers were separated by HPLC (silica gel, CHCl₃/EtOH 40:1, H₂O-saturated), to give β-anomer **2b**. The benzoylcytosine derivative **18** was obtained as a 1:1 mixture of anomers in 64% yield by treatment of **14** with bis-TMS-benzoylcytosine at 40°C for 1h, employing TMSOTf (1.8 eq.). The anomers were separated, after conversion of **18** to the thioacetate **19** (87%), by HPLC (silica gel, EtOAc/hexane/H₂O 7:3:0.07) to furnish pure β-anomer **3b**. Uracil derivative **20** (as a 1:1 mixture of anomers) was synthesized by reaction of **14** with bis-TMS-Uracil and TMSOTf (0.3 eq.) for 15 min. at room temperature (75%). Conversion of **20** to thioacetate **21** (78%), followed by HPLC-resolution of the anomers (silica gel, EtOAc/hexane/H₂O 7:3:0.07) afforded pure β-anomer **4b**.¹⁰

Structure assignments: The N⁷- and N⁹-isomers of the purine nucleoside analogs were distinguished by NMR spectroscopy. Uniformly in N-alkylated purines, the H-8 and H-1' proton signals and the C-4, C-8 and C-1' carbon signals of the N⁷-isomers are shifted downfield relative to the corresponding resonances of the N⁹-isomers, while the signal for C-5 of the N⁷-isomer is more shielded relative to the signal of the N⁹-isomer.¹¹ In the case of the analogs of guanosine, the N⁷-isomers **22** and the N⁹-isomers **17** were isolated. The α- and β-anomers of **22** were separated by crystallization from CH₂Cl₂/pentane, and the β-anomer converted to thioacetate β-**23**. The H-8 and H-1' proton signals and the C-4, C-8 and C-1' carbon signals of the chromatographically more mobile compound, assigned as the N⁷-isomer **23**, are downfield, and the C-5 resonance upfield relative to the corresponding resonances of its isomer **2b**, assigned as the the N⁹-isomer.

For the adenine derivative **1b**, the ¹³C NMR spectrum of **1b** was compared to the spectrum of N⁶-benzoyl-2'-deoxyadenosine and other benzoylated adenosines.¹² The chemical shifts of the purine ¹³C signals of **2b** and N⁶-benzoyl-2'-deoxyadenosine are similar within a range of 1.5 ppm, but differ, in the manner outlined above, from the chemical shifts reported for N⁷-ribofuranosyl-adenine.¹³

The anomeric configurations of compounds and **1b-4b** were assigned by NOE-difference spectroscopy. In all cases irradiation at H-1' gave a significant enhancement of the H-4' and H-2'α protons for the compound assigned the β-anomeric configuration, whereas the isomers assigned the α-anomeric configuration showed significant enhancement upon irradiation of H-1' only at H-2'β. The anomeric configuration of the N⁷-isomer **23** was tentatively assigned as β from the chemical shift of the H-4' signal. It was found that for all nucleoside analogs synthesized here, the H-4' signal of the α-anomer is shifted downfield relative to the signal of the β-anomer, presumably because of the anisotropic effect of the bases in the α-position. In the case of **3b** and **4b**, upon irradiation of H-1' NOE-enhancement is also found at H-6, which confirms that the N¹-isomers were obtained.

In conclusion, building blocks **1b-4b** have been synthesized as the necessary first step for the synthesis of analogs of oligonucleotides having -CH₂-S-CH₂-, -CH₂-SO-CH₂-, and -CH₂-SO₂-CH₂- units replacing

the $-O-PO_2-O-$ groups in natural oligonucleotides. Preliminary work on the coupling of these building blocks has produced tetramers in satisfactory yield. Progress in this area will be reported separately.

Notes and References

1. See for example: Jones, A. S.; MacCoss, M.; Walker, R. T. *Biochem. Biophys. Acta* **1973**, *365*, 365-377. Mungall, W. S.; Kaiser, J. K. *J. Org. Chem.* **1977**, *42*, 703-706. Cormier, J. F.; Ogilvie, K. K. *Nucl. Acids Res.* **1988**, *16*, 4583-4594. Letsinger, R. L.; Bach, S. A.; Eadie J. S. *Nucl. Acids Res.* **1986**, *14*, 3487-3499. Miller, P. S.; Agris, C. H.; Aurelian, L.; Blake, K. R.; Murakami, A.; Reddy, M. P.; Spitz, S. A.; Ts'o, P.O.P. *Biochimie* **1985**, *67*, 769-776. Moody, H. M.; van Genderen, M. H. P.; Koole, L. H.; Kocken, H. J. M.; Meijer, E. M.; Buck, H. M. *Nucl. Acids Res.* **1989** *17*, 4769-4782.
2. Paterson, B. M.; Roberts, B. E.; Kuff, E. L. *Proc. Nat. Acad. Sci. U.S.A.* **1977**, *74*, 4370-4374. Stephenson, M. L.; Zamecnik, P. C. *Proc. Nat. Acad. Sci. U.S.A.* **1978**, *75*, 285-288.
3. Nambiar, K. P.; Stackhouse, J.; Stauffer, D. M.; Kennedy, W. P.; Eldredge, J. K.; Benner, S. A. *Science*, **1984**, *223*, 1299-1301.
4. See for example: Schaller, H.; Weimann, G.; Lerch, B.; Khorana, H. G. *J. Am. Chem. Soc.* **1963**, *85*, 3821-3827.
5. Blancou, H.; Casadevall, E. *Tetrahedron* **1976**, *32*, 2907-2913.
6. See for example: Gais, H. J.; Lukas, K. L.; Ball, W. A.; Braun, S.; Lindner, H. J. *Liebigs Ann. Chem.* **1986** 687-716.
7. Schneider, K. C.; Huang, Z.; Benner, S. A., manuscript in preparation.
8. Vorbrüggen, H.; Bennua, B. *Chem. Ber.* **1981**, *114*, 1269-1286.
9. Volante, R. P. *Tetrahedron Letters* **1981**, *22*, 3119-3122.
10. All compounds were fully characterized by elemental analysis and/or UV-, mass-, IR-, and NMR-spectroscopy: **1b**: 1H -NMR ($CDCl_3$) δ 1.98-2.21 (m, 2H), 2.32 (s, 3H), 2.57 (ddd, $J=7,9,13.5$ Hz, 1H), 2.78-2.99 (m, 3H, C_3' -H), 3.18 (ddd, $J=5, 5, 13.5$ Hz, 1H), 4.16 (td, $J=3.5,8.5$ Hz, 1H), 4.47 (m, 2H), 6.37 (dd, $J=3,7$ Hz, 1H), 7.45-7.65 (m, 6H), 8.04 (m, 4H), 8.29 (s, 1H), 8.81 (s, 1H), 9.01 (s, 1H); ^{13}C NMR ($CDCl_3$) δ 25.9, 30.6, 35.3, 35.9, 42.2, 64.4, 82.7, 85.2, 123.9, 127.9, 128.6, 128.8, 129.6, 132.7, 133.4, 133.7, 141.5, 149.6, 151.2, 152.5, 164.8, 166.3, 195.5. **2b**: 1H NMR ($CDCl_3$) δ 1.27, 1.28 (2d, $J=8.5$ Hz, 6H), 1.97, 2.12 (2m, 2H), 2.29 (s, 3 H), 2.40 (ddd, $J=13.5,8.5,7.5$ Hz, 1H), 2.72 (m, 2H), 2.95 (m, 2H, C_3' -H), 3.11 (ddd, $J=5,5,13.5$, 1H), 4.05 (ddd=td, $J=8,3.5$ Hz, 1H), 4.39 (mc, 2H), 6.05 (dd, $J=7.5,3.5$ Hz, 1H), 7.49 (mc, 2 H), 7.60 (mc, 1 H), 7.80 (s, 1H), 8.87 (s, 1H), 12.08 (s, 1H). ^{13}C NMR ($CDCl_3$) δ 19.0, 19.1, 25.6, 30.6, 34.7, 35.6, 36.2, 41.8, 64.9, 82.5, 84.7, 121.8, 128.5, 129.5, 133.3, 137.6, 147.7, 147.8, 155.8, 166.3, 179.5, 196.0. **3b**: 1H NMR ($CDCl_3$) δ 1.97-2.24 (2m, 2 H), 2.30-2.50 (s, m, 5H), 2.59 (13.5,8.6,5 Hz, 1H), 2.94 (ddd, $J=13.5,8.5,7.5$, 1H), 3.29 (ddd, $J=13.5,9.5,7$ Hz, 1H), 4.10 (td, $J=10,7,3$ Hz, 1H), 4.40 (m, 2H), 6.11 (dd, $J=6.5,3$ Hz, 1H), 7.44 (m, 7H), 7.91 (d, $J=8$ Hz, 2H), 8.02 (m, 2H), 8.20 (d, $J=7.5$ Hz, 1H). **4b**: 1H NMR ($CDCl_3$) δ 1.91-2.17(2m, 2H), 2.28 (m, 1H), 2.33 (s, 3H), 2.42 (m, 2H), 2.89, 3.18 (2mc, 2H), 3.99 (ddd=td, $J=4.5,4.5,1.5$ Hz, 1H), 4.38 (mc, 2H), 5.81 (d, $J=8$ Hz), 6.09 (dd, $J=6.5,4$ Hz, 1H), 7.26 (m, 2H), 7.39 (m, 1H), 7.56 (d, $J=8$ Hz), 8.02 (m, 2H), 9.04 (s, 1H). **23**: 1H NMR ($CDCl_3$) δ 1.22 (2d=t, $J=7$ Hz, 6 H), 1.99-2.17 (m, 2H), 2.32 (s, 3H), 2.50-2.64 (m, 3H), 3.00 (m, 2H), 3.21 (dddd=sept., $J=5,8.5,14$ Hz,1H), 4.12 (dt, 3,9 Hz, 1H), 4.37, 4.45 (2dd, $J=5,11$ Hz, 2H), 6.53 (t, $J=5$ Hz, 1H), 7.46 (m, 3H), 7.59 (m, 2H), 8.18 (s, 1H), 10.52 (s, 1H), 12.39 (s, 1H); ^{13}C NMR ($CDCl_3$) δ 19.1, 19.2, 26.1, 30.6, 35.2, 35.9, 38.3, 41.6, 64.2, 83.0, 128.6, 129.6, 133.4, 140.7, 148.1, 153.1, 158.0 (C-4), 166.3, 180.1, 195.5.
11. Chenon, M.-T.; Pugmire, R. J.; Grant, D. M.; Panzica, R. P.; Townsend, L. B. *J. Am. Chem. Soc.* **1975**, *97*, 4628-4636. Kjellberg, J.; Johansson, N. G., *Tetrahedron* **1986**, *42*, 6541-6544. Garner, P.; Ramakanth, S. *J. Org. Chem.* **1988**, *53*, 1294-1298.
12. Seela, F.; Ott, J.; Potter, B. V. L. *J. Am. Chem. Soc.* **1983** *105*, 5879-5886.
13. Rousseau, R. J.; Robins, R. K.; Townsend, L. B. *J. Am. Chem. Soc.* **1968**, *90*, 2661-2668.

(Received in Germany 16 October 1989)