

Two Useful Photolabile Surfaces for Solid-Phase Synthesis


Troy D. Ryba and Patrick G. Harran*

Department of Biochemistry, The University of Texas Southwestern Medical Center at Dallas, Dallas, Texas 75390-9038

pharra@biochem.swmed.edu

Received February 4, 2000

ABSTRACT



o-Nitrobenzyl-based photolabile surfaces **3** and **4** have been synthesized from 2-phenylcyclohexanone and aminopropylsiloxane-grafted controlled pore glass. The procedures are simple and inexpensive and generate materials whose minimally functionalized reactive chromophore should tolerate a range of subsequent chemistries. Time- and solvent-dependent analysis of photodegradation demonstrates a performance comparable to related photolabile systems.

We are attempting to develop an artificial analogy to small molecule, processive biosynthesis: one that operates through synthetic restructuring of functionalized polymers. In these experiments, surface-bound oligopeptides are arrayed as starting materials for synthesis rather than being individual end points. Therefore, a format which enables efficient assembly of immobilized, unprotected peptides and also provides for versatility in their manipulation is vital. Because radicals, carbenoids, and active oxidants will play a role in this chemistry, commonly used organic polymer matrices are not attractive solid supports. Moreover, the system is designed to interface with whole-cell and receptor-based activity screens. Synthetic linkers degradable with acid and/or base can complicate this objective which, in other contexts, has been accomplished using photolabile inserts and product release using light.¹ The same tactic seemed appropriate for the current application, and we first employed a material formed from aminopropylsiloxane-grafted controlled pore

glass (**1**, amino-CPG) and *o*-nitrobenzylamine **2** (Figure 1) for solid-phase peptide synthesis. This combination performs well; however, our synthesis of **2**² is time-consuming and inefficient. Herein, we describe syntheses and preliminary performance of related photolabile reagents **3** and **4**: materials which retain a reactive chromophore of relatively high





Figure 1.

(1) The preparation and utility of photolabile linkers has been reviewed: (a) Gordon, K.; Balasubramanian, S. *J. Chem. Technol. Biotechnol.* **1999**, *74*, 835–851. (b) Backes, B. J.; Ellman, J. A. *Curr. Opin. Chem. Biol.* **1997**, *1*, 86–93. (c) Lloyd-Williams, P.; Albericio, F.; Giralt, E. *Tetrahedron* **1993**, *49*, 11065–11133. (d) Pallai, V. N. R. *Synthesis* **1980**, 1–26.

oxidation potential (in comparison to commercial veratryl-based systems) and are produced concisely from a common precursor.

Commercial 2-phenylcyclohexanone is nitrated with 85% nitronium tetrafluoroborate to afford an excess of 2' isomer **5** (Scheme 1).³ Purified **5** is then either converted to



^a (a) NO₂BF₄, CH₃NO₂, 10 °C, 15 min (46% + 12% 4' isomer); (b) *m*-CPBA, Na₂HPO₄, CH₂Cl₂, rt, 3 h (99%); (c) H₂SO₄, NaN₃, PhH, rt, 12 h (66%); (d) (Boc)₂O, DMAP, THF, rt (91%); (e) LiOH, THF/H₂O (82% for **9**, 93% for **10**).

caprolactone **6** via Baeyer–Villiger oxidation (*m*-CPBA) or to caprolactam **7** via Schmidt rearrangement (H₂SO₄, NaN₃). In both instances, ring expansion proceeds regioselectively with migration of the more substituted cyclohexanone α-carbon (C₂).⁴ Lactone **6** is saponified and the resultant hydroxy acid (**10**) transferred onto the surface of amino-CPG (Biosearch Technologies) through amidation (TBTU,⁵ DIPEA, HOBt). The composite formed (**3**) is functionally identical to a known polystyrene-based resin (six steps from *o*-nitrobenzaldehyde)⁶ although the current preparation is advantageous in that sensitive organometallic reagents and protecting group manipulations are not required. To complete the amine congener of **3**, lactam **7** is N-acylated with di-

tert-butyl dicarbonate and the formed imide (**8**) treated with LiOH to generate ω-carbamoyl carboxylic acid **9**. Amidation of **9** with amino-CPG (TBTU, DIPEA, HOBt) followed by exposing the resin to anhydrous CF₃CO₂H gives target material **4**. On a larger (10–15 g) scale, the conversion of **5** to **9** is carried out without intermediate purifications.

The density of reactive sites presented by batches of **3** (50–60 μmol/g)⁷ and **4** (120–125 μmol/g) is determined by acylation with *N*-Fmoc-Gly-OH and quantifying the dibenzofulvene released upon DBU treatment (2% volume in DMF).⁸ Photolyses are performed as stirred resin suspensions (borosilicate vials) in degassed solvent (vide infra) using a 450-W medium-pressure mercury vapor lamp positioned 5–8 cm from the reaction vessel. No attempt is made to attenuate power levels⁹ although a long-pass filter (cut-on 348 nm, Oriel Corp.) is routinely utilized.

A cited advantage of veratryl-based photolabile inserts is an increased photocleavage rate, due to higher molar absorptivity, at wavelengths (>350 nm) desirable for combinatorial chemistry applications.¹⁰ Interestingly, photophysical measurements in solution indicate that this increase in absorptivity can, in certain instances, be offset by a marked decrease in quantum yield.¹¹ Immobilization introduces additional variables. For example, Holmes has demonstrated that the photodecomposition of **11** (Hg(Xe) arc) occurs with *t*_{1/2} = 0.66 min in pH 7.4 phosphate-buffered saline (Figure 2).^{10b} In contrast, a solid-phase analogy of the process (i.e.,




Figure 2.

photoinduced release of **13** from **12**) requires 3 h to reach 95% conversion. For comparison, resin **4** loaded with *N*-Fmoc-Gly-OH evolves *N*-Fmoc-Gly-NH₂ on photolysis (*p*-dioxane, Hg vapor lamp, ≥348 nm) wherein yield peaks at 75% after roughly 90 min (Figure 3).¹² In *p*-dioxane, the

(2) Compound **2** is prepared from known 2,2-dimethyl-3-(*tert*-butoxycarbonyl)amino-3-(2'-nitrophenyl)propionic acid. Sternson, S. M.; Schreiber, S. L. *Tetrahedron Lett.* **1998**, 39, 7451–7454.

(3) The use of 95% NO₂BF₄ (Aldrich) results in lower regioselectivity (ortho:para = 1.4:1). The 85% reagent (Aldrich) is contaminated primarily with NOBF₄. Reaction mixtures using 95% NO₂BF₄ can be doped with 10 mol % of NOBF₄ to reconstitute, in part, a more selective nitration mixture. Nonselective nitration of 2-phenylcyclohexanone has been described: Prager, R. H.; Tippet, J. M.; Ward, A. D. *Aust. J. Chem.* **1978**, 31, 1989–2001.

(4) The conversion of **5** to **7** generates trace byproducts, none of which account for more than 5% of unrecovered mass. It is possible that one of these materials is a regioisomeric lactam.

(5) TBTU = *O*-benzotriazol-1-yl-*N,N,N',N'*-tetramethyluronium tetrafluoroborate, DIPEA = *N,N*-diisopropylethylamine, HOBt = *N*-hydroxybenzotriazole.

(6) Rodebaugh, R.; Fraser-Reid, B.; Geysen, H. M. *Tetrahedron Lett.* **1997**, 38, 7653–7656.

(7) The discrepancy in available sites for CPG derivatized with **9** and **10** is unexplained. In solution, the condensation of **10** with *n*-BuNH₂ (TBTU, DIPEA, HOBt, DMF) affords the corresponding hydroxy amide in 82% isolated yield. Self-condensation and/or re-lactonized products are not detected. Moreover, there is no significant difference in immobilized yields using **9** or **10** on organic polymer supports (Tentagel, Argopore).

(8) Newcomb, W. S.; Deegan, T. L.; Miller, W.; Porco, J. A., Jr. *Technical Bulletin 012*; Argonaut Technologies Inc.: San Carlos, CA. This information can be downloaded from www.argotech.com.

(9) 983 mW/cm² at the 366 nm line (50 cm distance from the lamp).

(10) (a) Holmes, C. P.; Jones, D. G. *J. Org. Chem.* **1995**, 60, 2318–2319. (b) Holmes, C. P. *J. Org. Chem.* **1997**, 62, 2370–2380.

(11) Krafft, G. A.; Randall Sutton, W.; Cummings, R. T. *J. Am. Chem. Soc.* **1988**, 110, 301–303.




Figure 3. Time and solvent-dependent photorelease of Fmoc-Gly-NH₂ from **4** acylated with Fmoc-Gly-OH (450W Hg vapor lamp, rt, ≥ 348 nm). Aliquots (5 μ L) of reaction solution (10 mg of resin suspended in 0.5 mL of degassed *p*-dioxane) removed by syringe and analyzed by HPLC (4.6 \times 250 mm Vydac C₁₈, linear gradient 40 \rightarrow 60% 1:1 *i*-PrOH/CH₃CN in H₂O at 0.5 mL min⁻¹) using Fmoc-Gly-OH (Bachem) as external standard. Inset: The same photochemical experiment with varying solvents and product analysis after 2 h of irradiation: lane 1, *p*-dioxane; lane 2, 4:1 H₂O/MeOH; lane 3, CH₃CN; lane 4, 1:1 THF/H₂O; lane 5, THF.

purity of isolated product is >95% (¹H NMR and HPLC), although both yield and purity do vary with reaction medium. Of the solvents examined, *p*-dioxane has proven most generally effective (Figure 3). Hydroxyl-presenting particles **3** loaded with *N*-Fmoc-Gly-OH release unchanged acid in 77% yield (85% purity) after 2.0 h of photolysis in *p*-dioxane. This contrasts with the 12–24 h required to similarly

(12) It is common to detect residual dibenzofulvene, accounting for ~15% of the original loading capacity, when photolyzed glass particles (dark red after 2 h of irradiation) are re-subjected to DBU treatment.

photodegrade acylated **10** immobilized on grafted polystyrene (THF/H₂O).⁶

Solid-phase peptide synthesis on resin **4** is facile and efficient. Using Fmoc-protected monomers and TBTU-mediated couplings, semiautomated syntheses of four- and five-residue oligomers accommodate the incorporation of all proteinogenic and designed residues thus far examined. The glass matrix tolerates a corrosive reagent mixture (95% CF₃CO₂H, 5% thioanisole, 5% 1:1 HS(CH₂)₂SH/H₂O) used for side-chain deprotections. Protected Leu-enkephalin-amide (YGGFL-NH₂) is synthesized on **4** and, following exposure to the above mixture (3h, rt), photoreleased in 40% yield (93% per step average) and 88% purity (HPLC, Vydac C₁₈).

The combination of an inorganic matrix and a minimally functionalized linking group has advantages in our experiments that should also be useful in other applications. The resins load at low density, there are no issues of solvent-dependent swelling, nor is there a need to protect **3/4** from normal working light. The extent of their chemical and mechanical stability remains to be determined although strong aqueous base (NaOH, KOH) should be avoided. Finally, the use of **9** and **10** is not limited to immobilization on glass. The simplicity and low cost of synthesizing each from 2-phenylcyclohexanone will hopefully facilitate their use with other solid supports in the range of solid-phase chemistry now being explored by the community at large.

Acknowledgment. Financial support provided by the Advanced Research Program of the Texas Higher Education Coordinating Board, The Robert A. Welch Foundation, and junior faculty awards administered through the Howard Hughes Medical Institute and the University of Texas Southwestern Medical Center are gratefully acknowledged. P.G.H. is the recipient of an NSF-CAREER award.

Supporting Information Available: Experimental procedures, characterization data, and ¹H NMR spectra for **5–10**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0056224