mmol) in THF (20 mL) was stirred at ambient temperature for 10 min and then at reflux for 40 min. The progress of the reaction was monitored by TLC on Kieselgel (eluent, ether/hexane, 5/95), showing a yellow-green spot, 14/15 at $R_f 0.10$. Chromatography on alumina gel (eluent, ether/hexane, 20/80) yielded dark green crystals, 14/15 (0.1368 g, 0.12 mmol, 61%): decomp pt 118-120 °C; ¹³C NMR (CD₂Cl₂) δ 211 (br, exchanging molybdenum and cobalt carbonyls), 181.8 (ester carbonyl), 141-124 (phenyls and indenyl aromatic CH's), 94.2/93.7 (indenyl C-2), 84.7/84.5, 82.5/82.3 (indenyl C-1, C-3), 75.2 (C-3), 47.6/47.1 (C-4), 41.5/41.1 (C-2), 34.8 (C-6), 32.0 (C-1), 26.0/25.9 (C-8), 25.4 (CH₂P), 23.7 (CH₂As), 23.2 (C-5), 22.3 (C-7), 20.8 (C-10), 16.3 (C-9), 31 P NMR $(CD_2Cl_2, 300 \text{ K})$ 43.5 and 42.0 ppm; IR $(CH_2Cl_2) \nu_{CO}$ 2030 (w), 2010 (m), 1985 (vs), 1975 (vs), 1945 (m), 1910 (m), 1740 (sh), 1720 (s), 1635 (ester) cm⁻¹; FAB mass spectrum, m/z (relative intensity) $1052 (37, (M - 3CO)^{+}), 1024 (100, (M - 4CO)^{+}), 996 (7, (M - 4CO)^{+}))$ $5CO)^+$, 968 (6, (M - 6CO)⁺). Anal. Calcd for $C_{53}H_{50}O_8Co_2MoAsP$: C, 56.10; H, 4.44. Found: C, 56.36; H, 4.48.

CpMoCo₂(CO)₇(P(OMe)₃)CCO₂(menthyl) (16). A solution of 10 (0.184 g, 0.26 mmol) and trimethyl phosphite (0.032 mL, 0.26 mmol) was stirred together in THF (20 mL) for 47 h at room temperature. The reaction was monitored by TLC on Kieselgel (eluent, ether/hexane, 12/88) showing two spots: dark green 16, R_f 0.29; green 10, R_f 0.14. The solvent was removed and the resulting green oil separated by flash chromatography on silica gel (eluent, ether/hexane, 22/78). The product was obtained as a green oil, 16 (0.0836 g, 0.11 mmol, 42%): ³¹P NMR (CD₂Cl₂, 300 K) 162.4 and 157.4 ppm; IR (CH₂Cl₂) ν_{CO} 2055 (s), 2000 (vs), 1990 (sh), 1970 (sh), 1920 (m), 1865 (m), 1655 (ester) cm⁻¹. The FAB mass spectrum indicates the presence of a very small amount of disubstituted product. FAB mass spectrum: m/z (relative intensity) 797 (12) $(M + 1)^+$), 768 (10, $(M - CO)^+$); 740 (14, (M $(M - 2CO)^+$; 712 (44, $(M - 3CO)^+$), 684 (71, $(M - 4CO)^+$), 656 (19, $(M - 5CO)^+$), 628 (65, $(M - 6CO)^+$), 600 (25, $(M - 7CO)^+$). CpMoCo₂(CO)₆(P(OMe)₃)₂CCO₂(menthyl): FAB mass spectrum, m/z (relative intensity) 836 (5, (M - 2CO)⁺), 808 (7, (M - 3CO)⁺), 780 (25, $(M - 4CO)^+$), 752 (4, $(M - 5CO)^+$), 724 (11, $(M - 6CO)^+$). $CpMoCo_2(CO)_7(P(C_6H_{11})_3)CCO_2(menthyl)$ (17). Tricyclohexylphosphine (0.0780 g, 0.28 mmol) and 10 (0.1825 g, 0.26 mmol) were stirred together in THF (20 mL) at ambient temperature for 38 h. The progress of the reaction was monitored by TLC on Kieselgel (eluent, ether/hexane, 16/84) showing three spots: yellow-green, 17, $R_f 0.65$; UV visible, $P(C_6H_{11})_3$, $R_f 0.40$; dark green, 10, R_f 0.25. Flash chromatography on silica (eluent, ether/hexane, 7/93) was used to separate the product from the starting materials. A dark green solid, 17, was obtained (0.1436 g, 0.15 mmol, 58%): mp 62–64 °C; ³¹P NMR (CD₂Cl₂, 300 K) 52.0 and 50.9 ppm; IR (CH₂Cl₂) ν_{C0} 2090 (w), 2075 (w), 2050 (s), 2025 (s), 2000 (s), 1970 (m), 1930 (m), 1855 (w), 1725 (sh), 1645 (ester) cm⁻¹; FAB mass spectrum: m/z (relative intensity) 896 (5, (M - 2CO)⁺), 840 (8, $(M - 4CO)^+$, 784 (71, $(M - 6CO)^+$), 758 (100, $(M - 7CO)^+$, 728 $(5, (M - 5CO - C_6H_{11})^+), 702 (89, (M - 6CO - C_6H_{11})^+), 676 (27, 10)$ $(M - 7CO - C_6H_{11})^+)$, 648 (38, $(M - 5CO - 2C_6H_{11})^+)$. Anal. Calcd for C₄₂H₅₇O₉Co₂MoP: C, 53.06; H, 6.04. Found: C, 52.94; H, 6.28.

Acknowledgment. We thank the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research. Financial support from the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged. K.A.S. and D.T.C. thank NSERC for graduate fellowships. It is a pleasure to acknowledge Richard Perrier and Dr. Donald Hughes for assistance with the 2-D NMR spectra and Dr. Richard Smith of the Southern Ontario Regional Mass Spectrometry Centre for the FAB mass spectra.

Registry No. 7, 117707-13-8; 8, 117627-32-4; 9a, 117708-19-7; **9b**, 117627-38-0; **10**, 117707-15-0; **11**, 117627-33-5; **12** (isomer 1), 117627-34-6; 12 (isomer 2), 117707-16-1; 13, 117627-35-7; 14, 117627-36-8; 15, 117707-14-9; 16 (isomer 1), 117627-37-9; 16 (isomer 2), 117707-17-2; 17 (isomer 1), 117627-39-1; 17 (isomer 2), 117707-18-3; [Mo(CO)₃(*i*-PrCp)]₂, 105046-06-8; [Mo(CO)₃(C₉H₇)]₂, 12098-65-6; Co, 7440-48-4; Mo, 7439-98-7; ((+)-methyl)-O₂CCl₃, 117652-31-0; Co₂(CO)₈, 10210-68-1; (+)-menthol, 15356-60-2.

Synthesis and Chemistry of Thiagermiranes

Takeshi Tsumuraya, ^{1a} Sadao Sato, ^{1b} and Wataru Ando*, ^{1a}

Department of Chemistry, The University of Tsukuba, Tsukuba, Ibaraki 305, Japan, and Analytical and Metabolic Research Laboratories, Sankyo Co., Ltd., Shinagawa, Tokyo 140, Japan

Received April 26, 1988

Dimethylgermylene (1) reacts with thicketones 3 and 9 to give 1,3,2,4-dithiadigermolanes 4 and 10, the ultimate products of formation of thiagermiranes 6 and 11 and subsequent addition of dimethylgermanethione (7). Dimesitylgermylene (16) reacts with adamantanethione (3) to give 3-(2-adamantyl)-2.2-dimesityl-1,2-thiagermirane (18). Photolysis of thiagermirane 18 yields dimesitylgermylene (16) and adamantanethione (3) derived from germathiocarbonyl ylide 22. Oxidation of thiagermirane 18 by mCPBA gives 3-(2adamantyl)-4,4-dimesityl-1,2,4-oxathiagermetane S-oxide (27). X-ray analyses were done for thiagermirane 18 and 1,2,4-oxathiagermetane S-oxide 27.

Introduction

A great deal of attention both experimental and theoretical has been devoted to cyclic compounds containing metal and hetero atoms, most particularly cyclopropane and cyclobutane systems and their derivatives.²⁻⁷ Since the introduction of strained rings into a molecular environment can have a profound effect on the chemistry of the

 ^{(1) (}a) The University of Tsukuba. (b) Sankyo Co., Ltd.
 (2) (a) Gordon, M. S.; George, C. J. Am. Chem. Soc. 1984, 106, 609.
 (b) Grev, R. S.; Schaefer, H. F. J. Am. Chem. Soc. 1987, 109, 6577.

^{(3) (}a) Baudler, M.; Jongebloed, H. Z. Anorg. Allg. Chem. 1979, 458, (b) Hluchy, J.; Klingebiel, U. Angew. Chem., Int. Ed. Engl. 1982, 21, 301.
 (c) Ando, W.; Hamada, Y.; Sekiguchi, A.; Ueno, K. Tetrahedron Lett. 1982, 23, 5323.
 (d) Ando, W.; Hamada, Y.; Sekiguchi, A.; Ueno, K. Tetrahedron Lett. 1983, 24, 4033. (e) Michalczyk, M. J.; West, R.; Michl, J. J. Chem. Soc., Chem. Commun. 1984, 1525. (f) West, R.; De Young, D. J.; Haller, K. J. J. Am. Chem. Soc. 1985, 107, 4942. (g) Yokelson, H. B.; Millevolte, A. J.; Gillette, G. R.; West, R. J. Am. Chem. Soc. 1987, 109, 6865.



overall system, interest in strained cyclic compounds in which one or more carbons is replaced by metal and hetero atoms is likely to increase. Recently we prepared several strained cyclic molecules containing one or more silicon or germanium atoms with hetero atoms in the ring.^{3c,d,5e,6} In this paper, we report the first synthesis of thiagermiranes from the reactions of germylenes with thioketones.

Results and Discussion

Reactions of Dimethylgermylene (1) with Thioketones.^{6a} When 7,7-dimethyl-7-germanorbornadiene 2⁸ (1 mmol) was decomposed in the presence of adamantanethione (3)⁹ (2 mmol) in benzene at 70 °C, a major product was 1,3,2,4-dithiadigermolane 4 (38%), along with adamantylideneadamantane $(5)^{10}$ (4%) and 1,2,3,4-tetraphenylnaphthalene (eq 1). Most of this reaction can be



explained in quite conventional fashion: dimethyl-

(4) (a) Lambert, R. L.; Seyferth, D. J. Am. Chem. Soc. 1972, 94, 9246. (b) Conlin, R. T.; Gaspar, P. P. J. Am. Chem. Soc. 1976, 98, 3715. (c) Masamune, S.; Hanzawa, Y.; Murakami, S.; Bally, T.; Blount, J. F. J. Am. Chem. Soc. 1982, 104, 1150. (d) Ishikawa, M.; Sugisawa, H.; Kumada, M.; Higuchi, T.; Matsui, K.; Hirotsu, K.; Iyoda, J. Organometallics 1983, 2, 174. (e) Masamune, S.; Murakami, S.; Tobita, H.; Williams, D. J. J. Am. Chem. Soc. 1983, 105, 7776.

(5) (a) Baudler, M. Pure Appl. Chem. 1980, 52, 755. (b) Krebs, A.; Berndt, J. Tetrahedron Lett. 1983, 24, 4083. (c) Masamune, S.; Hanzawa, Y.; Williams, D. J. J. Am. Chem. Soc. 1982, 104, 6136. (d) Snow, J. T.; Murakami, S.; Masamune, S.; Williams, D. J. Tetrahedron Lett. 1984, 25, 4191. (e) Ando, W.; Tsumuraya, T.; Goto, M. Tetrahedron Lett. 1986, 27. 5105.

(6) (a) Ando, W.; Tsumuraya, T.; Sekiguchi, A. Tetrahedron Lett. 1985, 26, 4523. (b) Ando, W.; Tsumuraya, T . Tetrahedron Lett. 1986, 27, 3251. (c) Tsumuraya, T.; Sato, S.; Ando, W. Organometallics, 1988, 7, 2015

(7) Ando, W.; Tsumuraya, T. J. Chem. Soc., Chem. Commun. 1987, 1514

(8) Schriewer, M.; Neumann, W. P. J. Am. Chem. Soc. 1983, 105, 897.
(9) Greidanus, J. W. Can. J. Chem. 1979, 48, 2530.
(10) Geluk, H. W. Synthesis 1970, 652.



Table I. Yield of the Products in Thermolysis of 3 and 2

	yield (%)			
mole ratio $(3/2)$	12	4	5	
0.46	52	5		
1.2	60	6		
4.4	12	33	5	

germylene (1) adds to adamantanethione (3) to form thiagermirane 6, which reacts with another molecule of germylene to eliminate sulfur to form dimethylgermanethione (7). 7 reacts with 6 to give 4 (Scheme I).^{11,12} Adamantylideneadamantane (5) is probably derived from pseudo Wittig-type reaction of 3 and germaethene 8.13

The results obtained in the reaction of dimethylgermylene (1) with 1,1,3,3-tetramethyl-2-indanthione $(9)^{14}$ were similar to those with adamantanethione (3). The main product was 1,3,2,4-dithiadigermolane 10 (11%), presumably derived from the reaction of dimethylgermanethione (7) and thiagermirane 11 (Scheme II).

Although dimethylgermylene (1) gives a significant amount of polymeric products under the reaction conditions, 15 2 was copyrolyzed with adamantanethione (3) at 250 °C to give 1,2,3-thiadigermetane 12 along with 1,3,2,4-dithiadigermolane 4 (eq 2). The yield of 12 was



⁽¹¹⁾ Reaction of thiirane with germylene was reported to yield the similar insertion product: Barrau, J.; Bouchaut, M.; Lavayssiere, H.; Dousse, G.; Satgé, J. J. Organomet. Chem. 1983, 243, 281.

(12) It is well-known that dimethylgermanethione (7) trimerizes very quickly to (Me₂GeS)₃, but we could not detect this trimer.

(13) Barrau, J.; Hamida, N. B.; Satgé, J. J. Organomet. Chem. 1985, 282, 315.

(14) Steliou, K.; Mrani, M. J. Am. Chem. Soc. 1982, 104, 3104.

(15) Kocher, J.; Neumann, W. P. Organometallics 1985, 4, 400.

Synthesis and Chemistry of Thiagermiranes

maximum at equal mole ratio of 2 and 3 but decreased with increasing concentration of 3 as shown in Table I. In this system dimethylgermylene (1) dimerizes to form tetramethyldigermene which is captured by adamantanethione (3) to give 12. Another conceivable route is germylene insertion into thiagermirane intermediate 6.16

Reactions of Diarylgermylenes with Adamantanethione (3).^{6b} Thermolysis of a benzene solution of 7,7diphenyl-7-germanorbornadiene 13 (0.6 mmol) and adamantanethione (3) (2 mmol) at 70 °C gave ethoxygermane 14, a product of the addition of ethanol to thiagermirane 15 in 50% yield after treatment by ethanol at room temperature (Scheme III). These results suggest that thiagermirane 15 is stable at room temperature in the absence of alcohols. However the isolation of 15 failed probably because it was unstable toward moisture or oxygen.

We next tried the reaction of a bulky substituted germylene: dimesitylgermylene (16) with adamantanethione (3). When a cyclohexane solution of hexamesitylcyclotrigermane (17)⁷ (0.4 mmol) and adamantanethione (3) (1.2 mmol) in a quartz tube was irradiated at room temperature with a low-pressure mercury lamp for 2.5 h, the major products were 3-(2-adamantyl)-2,2-dimesityl-1,2-thiagermirane (18) (53%), 1,2,3-thiadigermetane 19 (11%), adamantanethione dimer 20 (37%), and unreacted 17 (35%) (eq 3). The structure of 18 was confirmed by ¹H



and ¹³C NMR and mass spectroscopic analyses and elemental analysis. It is noted that the chemical shift attributed to the ring carbon (56.3 ppm) is similar to that of the corresponding thiasilirane (53.6 ppm). Thiagermirane 18 is stable in an atmosphere containing oxygen or moisture and does not decompose even when heated to its melting point (mp 208-209 °C).

Compound 18 presumably arises from the direct reaction of dimesitylgermylene (16) with adamantanethione (3). Tetramesityldigermene generated in the photolysis of cyclotrigermane 17 reacted with adamantanethione (3) to yield 1,2,3-thiadigermetane 19 as the [2 + 2] addition product. In a separate experiment tetramesityldigermene, prepared from the photolysis of cyclotrigermane 17, reacted with adamantanethione at room temperature to yield 1,2,3-thiadigermetane 19.

A single-crystal X-ray diffraction study of 18 confirmed the proposed structure. The asymmetric unit contains two crystallographically independent molecules (molecules A and B), and they were quite similar in conformation, bond angles, and bond lengths. Their molecular structure with atomic labeling schemes are shown in Figure 1.



Molecule A



Molecule B

Figure 1. Molecular conformation and atomic labeling of 18.

The bond lengths in the three-membered rings are 2.222 (1) Å for Ge–S bonds, 1.973 (5) and 1.970 (5) Å for Ge–C bonds, and 1.880 (5) and 1.877 (5) Å for S–C bonds. The endocyclic bond angles of Ge, C, S are 52.9 (1), 70.4 (1), and 56.8 (1)° for molecule A and 52.8 (1), 70.5 (1), and 56.7 (1)° for molecule B, respectively. The two aryl substituents form dihedral angles of 107.7 (1) and 66.2 (2)° for molecule A and 104.7 (2) and 65.4 (1)° for molecule B with the central thiagermirane ring. The dihedral angles between mesityl rings are 77.3 (2) and 73.1 (2)° for molecules A and B. The independent Ge–C_{Ar} bond lengths are 1.956 (4) and 1.958 (4) Å for molecule A and 1.954 (5) and 1.969 (5) Å for molecule B. Other bond lengths and significant bond angles for 18 are listed in Tables II and III.

Photolysis and Pyrolysis of Thiagermirane 18. Irradiation of a cyclohexane solution of 18 (0.2 mmol) and 2,3-dimethyl-1,3-butadiene (2 mmol) with a low-pressure mercury lamp at room temperature produced germacyclopentene 21^{17} and adamantanethione dimer 20, both in 37% yield. Decomposition of 18 at 495 °C under flowing nitrogen yielded 21 (34%) and adamantanethione (3) (29%).

Since 21 appears to be derived from the addition of dimesitylgermylene (16) to 2,3-dimethyl-1,3-butadiene, it is clear that 18 undergoes cycloelimination on photolysis or pyrolysis to produce dimesitylgermylene (16) and adamantanethione (3) either via the simultaneous cleavage of both the Ge-C and Ge-S bonds or via the germathio-carbonyl ylide 22 (Scheme IV).

Since the matrix-isolation technique has been used widely for the observation of the reactive species,¹⁸ we tried

^{(16) (}a) Barrau, J.; Bouchaut, M.; Lavayssiere, L.; Dousse, G.; Satgé, J. Helv. Chim. Acta 1979, 62, 152. (b) Riviere, P.; Riviere, M.; Satgé, J. "In Comprehensive Organometallic Chemistry; Pergamon: Oxford, 1983; Chapter 10, p 399.

⁽¹⁷⁾ Riviere, P.; Castel, A.; Satgé, J. J. Organomet. Chem. 1982, 232, 123.

Table II. Bond Lengths (Å) in 18 (Esd's in Parentheses)

molecule A		molecule B	
Ge(1)-S(1)	2.222 (1)	Ge(2)-S(2)	2.222 (1)
Ge(1)-C(1)	1.956 (4)	Ge(2)-C(31)	1.954(5)
Ge(1) - C(7)	1.958 (4)	Ge(2)-C(37)	1.969 (5)
Ge(1)-C(13)	1.973 (5)	Ge(2) - C(43)	1.970 (5)
S(1)-C(13)	1.880 (5)	S(2)-C(43)	1.877 (5)
C(1)-C(2)	1.387 (6)	C(31)-C(32)	1.395 (7)
C(1) - C(6)	1.425(7)	C(31)-C(36)	1.430 (7)
C(2)-C(3)	1.385(6)	C(32)-C(33)	1.372(7)
C(2)-C(2')	1.516 (7)	C(32)-C(32')	1.515 (8)
C(3)-C(4)	1.389 (8)	C(33)~C(34)	1.394 (8)
C(4) - C(5)	1.381(7)	C(34)-C(35)	1.374 (8)
C(4)-C(4')	1.522(7)	C(34)-C(34')	1.511 (8)
C(5) - C(6)	1.391 (6)	C(35)-C(36)	1.380(7)
C(6)-C(6')	1.509 (7)	C(36)-C(36')	1.504(7)
C(7) - C(8)	1.409(7)	C(37) - C(38)	1.385 (7)
C(7)-C(12)	1.407 (6)	C(37)-C(42)	1.414 (7)
C(8) - C(9)	1.403(7)	C(38)-C(39)	1.397 (8)
C(8)-C(8')	1.513(7)	C(38)-C(38')	1.526 (8)
C(9) - C(10)	1.383(7)	C(39)-C(40)	1.398 (9)
C(10)-C(11)	1.378 (8)	C(40) - C(41)	1.356 (8)
C(10) - C(10')	1.522(8)	C(40) - C(40')	1.519 (9)
C(11)-C(12)	1.401 (7)	C(41)-C(42)	1.403 (8)
C(12)-C(12')	1.521(7)	C(42)-C(42')	1.510 (8)
C(13)-C(14)	1.523(7)	C(43)-C(44)	1.509 (7)
C(13) - C(18)	1.523(6)	C(43) - C(48)	1.529(7)
C(14) - C(15)	1.553 (8)	C(44) - C(45)	1.537 (8)
C(14)-C(21)	1.545 (8)	C(44) - C(51)	1.540 (8)
C(15)-C(16)	1.525(7)	C(45) - C(46)	1.519 (8)
C(16) - C(17)	1.527 (8)	C(46) - C(47)	1.522 (8)
C(16) - C(22)	1.526 (8)	C(46) - C(52)	1.544 (9)
C(17) - C(18)	1.533 (7)	C(47) - C(48)	1.550 (8)
C(18) - C(19)	1.539 (8)	C(48) - C(49)	1.545 (8)
C(19) - C(20)	1.547 (8)	C(49) - C(50)	1.517 (9)
C(20) - C(21)	1.535 (7)	C(50) - C(51)	1.531 (8)
C(20)-C(22)	1.527 (8)	C(50) - C(52)	1.532 (9)

 Table III. Significant Bond Angles (deg) in 18 (Esd's in Parentheses)

molecule A		molecule B	
S(1)-Ge(1)-C(1)	110.1 (1)	S(2)-Ge(2)-C(31)	112.2 (1)
S(1)-Ge(1)-C(7)	119.1 (1)	S(2)-Ge(2)-C(37)	118.6 (1)
S(1)-Ge(1)-C(13)	52.9 (1)	S(2)-Ge(2)-C(43)	52.8 (1)
C(1)-Ge(1)-C(7)	115.1(1)	C(31)-Ge(2)-C(37)	115.4 (1)
C(1)-Ge(1)-C(13)	126.2(1)	C(31)-Ge(2)-C(43)	128.6(1)
C(7)-Ge(1)-C(13)	116.9 (1)	C(37)-Ge(2)-C(43)	113.4 (1)
Ge(1)-S(1)-C(13)	56.8 (1)	Ge(2)-S(2)-C(43)	56.7 (1)
Ge(1)-C(1)-C(6)	116.7(2)	Ge(2)-C(31)-C(36)	115.8(2)
Ge(1)-C(1)-C(2)	124.6 (3)	Ge(2)-C(31)-C(32)	126.3 (3)
C(2)-C(1)-C(6)	118.5(2)	C(32)-C(31)-C(36)	117.9 (3)
Ge(1)-C(7)-C(8)	115.0(2)	Ge(2)-C(37)-C(38)	126.7(2)
Ge(1)-C(7)-C(12)	126.1(2)	Ge(2)-C(37)-C(42)	114.0(2)
C(8)-C(7)-C(12)	118.9 (3)	C(38)-C(37)-C(42)	119.2 (2)
C(13)-C(18)-C(17)	109.4(2)	C(43)-C(48)-C(47)	108.8 (2)
C(13)-C(18)-C(19)	109.1 (2)	C(43)-C(48)-C(49)	109.1 (2)
C(17)-C(18)-C(19)	109.5 (3)	C(47)-C(48)-C(49)	108.6 (3)
Ge(1)-C(13)-C(14)	126.6 (2)	S(2)-C(43)-C(44)	114.8 (1)
Ge(1)-C(13)-C(18)	116.1 (2)	S(2)-C(43)-C(48)	111.5 (2)
S(1)-C(13)-C(14)	113.4(1)	Ge(2)-C(43)-S(2)	70.5(1)
S(1)-C(13)-C(18)	112.3(2)	Ge(2)-C(43)-C(44)	126.4(2)
Ge(1)-C(13)-S(1)	70.4(1)	Ge(2)-C(43)-C(48)	115.1(2)
C(14)-C(13)-C(18)	110.8 (3)	C(44)-C(43)-C(48)	111.3(3)
C(13)-C(14)-C(15)	110.0(2)	C(43)-C(44)-C(45)	108.5(2)
C(13)-C(14)-C(21)	108.0 (2)	C(43)-C(44)-C(51)	109.4 (2)
C(15)-C(14)-C(21)	108.2 (3)	C(45)-C(44)-C(51)	107.9 (3)

the photochemical decomposition of 18 in a hydrocarbon matrix at 77 K. Irradiation of thiagermirane 18 in 3methylpentane (3-MP) matrix at 77 K with a low-pressure mercury lamp led to the appearance of a new band at 690





Figure 2. UV-vis spectrum in 3-MP at 77 K: (A) prior photolysis of 18; (B) after photolysis with 254 nm for 30 min, 77 K; (C) after annealing and recooling to 77 K.







nm in the UV-vis spectrum, and the matrix became blue in color (Figure 2). This absorption band was indefinitely stable at 77 K.

After careful annealing of the matrix and recooling, this band diminished and a new band was formed with λ_{max} at 406 nm, which was assigned to tetramesityldigermene.¹⁹ Complete warming to room temperature resulted in disappearance of this band. Tetramesityldigermene probably reacted with adamantanethione at room temperature to yield 19. In the presence of 2,3-dimethyl-1,3-butadiene in 3-MP glass, melting of the matrix gave addition product 21.

Recently, silacarbonyl or silathiocarbonyl ylides 23 or 24 have been detected spectroscopically from the photolysis of oxasilirane 25 or thiasilirane 26 in a hydrocarbon matrix at 77 K.²⁰ On the basis of these results, it is quite reasonable to assume that the blue intermediate is most

⁽¹⁹⁾ Dimesitylgermylene shows an absorption band at 550 nm in 3-MP matrix at 77 K and dimerizes to tetramesityldigermene (λ_{max} 406 nm at 77 K) on annealing the matrix: Ando, W.; Tsumuraya, T.; Sekiguchi, A. Chem. Lett. 1987, 317.

⁽²⁰⁾ Ando, W.; Hagiwara, K.; Sekiguchi, A. Organometallics 1987, 6, 2270.

⁽²¹⁾ Gray, M. D. M.; Russel, D. D. R.; Smith, D. J. H.; Durst, T.; Gimbarzevsky, B. J. Chem. Soc., Perkin Trans. 1 1981, 1826.

Synthesis and Chemistry of Thiagermiranes



probably germathiocarbonyl ylide 22, which decomposes thermally to dimesitylgermylene (16) and adamantanethione (3) as shown in Scheme V.

Oxidation of 18 by m-Chloroperbenzoic Acid. Thiagermirane 18 is fairly air-stable but is easily oxidized by treatment with *m*-chloroperbenzoic acid (mCPBA). The reaction of 18 with 1 equiv of mCPBA in dichloromethane at 0 °C gave a colorless crystalline compound, 3-(2-adamantyl)-4,4-dimesityl-1,2,4-oxathiagermetane Soxide (27), in 50% yield, along with a 50% yield of unreacted 18. When 2 equiv of mCPBA were used under the same conditions, 18 was completely consumed and quantitative yield of 27 was obtained. 27 was further oxidized by the treatment with mCPBA to afford 3-(2adamantyl)-4,4-dimesityl-1,2,4-oxathiagermetane S,S-dioxide (28). Although thiagermirane S-oxide 29 was not isolated, the formation of compound 27 can be explained by either direct oxidation of 29 or oxidation of its rearrangement product 30 (Scheme VI). The structure of compound 27 was confirmed by an X-ray diffraction study, and the molecular structure with an atomic labeling scheme is shown in Figure 3. The four-membered ring consisting of O1, S, C, and Ge has a puckered structure in which the dihedral angle formed by plane Ge, S, C and plane Ge, S, O1 is 24.3 (3)°. The bond lengths in the four-membered ring are 1.876 (3) Å for the Ge–O1 bond, 2.033 (5) Å for the Ge-C bond, 1.844 (4) Å for the S-C bond, and 1.603 (4) Å for the S–O1 bond. The slightly long Ge–C bond length is probably due to the steric repulsion of bulky adamantyl and mesityl groups. The S-O2 bond length (1.461 (5) Å) is consistent with that of 1,2-oxathietane S-oxide 31 (1.466 Å).²¹ The endo cyclic bond



angles of Ge, S, C, and O1 atoms are 77.9 (1), 90.8 (1), 86.7 (1), and 99.7 (1)°, respectively. Other bond lengths and angles for 27 are listed in Tables IV and V.

Experimental Section

General Procedure. All reactions were carried out under an atmosphere of dry argon. The ¹H NMR spectra were recorded on Varian EM 360A and JEOL PMX 60 SI spectrometers. ¹³C NMR spectra were recorded on a JEOL FX-90Q spectrometer. Mass spectra were obtained on a Hitachi RMU-6M mass spectrometer. UV spectra were measured in cyclohexane on a Shimazu UV-365 spectrometer. IR spectra were recorded on a Hitachi 260-50 spectrometer. Preparative HPLC was carried out on a LC-08 (Japan Analytical Ind. Co., Ltd.). Preparative gas chromatographic separations were carried out on a Ohkura gas chromatograph with a 8 mm × 1.5 m glass column of 10% SF-96 on Celite 545. All melting points were uncorrected.

Materials. 2,3-Benzo-7,7-dimethyl-1,4,5,6-tetraphenyl-7germanorborna-2,5-diene (2) was prepared according to the literature procedures.⁸ 2,3-Benzo-1,4,5,6,7,7-hexaphenyl-7germanorborna-2,5-diene (13) was synthesized by the reaction of 1,1,2,3,4,5-hexaphenylgermole and benzyne generated by 1-



Figure 3. Molecular conformation and atomic labeling of 27.



Table IV. Bond Lengths (Å) in 27 (Esd's in Parentheses)

GeO(1)	1.876 (3)	C(8)–C(8')	1.505 (8)
Ge-C(1)	1.959 (5)	C(9) - C(10)	1.399 (9)
Ge-C(7)	1.971 (5)	C(10)-C(11)	1.369 (7)
Ge-C(13)	2.033 (5)	C(10)-C(10')	1.512 (8)
S-O(1)	1.603 (4)	C(11)-C(12)	1.390 (7)
S-O(2)	1.461 (5)	C(12)-C(12')	1.513 (7)
S-C(13)	1.844 (4)	C(13)-C(14)	1.542 (8)
C(1)-C(2)	1.405 (7)	C(13)-C(18)	1.551(7)
C(1)–C(6)	1.427 (8)	C(14) - C(15)	1.531 (8)
C(2) - C(3)	1.390 (7)	C(14)–C(19)	1.537 (7)
C(2)-C(2')	1.506 (8)	C(15)-C(16)	1.534 (8)
C(3) - C(4)	1.382 (9)	C(16)-C(17)	1.521 (9)
C(4) - C(5)	1.378 (8)	C(16)-C(21)	1.531 (8)
C(4)-C(4')	1.507 (8)	C(17)-C(18)	1.535 (8)
C(5) - C(6)	1.385 (7)	C(18)-C(22)	1.557 (7)
C(6)–C(6')	1.513(7)	C(19)-C(20)	1.526 (8)
C(7)–C(8)	1.395 (7)	C(20)-C(21)	1.534 (9)
C(7)-C(12)	1.398 (8)	C(20)-C(22)	1.530 (9)
C(8)-C(9)	1.399 (7)		

Table V. Significant Bond Angles (deg) in 27 (Esd's in Parentheses)

,				
111.3 (1)	Ge-C(7)-C(8)	116.0 (1)		
102.6 (1)	C(8)-C(7)-C(12)	119.8 (3)		
77.9 (1)	Ge-C(13)-C(18)	108.5 (3)		
113.4 (1)	S-C(13)-C(14)	111.6 (3)		
112.8 (1)	S-C(13)-C(18)	110.3 (1)		
129.8 (1)	Ge-C(13)-S	86.7 (1)		
109.3 (1)	Ge-C(13)-C(14)	128.8 (2)		
90.8 (1)	C(14)-C(13)-C(18)	108.6 (3)		
105.1 (1)	C(13)-C(14)-C(15)	108.4 (2)		
99.7 (1)	C(13)-C(14)-C(19)	110.0 (4)		
117.2 (2)	C(15)-C(14)-C(19)	107.9 (3)		
123.9 (4)	C(13)-C(18)-C(17)	110.6 (4)		
118.6 (5)	C(13)-C(18)-C(22)	109.1 (2)		
124.1 (3)	C(17)-C(18)-C(22)	108.2 (3)		
	111.3 (1) 102.6 (1) 77.9 (1) 113.4 (1) 112.8 (1) 129.8 (1) 109.3 (1) 90.8 (1) 105.1 (1) 99.7 (1) 117.2 (2) 123.9 (4) 118.6 (5) 124.1 (3)	$\begin{array}{c ccccc} 111.3 & (1) & Ge-C(7)-C(8) \\ 102.6 & (1) & C(8)-C(7)-C(12) \\ 77.9 & (1) & Ge-C(13)-C(14) \\ 112.8 & (1) & S-C(13)-C(14) \\ 112.8 & (1) & S-C(13)-C(18) \\ 129.8 & (1) & Ge-C(13)-S \\ 109.3 & (1) & Ge-C(13)-C(14) \\ 90.8 & (1) & C(14)-C(13)-C(18) \\ 105.1 & (1) & C(13)-C(14)-C(15) \\ 99.7 & (1) & C(13)-C(14)-C(19) \\ 117.2 & (2) & C(15)-C(14)-C(19) \\ 117.2 & (2) & C(15)-C(14)-C(19) \\ 123.9 & (4) & C(13)-C(18)-C(17) \\ 118.6 & (5) & C(13)-C(18)-C(22) \\ 124.1 & (3) & C(17)-C(18)-C(22) \\ \end{array}$		

aminobenzotriazole and Pb(OAc)₄.²² Hexamesitylcyclotrigermane (17) was prepared by reductive coupling of Mes₂GeCl₂ with Mg and MgBr₂ in THF as reported previously.⁷ Adamantanethione (**3**)⁹ and 1,1,3,3-tetramethyl-2-indanthione (**9**)¹⁴ were synthesized according to the literature procedures. Benzene, cyclohexane, and 3-methylpentane used as solvents were dried over lithium aluminium hydride and distilled before use. Dichloromethane was dried over calcium chloride and then distilled.

Thermolysis of 2,3-Benzo-7,7-dimethyl-1,4,5,6-tetraphenyl-7-germanorborna-2,5-diene (2) with Adamantanethione (3). A solution of 2 (535 mg, 1 mmol), 3 (332 mg, 2 mmol), and benzene (7 mL) was heated at 70 °C for 3 h. Separation of the reaction mixture by preparative GLC gave 4 (38%) and 5 (4%). 4: colorless oil; ¹H NMR (CCl₄) δ 0.72 (s, 6 H), 0.82 (s, 6 H), 1.3-2.9 (m, 14 H); ¹³C NMR (CDCl₃) δ 5.9 (q), 9.0 (q), 27.1 (d), 27.9 (d), 34.3 (t), 36.7 (d), 36.8 (t), 38.9 (t), 63.4 (s); MS, m/e (relative intensity) 404 (22, M⁺), 389 (50, M⁺ - 15), 223 (M⁺ -Ad=S - 15), 119 (98, Me₃Ge). HR-MS Calcd for C₁₄H₂₈S₂⁷²Ge⁷⁴Ge: 403.9909. Found: 403.9920.

All spectral data obtained for 5 were identical with those of an authentic sample. $^{10}\,$

Thermolysis of 2 with 1,1,3,3-Tetramethyl-2-indanthione (9). A solution of 2 (535 mg, 1 mmol), 9 (408 mg, 2 mmol), and benzene (7 mL) was heated at 70 °C for 3 h. Separation of the reaction mixture by preparative GLC gave 10 in 11% yield. 10: mp 144.5–145.5 °C; ¹H NMR (CCl₄) δ 0.78 (s, 6 H), 0.92 (s, 6 H), 1.56 (s, 12 H), 7.20 (s, 4 H); MS, m/e (relative intensity) 442 (40, M⁺), 427 (29, M⁺ – 15), 255 (21, M⁺ – C₁₃H₁₆ – 15), 238 (43, M⁺ – C₁₃H₁₆S), 223 (59, M⁺ – C₁₃H₁₆S – 15), 119 (100, Me₃Ge). Anal. Calcd for C₁₇H₂₈S₂Ge₂: C, 46.22; H, 6.38. Found: C, 46.23; H, 6.40.

Flow Pyrolysis of 2 with 3. The pyrolysis apparatus consisted of a 28 cm \times 1 cm Pyrex tube packed with Pyrex chips. The upper end of the tube was equipped with a rubber cap for syringe introduction of the sample and a nitrogen inlet. The pyrolysis tube was maintained at 250 °C, and the nitrogen flow was ca. 10 mL/min. The sample was introduced drop by drop using a syringe. The pyrolysates were trapped in a receiver cooled by dry ice-MeOH. A solution of 2 (53.5 mg, 0.1 mmol), 3 (7.6 mg, 0.046 mmol), and benzene (0.7 mL) was pyrolyzed at 250 °C under flowing nitrogen. Separation of the reaction mixture by preparative gas chromatography to give 12 (52%) and 4 (5 $\ddot{\%}$). The same reactions were carried out for 2 (0.1 mmol)/3 (0.44 mmol)and 2 (0.1 mmol)/3 (0.12 mmol). 12: colorless oil; ¹H NMR (CCl₄) δ 0.60 (s, 6 H), 0.70 (s, 6 H), 1.2-2.6 (m, 14 H); MS, m/e (relative intensity) 372 (16, M⁺), 357 (42, M⁺ - 15), 238 (16, Ad=GeMe₂), 223 (14, M⁺ - Ad - 15), 206 (9, Me₂Ge=GeMe₂), 166 (25, Ad=S), 119 (100, Me_3Ge). Elemental analysis could not be achieved since compound 12 rapidly decomposed under air. The reaction of 12 and sulfur in benzene at 80 °C gave 4, an insertion product of sulfur into a Ge-Ge bond of 12.

Thermolysis of 2,3-Benzo-1,4,5,6,7,7-hexaphenyl-7germanorborna-2,5-diene (13) with Adamantanethione (3). A solution of 13 (395 mg, 0.6 mmol), 3 (332 mg, 2 mmol), and benzene (5 mL) was heated at 70 °C for 3 h. Addition of ethanol at room temperature and separation by preparative HPLC gave colorless crystals of 14 in 50% yield. 14: mp 151-152 °C; ¹H NMR (CDCl₃) δ 1.14 (t, 3 H), 1.3-3.2 (m, 14 H), 1.40 (s, 1 H), 3.60 (q, 2 H), 7.6-8.3 (m, 10 H); ¹⁸C NMR (CDCl₃) δ 19.1 (q), 27.1 (d), 27.9 (d), 32.4 (t), 35.9 (t), 37.1 (d), 39.2 (t), 52.0 (s), 61.4 (t), 128.0 (d), 131.2 (d), 134.5 (s), 135.4 (d); MS, *m/e* (relative intensity) 394 (25, M⁺ - EtOH), 260 (6, Ph₂Ge=S), 228 (20, Ph₂Ge), 166 (100, Ad=S). Anal. Calcd for C₂₄H₃₀OSGe: C, 65.64; H, 6.88. Found: C, 65.42; H, 6.98.

Photolysis of Hexamesitylcyclotrigermane (17) with 3. A solution of 17 (373 mg, 0.4 mmol), 3 (199 mg, 1.2 mmol), and cyclohexane (5 mL) in a quartz tube was irradiated with a low-pressure mercury lamp at room temperature for 2.5 h. The resulting reaction mixture was concentrated and separated by preparative HPLC to give 18 (53%), 19 (11%), 20 (37%), and unreacted 17 (35%). 18; mp 208-209 °C; ¹H NMR (CDCl₃) δ 1.5-2.5 (m, 14 H), 2.23 (s, 6 H), 2.60 (s, 12 H), 6.87 (s, 4 H); ¹³C NMR (CDCl₃) δ 20.9 (q), 24.5 (q), 27.1 (d), 28.0 (d), 35.6 (t), 38.2

(t), 39.1 (t), 39.1 (d), 56.3 (s), 128.8 (d), 132.5 (s), 139.1 (s), 143.1 (s); MS, m/e (relative intensity) 478 (5, M^+), 344 (17, Mes₂Ge=S), 312 (29, Mes₂Ge), 166 (100, Ad=S). Anal. Calcd for C₂₈H₃₆SGe: C, 70.46; H, 7.60. Found: C, 70.67; H, 7.77. 19; mp 187–189 °C; ¹H NMR (CDCl₃) δ 1.2–2.0 (m, 14 H), 2.20 (s, 12 H), 2.40 (s, 24 H), 6.80 (s, 8 H); ¹³C NMR (CDCl₃) δ 20.8 (q), 26.0 (q), 27.1 (d), 28.0 (d), 37.1 (t), 38.2 (t), 39.4 (t), 40.0 (d), 66.2 (s), 129.2 (d), 137.1 (s), 137.6 (s), 141.5 (s), 143.8 (s), 145.5 (s); MS m/e (relative intensity) 431 (21, Mes₃Ge), 334 (9, Mes₂Ge=S), 312 (100, Mes₂Ge). Anal. Calcd for C₄₆H₅₈SGe₂: C, 70.09; H, 7.41. Found: C, 69.98; H, 7.48.

All spectral data obtained for 20^9 and 17^7 were identical with those of authentic samples.

Photolysis of 18 with 2,3-Dimethyl-1,3-butadiene. A solution of 18 (100 mg, 0.2 mmol), 2,3-dimethyl-1,3-butadiene (164 mg, 2 mmol), and cyclohexane (5 mL) was irradiated with a low-pressure mercury lamp at room temperature for 5 h. Separation of the reaction mixture by preparative HPLC gave 21 (39%) and 20 (39%). All spectral data obtained for 21^{17} and 20 were identical with those of authentic samples.

Pyrolysis of 18 with 2,3-Dimethyl-1,3-butadiene under Flowing Nitrogen. The pyrolysis was carried out by the same procedure used for flow pyrolysis of 2 and 3. A solution of 18 (47.7 mg, 0.1 mmol), 2,3-dimethyl-1,3-butadiene (410 mg, 5 mmol), and benzene (1 mL) was pyrolyzed at 495 °C under flowing nitrogen. Separation of the reaction mixture by preparative TLC (silica gel, benzene/hexane = 1/3) gave 21 (25%), 3 (21%), and unreacted 18 (27%).

All spectral data obtained for 21, 3, and 18 were identical with those of authentic samples.

Photolysis of 18 in 3-MP Matrix at 77 K. A mixture of 18 (ca. 1 mg) and 3-MP (5 mL) was placed in a quartz UV cell and degassed for several times. The mixture was cooled to 77 K, and the resulting matrix was irradiated with a low-pressure mercury lamp. The UV-vis absorptions were taken at periodic intervals. An absorption band with a maximum at 690 nm appeared and grew on irradiation as shown in Figure 2. On annealing the matrix this band immediately disappeared with concurrent formation of a new band with $\lambda_{max} = 406$ nm.

Photolysis of 18 with 2,3-Dimethyl-1,3-butadiene in 3-MP Matrix at 77 K. A mixture of 18 (ca. 5 mg), 2,3-dimethyl-1,3butadiene (0.5 mg), and 3-MP (5 mL) was placed in a quartz tube and degassed several times. The mixture was irradiated with a low-pressure mercury lamp for 10 min. The matrix turned to blue, and a broad band with $\lambda_{max} = 690$ nm was observed. When matrix was warmed to melt, the color disappeared, and 21 was detected by GLC-mass spectrometry.

Oxidation of 18 with mCPBA. To a mixture of 18 (47.7 mg, 0.1 mmol) and dichloromethane (3 mL) was added m-chloroperbenzoic acid (mCPBA) (0.1 mmol) at 0 °C. The reaction mixture was separated by preparative HPLC to give 27 (50%) and unreacted 18 (50%). Similar reaction of 18 (47.7 mg, 0.1 mmol) and mCPBA (0.2 mmol) gave quantitative yield of 27. 27 was further oxidized to 28 by the treatment with mCPBA at room temperature. 27: mp 225.5-227 °C; ¹H NMR (CDCl₃) δ 1.1-2.1 (m, 12 H), 2.1-2.6 (m, 18 H), 2.6-3.3 (m, 2 H), 6.88 (s, 4 H); ¹³C NMR (CDCl₃) δ 21.0 (q), 23.0 (q), 26.4 (d), 26.8 (d), 30.6 (d), 32.1 (d), 35.0 (t), 35.1 (t), 35.8 (t), 36.0 (t), 36.8 (t), 96.8 (s), 129.0 (d), 129.6 (d), 133.5 (s), 138.8 (s), 140.0 (s), 140.1 (s), 140.8 (s), 142.3 (s); IR (CDCl₃) 1120 cm⁻¹ (-SO-O-); MS, m/e (relative intensity) 446 (14, M⁺ - SO₂), 312 (100, Mes₂Ge). Anal. Calcd for C₂₈H₃₆SO₂Ge: C, 66.04; H, 7.12. Found; C, 66.08; H, 7.29. 28: $\begin{array}{l} & \text{mp} > 300 \ ^{\circ}\text{C}; \ ^{1}\text{H} \ \text{NMR} \ (\text{CDCl}_{3}) \ \delta \ 1.5 - 3.0 \ (\text{m}, \ 14 \ \text{H}), \ 2.27 \ (\text{s}, \ 6 \ \text{H}), \\ & 2.45 \ (\text{s}, \ 12 \ \text{H}), \ 6.90 \ (\text{s}, \ 4 \ \text{H}); \ ^{13}\text{C} \ \text{NMR} \ (\text{CDCl}_{3}) \ \delta \ 21.0 \ (\text{q}), \ 23.3 \ (\text{q}), \\ \end{array}$ 26.5 (d), 33.0 (d), 34.6 (t), 37.0 (t), 37.3 (t), 94.9 (s), 129.7 (d), 132.6 (s), 140.8 (s), 142.1 (s); IR (CDCl₃) 1305, 1155 cm⁻¹ ($-SO_2-O-$); MS, m/e (relative intensity) 526 (10, M⁺), 312 (100, Mes₂Ge). Anal. Calcd for C₂₈H₃₆SO₃Ge: C, 64.02; H, 6.90. Found: C, 63.77; H, 6.96.

X-ray Crystallography of 18 and 27. Crystals of dimensions $0.4 \times 0.4 \times 0.15$ mm for 18 and $0.4 \times 0.3 \times 0.2$ mm for 27, obtained from hexane, were used for the X-ray analysis. Lattice constants were determined by least-squares fit of angular settings of 20 reflections within range $30 < 2\theta < 50^{\circ}$. Intensity data were obtained on a Rigaku AFC-5R equipped with graphite-mono-chromatized Cu K α radiation and using the θ -2 θ scan technique

atom	x	У	z	B_{eq}
		Molecule A		
Ge(1)	1705.5 (2)	1204.7 (5)	3668.1(3)	2.79
S(1)	1498 (1)	1654 (1)	2507 (1)	4.32
C(1)	2484 (2)	680 (5)	3892 (2)	2.86
C(2)	2915 (2)	1426 (5)	4214 (3)	3.26
C(3)	3463 (2)	1040 (5)	4254 (3)	3.81
C(4)	3594 (2)	-111 (5)	3996 (3)	4.16
C(5)	3167 (2)	-897 (5)	3713 (3)	3.98
C(6)	2613 (2)	-526 (5)	3659 (3)	3.27
C(2')	2816 (2)	2682 (5)	4532 (3)	4.16
C(4')	4198 (2)	-502 (7)	4017 (4)	6.11
C(6')	2165 (2)	-1420(5)	3340 (3)	4.18
C(7)	1178 (2)	229 (4)	4128 (2)	2.74
C(8)	1253 (2)	302 (4)	4003 (3)	3.18
C(9)	910 (2) 501 (2)	-393 (3)	0200 (0) 4099 (9)	3.19 4.07
C(10)	496 (2)	-1100(0)	4522 (3)	4.07
C(11)	755(2)	-542(5)	3776(3)	3.40
C(8')	1697(2)	1107(5)	5306 (3)	4 35
C(10')	139(2)	-1929(6)	5352 (3)	5.34
C(12')	634(2)	-685 (5)	2963 (3)	4.31
C(13)	1436 (2)	2804 (4)	3241 (2)	3.05
C(14)	1780 (2)	3972 (5)	3193 (3)	3.74
C(15)	1742 (2)	4823 (5)	3848 (3)	4.53
C(16)	1134 (2)	5156 (5)	3872 (3)	4.62
C(17)	811 (2)	3961 (5)	3943 (3)	4.25
C(18)	832 (2)	3129 (5)	3288 (3)	3.54
C(19)	580 (2)	3824 (5)	2602 (3)	4.69
C(20)	923 (2)	5017 (5)	2538 (3)	5.00
C(21)	1532 (3)	4678 (5)	2505 (3)	5.15
C(22)	889 (3)	5443 (5)	3188 (3)	5.39
		Molecule B		
Ge(2)	1866.2(3)	1487.4 (5)	8558.6 (3)	3.47
S(2)	1602 (1)	1142 (1)	7394 (1)	4.48
C(31)	2630 (2)	2104 (5)	8766 (3)	3.54
C(32)	3097 (2)	1428 (5)	9070 (3)	4.18
C(33)	3618 (2)	1954 (5)	9156 (3)	4.64
C(34)	3704 (2)	3170 (6)	8947 (3)	4.80
C(35)	3245 (2)	3840 (5)	8650 (3)	4.34
C(36)	2712 (2)	3361 (5)	8000 (3)	3.67
C(32')	3049 (3)	100 (0)	9328 (3)	0.07 6.90
C(34)	4270 (3)	1182 (1)	8263 (3)	4.46
C(37)	1332(2)	2267(5)	9111 (3)	3.50
C(38)	864(2)	2944 (5)	8836 (3)	3.98
C(39)	498 (3)	3355 (6)	9291 (3)	5.20
C(40)	594 (3)	3078 (6)	10025 (3)	5.84
C(41)	1056 (3)	2425 (6)	10294 (3)	5.33
C(42)	1437 (2)	2010 (5)	9855 (3)	4.28
C(38')	707 (2)	3290 (6)	8046 (3)	4.82
C(40')	188 (3)	3481 (8)	10520 (4)	8.67
C(42')	1945 (3)	1294 (7)	10189 (3)	6.12
C(43)	1596 (2)	-83 (5)	8100 (3)	3.45
U(44)	1943 (2)	-1214 (5)	8003 (3)	4.41
C(45)	1668 (3)	-1819 (6)	7333 (3)	5.09
C(47)	1079 (3)	-2303 (6)	7410 (J) 7516 (9)	0.3Z 1 70
C(42)	1001 (2)	-1100 (0)	8192 (3)	3 01
C(40)	1021(2)	-1298 (6)	8852 (3)	5 75
C(50)	1356 (3)	-2451(6)	8743 (3)	6.58
C(51)	1951 (3)	-2078 (5)	8655 (3)	5.85
C(52)	1089 (3)	-3160 (6)	8074 (4)	6.65

 $(2\theta < 128^{\circ})$. During data collection three standards, measured before every 200 reflections, indicated no systematic variation of intensity with time. Of 8577 and 4171 independent reflections measured, only 7062 and 3687 for 18 and 27, respectively, were considered as observed on the basis of the criterion $F_o > 2\sigma(F_o)$. All intensities were corrected for Lorentz and polarization effects but not for absorption.

Crystal data for 18: C₂₈H₃₆SGe; mol wt 477.3; monoclinic; P2₁/n; a = 24.277 (5) Å, b = 10.789 (1) Å, c = 18.832 (2) Å, $\beta = 98.70$ (1)°; U = 4875.9 Å³; Z = 8; $\rho = 1.30$ g cm⁻³; μ (Cu K_a) = 27 cm⁻¹; F(000) = 2016; T = 297 K.

Table VII. Fractional Atomic Coordinates $(\times 10^4)$ and Thermal Parameters $(Å^2)$ for 27 (Esd's in Parentheses)

		<u>, , , , , , , , , , , , , , , , , , , </u>		
atom	x	У	2	B_{eq}
Ge	228 (1)	6235.6 (2)	7405 (1)	2.76
S	-1142 (2)	5862 (1)	4960 (2)	2.92
O(1)	-1286 (4)	5796 (1)	6438 (4)	2.73
O(2)	-2183 (5)	6245 (2)	4256 (4)	4.79
C(1)	1743 (6)	5939 (2)	8993 (5)	2.25
C(2)	2913 (6)	6234 (2)	9837 (5)	2.78
C(3)	4140 (7)	6038 (2)	10879 (6)	3.02
C(4)	4239 (7)	5553 (2)	11121 (6)	3.42
C(5)	3063 (7)	5266 (2)	10321 (6)	3.18
C(6)	1821 (6)	5439 (2)	9256 (5)	2.48
C(7)	-1047 (6)	6734 (2)	7868 (6)	2.48
C(8)	-1992 (7)	6586 (2)	8613 (6)	3.10
C(9)	-3031 (7)	6909 (2)	8881 (6)	3.54
C(10)	-3123 (7)	7381 (2)	8428 (7)	3.77
C(11)	-2147 (7)	7519 (2)	7733 (6)	3.63
C(12)	-1108 (7)	7208 (2)	7435 (6)	3.04
C(13)	792 (6)	6145 (2)	5691 (5)	2.31
C(14)	1078 (7)	6527 (2)	4737 (6)	3.24
C(15)	1078 (8)	6283 (2)	3427 (6)	4.41
C(16)	2370 (8)	5906 (2)	3694 (6)	4.36
C(17)	2105 (7)	5529 (2)	4644 (6)	3.42
C(18)	2107 (7)	5766 (2)	5969 (6)	2.86
C(19)	2691 (8)	6758 (2)	5385 (7)	4.45
C(20)	3995 (8)	6388 (2)	5661 (7)	4.32
C(21)	3963 (8)	6144 (3)	4339 (7)	5.12
C(22)	3720 (7)	6012 (2)	6620 (6)	3.52
C(2')	2872 (7)	6767 (2)	9647 (7)	3.64
C(4')	5572 (8)	5342 (3)	12252 (7)	5.05
C(6′)	610 (7)	5084 (2)	8462 (6)	3.28
C(8′)	-1979 (8)	6087 (2)	9146 (7)	4.39
C(10')	-4270 (8)	7723 (2)	8714 (8)	5.74
C(12')	-63 (8)	7399 (2)	6680 (7)	4.35

Crystal data for 27: C₂₈H₃₆O₂SGe; mol wt 509.3; monoclinic, P2₁/c; a = 9.040 (1) Å, b = 28.061 (3) Å, c = 10.454 (1) Å, $\beta = 109.02$ (1)°; U = 2507.0 Å³; Z = 4; $\rho = 1.35$ g cm⁻³; μ (Cu K_{α}) = 27 cm⁻¹; F(000) = 1072; T = 297 K.

Both structures 18 and 27 were solved by MULTAN84²³ and refined by block-diagonal least-squares methods. Positions of the hydrogen atoms were estimated from standard geometry. The final refinements with anisotropic temperature factors for the non-hydrogen atoms and isotropic temperature factors for the hydrogen atoms lowered R values to 0.056 ($R_w = 0.063$, W = $1/\sigma^2(F_o)$) and 0.061 ($R_w = 0.056$, $w = 1/\sigma^2(F_o)$), for 18 and 27, respectively. The goodness of fit values were 3.45 and 2.32, and maximum and minimum heights in the final difference Fourier syntheses were +0.2, -0.2 and +0.2, -0.3 e $Å^{-3}$ for 18 and 27, respectively. The maximum and average shifts in the final cycle of refinements were 0.33, 0.05 and 0.21, 0.03 of the corresponding standard deviations for 18 and 27, respectively. The molecular structures with atomic labeling schemes, the bond lengths, the selected bond angles, and atomic coordinates for the non-hydrogen atoms are given in Figure 1 and Tables II, III, and VI for 18 and in Figure 3 and Tables IV, V, and VII for 27, respectively.

Acknowledgment. This research was supported by a Grant-in-Aid for Scientific Research (No. 62115003) from the Ministry of Education, Science, and Culture of Japan.

Registry No. 2, 76054-64-3; 3, 23695-65-0; 4, 104144-59-4; 5, 30541-56-1; 9, 74768-58-4; 10, 104144-60-7; 12, 104144-61-8; 13, 107824-82-8; 14, 107824-83-9; 16, 72176-49-9; 17, 107824-84-0; 18, 107824-85-1; 19, 117308-60-8; 20, 31023-85-5; 21, 83686-73-1; 27, 117308-61-9; 28, 117308-62-0; 2,3-dimethyl-1,3-butadiene, 513-81-5.

Supplementary Material Available: Tables of bond angles, anisotropic thermal factors, and H-atom parameters for 18 and 27 (7 pages); listings of structure factors (43 pages). Ordering information is given on any current masthead page.

⁽²³⁾ Main, P.; Germain, G.; Woolfson, M. M. MULTAN84: A System of Computer Programs for the Automatic Data; Universities of York: York, England, and Louvain: Louvain Belgium, 1984.