

Unimolecular Alcoholysis of Organosilicon Halides of the Type $(\text{Me}_3\text{Si})_2\text{C}-(\text{SiMe}_2\text{OMe})(\text{SiR}'_2\text{X})$. Anchimeric Assistance by and Migration of the OMe Group

Colin Eaborn,* Paul D. Lickiss, Sabah T. Najim, and M. Novella Romanelli

School of Chemistry and Molecular Sciences, University of Sussex, Brighton BN1 9QJ, U.K.


The compound $\text{R}_2\text{C}(\text{SiMe}_2\text{OMe})(\text{SiMe}_2\text{Cl})$ ($\text{R} = \text{SiMe}_3$ throughout) reacts more rapidly with $\text{CF}_3\text{CH}_2\text{OH}$ than with MeOH , and $\text{R}_2\text{C}(\text{SiMe}_2\text{OMe})(\text{SiPh}_2\text{Br})$ reacts with EtOH to give exclusively the rearranged product $\text{R}_2\text{C}(\text{SiMe}_2\text{OEt})(\text{SiPh}_2\text{OMe})$, indicating that the rate-determining step of the solvolyses involves formation of a methoxy-bridged cation; migration of the OMe group also occurs in the reaction of the bromide with AgBF_4 .

Organosilicon iodides of the type $\text{R}_3\text{CSiR}'_2\text{I}$ ($\text{R} = \text{SiMe}_3$ throughout), in which steric hindrance greatly inhibits direct nucleophilic attack at the functional silicon centre, react with silver salts AgY to give wholly ($\text{R}' = \text{Ph}$) or in part ($\text{R}' = \text{Et}$) rearranged products of the type $\text{R}_2\text{C}(\text{SiR}'_2\text{Me})(\text{SiMe}_2\text{Y})$, and it is thought that the reactions proceed *via* the bridged cations (**1**, $\text{Z} = \text{Me}$). Analogous rearrangement occurs in the trifluoroacetolysis of $\text{R}_3\text{CSiEt}_2\text{I}$, which thus appears to involve rate-determining ionization to a bridged cation of type (**1**, $\text{Z} = \text{Me}$; $\text{R}' = \text{Et}$).¹ In these reactions the migrating Me group is assumed to provide anchimeric assistance to the leaving of I^- .

The methanolysis of $\text{R}_3\text{CSiMe}_2\text{X}$ compounds with $\text{X} = \text{I}$, OClO_3 , or OSO_2CF_3 , is known not to be significantly accelerated by the presence of NaOMe ,^{2,3} and so at first it seemed likely that this reaction also involved anchimerically-

assisted formation of the bridge cation (**1**; $\text{R}' = \text{Me}$), but (a) solvolysis was much slower in the more electrophilic solvent $\text{CF}_3\text{CH}_2\text{OH}$, in which ionization would be expected to occur more readily, and (b) the hydrolysis and methanolysis of $\text{R}_3\text{CSiEt}_2\text{I}$ were found to proceed without rearrangement.⁴ It thus appears that the reaction does not proceed through a cation, and that the solvent is nucleophilically involved in the rate-determining step,⁴ possibly in nucleophilic solvation in an $\text{S}_{\text{N}}2$ (intermediate) type of process.⁵

In the hope of observing rate-determining unimolecular cleavage of a silicon-halogen bond in alcoholysis without nucleophilic participation by the solvent we turned to compounds of the type $\text{R}_2\text{C}(\text{SiMe}_2\text{OMe})(\text{SiR}'_2\text{X})$, in which the γ -OMe group could be expected to lead to much more stable bridged ions of the type (**1**, $\text{Z} = \text{OMe}$). (Compare the



calculations by Kos and Schleyer quoted in ref. 6.) The methoxychloride $R_2C(SiMe_2OMe)(SiMe_2Cl)$ was previously shown to undergo methanolysis $> 10^6$ times as rapidly as R_3CSiMe_2Cl ,⁶ and we have now found that (a) its rate of reaction with 9:1 v/v MeOH–dioxane is not very sensitive to the presence of NaOMe (the half-lives, measured by n.m.r. spectroscopy, at 35 °C with 0.00, 0.08, 0.16, 0.32, and 0.64 M NaOMe were 7.5, 5.1, 4.4, 6.1, and 6.4 min, respectively), and (b) its reaction with 8:1 v/v CF_3CH_2OH –dioxane [in which the alcohol:dioxane molar ratio is similar to that in the mixture in (a)], to give $R_2C(SiMe_2OMe)(SiMe_2OCH_2CF_3)$, is markedly faster, being complete before the first measurement can be made, *i.e.* in < 45 s; there is a subsequent slower reaction, to give $R_2C(SiMe_2OCH_2CF_3)_2$, which does not occur in the presence of Et_3N .

We have furthermore found that the reaction of $R_2C(SiMe_2OMe)(SiPh_2Br)$ with EtOH for 5 min at room temperature gives a single product [m.p. 168 °C; δ_H (CCl_4) 0.10 (s, $SiMe_2$), 0.18 (s, $SiMe_3$), 1.32 (t, CH_2Me), 3.55 (s, OMe), 3.65 (q, CH_2), and 7.2–8.0 (m, Ph)], which has been shown by X-ray diffraction⁷ to be the rearranged species $R_2C(SiMe_2OEt)(SiPh_2OMe)$. (The same product is formed in the reaction with $AgClO_4$ in EtOH.) It can thus be assumed that the reactions proceed *via* cations of type (1, Z = OMe; $R' = Me$, or Ph), which are rapidly attacked by the solvent nucleophile at the less hindered end of the bridge. It seems that there can be rate-determining ionization in these crowded systems of the type $R_2C(SiMe_2Z)(SiR'_2X)$ if there is either a weak bridging group (Z) such as Me in association with a strongly electrophilic solvent such as CF_3CO_2H , or a powerful bridging group such as OMe in association with a more weakly

electrophilic solvent such as MeOH, and presumably some intermediate combinations would also be effective.

Predominant, but not complete, rearrangement occurred when $R_2C(SiMe_2OMe)(SiPh_2Br)$ was treated with $AgBF_4$ in Et_2O , the product being a 4:1 mixture of $R_2C(SiMe_2F)(SiPh_2OMe)$ [for $SiMe_2F$, δ_H 0.22 (d), δ_F –138.1 p.p.m.] and $R_2C(SiMe_2OMe)(SiPh_2F)$ [for $SiMe_2OMe$, δ_H 0.11 (s); for $SiPh_2F$, δ_F –161.3 p.p.m.].

The observation that with sufficient anchimeric assistance alcoholysis of organosilicon halides can take place without nucleophilic involvement of the solvent in the rate-determining step extends the analogy between mechanisms of substitution at a saturated carbon atom and those at a sterically hindered silicon atom.⁴ It will be of interest to see whether variations in the nature of the solvent affect such reactions of organosilicon halides in the way they do S_N1 reactions of organic halides.⁸

We thank the S.E.R.C. for support, the Iraqi Ministry of Higher Education and the University of Basrah for the award of a postgraduate scholarship (to S. T. N.), and Mr. N. H. Buttrus and Dr. P. B. Hitchcock for the X-ray structural determination.

Received, 14th August 1985; Com. 1218

References

- 1 C. Eaborn, D. A. R. Happer, S. P. Hopper, and K. D. Safa, *J. Organomet. Chem.*, 1980, **188**, 179.
- 2 C. Eaborn and F. M. S. Mahmoud, *J. Chem. Soc., Perkin Trans. 2*, 1981, 1309.
- 3 S. A. I. Al-Shali, C. Eaborn, and F. M. S. Mahmoud, *J. Organomet. Chem.*, 1982, **232**, 215.
- 4 S. A. I. Al-Shali, C. Eaborn, F. A. Fattah, and S. T. Najim, *J. Chem. Soc., Chem. Commun.*, 1984, 318.
- 5 T. W. Bentley and G. E. Carter, *J. Am. Chem. Soc.*, 1982, **104**, 5741.
- 6 C. Eaborn and D. E. Reed, *J. Chem. Soc., Chem. Commun.*, 1983, 495.
- 7 N. H. Buttrus and P. B. Hitchcock, personal communication.
- 8 T. W. Bentley, G. E. Carter, and K. Roberts, *J. Org. Chem.*, 1984, **49**, 5183.