

Preliminary communication

SYNTHESIS OF METALLOCHIRAL TRIORGANOBISMUTHINES

P. BRAS, A. VAN DER GEN and J. WOLTERS

*Gorlaeus Laboratories, Department of Organic Chemistry, University of Leiden,
 P.O. Box 9502, 2300 RA Leiden (The Netherlands)*


(Received July 27th, 1983)

Summary

Triorganobismuthines containing a chiral centre at the metal atom have been prepared by selective bromodearylation of mixed triarylbismuthines, followed by reaction of the resulting chiral diarylbromobismuthines with arylmagnesium bromides. The NMR spectra of some of the new compounds exhibit diastereotopic effects, associated with the chirality, indicating that the bismuthines have a stable pyramidal configuration.

Selective heterolytic cleavage of carbon—metal bonds has been successfully used in the synthesis of chiral organometallic compounds such as chiral tetraorganotin [1] and tetraorganolead [2] compounds. We have found that the reaction of certain mixed triarylbismuthines (I) with hydrogen bromide in methanol proceeds with sufficient selectivity to be used for the synthesis of chiral diarylbromobismuthines (II). Reaction of compounds II with arylmagnesium bromides provides a novel type of compound, the metalochiral triorganobismuthines (III) (Scheme 1).

The synthesis of II was carried out by adding slowly a solution of hydrogen bromide in methanol (0.2–0.5 M) at 0–20°C to a solution of I in dichloro-



SCHEME 1

TABLE 1

SYNTHESIS OF CHIRAL BISMUTHINES: $(Ar')_2(Ar'')Bi \xrightarrow[(2) Ar''MgBr]{(1) HBr/MeOH} (Ar')(Ar')(Ar'')Bi$

Starting compound	Product	Yield (%)	M.P. (°C (uncorr.))
Ia (4-BrC ₆ H ₄)(α-C ₁₀ H ₇) ₂ Bi ^{a,b}	IIIa (4-BrC ₆ H ₄)(C ₆ H ₅)(α-C ₁₀ H ₇)Bi	44	129–130
Ib (4-ClC ₆ H ₄)(2-CH ₃ C ₆ H ₄) ₂ Bi ^{a,c}	IIIb (4-ClC ₆ H ₄)[2-(CH ₃) ₂ CHOC ₆ H ₄](2-CH ₃ C ₆ H ₄)Bi	10	70–70.5
Ic (4-ClC ₆ H ₄)(α-C ₁₀ H ₇) ₂ Bi ^{a,d}	IIIc (4-ClC ₆ H ₄)[2-(CH ₃) ₂ CHOC ₆ H ₄](α-C ₁₀ H ₇)Bi	39	130–133
Id (4-ClC ₆ H ₄)[2,4,6-(CH ₃) ₃ C ₆ H ₂] ₂ Bi ^{a,e}	IIId (4-ClC ₆ H ₄)[2-(CH ₃) ₂ CHOC ₆ H ₄][2,4,6-(CH ₃) ₃ C ₆ H ₂]Bi	45	110–112
Id (4-ClC ₆ H ₄)[2,4,6-(CH ₃) ₃ C ₆ H ₂] ₂ Bi ^{a,e}	IIIe (4-ClC ₆ H ₄)(2-C ₆ H ₄ CH ₂ OC ₆ H ₄)[2,4,6-(CH ₃) ₃ C ₆ H ₂]Bi	55	103–104

^a Prepared by adding 2 eq. of ArMgBr to Ar'BiCl₂ in THF at 0° C; Ar'BiCl₂ was prepared from Ar'₂Bi and 2 eq. of BiCl₃ in toluene/ether [3]; (4-BrC₆H₄)₂Bi was synthesized from 4-BrC₆H₄Li and BiCl₃ in ether (yield 67%, m.p. 145° C (uncorr.)) (lit 145° C [3]). ^b Yield 67%, m.p. 133–136° C (uncorr.). ^c 78%, 80–83° C. ^d 63%, 135° C; ^e 50%, 130–131° C.

methane or chloroform (0.2–0.5 M). Bismuthine IIb ($X = \text{Cl}$, $R = 2\text{-CH}_3$) precipitated after partial evaporation of the solvent and was obtained pure by filtration. The other bismuthines, sometimes as a mixture with the hydrocarbon formed, were isolated by complete evaporation of the solvent. The products thus obtained were used directly in the next step.

Compounds III were synthesized by treating II in tetrahydrofuran at 0°C with an excess of arylmagnesium bromide. The bismuthines were purified by crystallization from chloroform/ethanol and their structures were established by ^1H and ^{13}C NMR spectroscopy. The results are presented in Table 1*.

The selectivity of the reaction with hydrogen bromide is in agreement with the variations in reactivity found for electrophilic substitution in benzenoid compounds containing $(R)_n\text{E}$ ($\text{E} = \text{Si}, \text{Ge}, \text{Sn}, \text{Pb}, \text{Hg}$ etc.) as the leaving group [4].

Some PMR data for the prochiral isopropoxy group of IIIb,c,d and the prochiral benzyloxy group of IIIe are listed in Table 2. For both groups diastereotopic effects are observed as a consequence of the chirality of the bismuthines.

TABLE 2
PMR DATA FOR TRIARYLBISMUTHINES

TABLE 2

PMR DATA FOR TRIARYLBISMUTHINES

III_{b,c,d}

III_e

Bismuthine	R	Solvent	$\delta\text{-C(CH}_3)_2$ ^a (ppm, vs TMS)		$\Delta\delta$ (ppm)
IIIb	2-CH ₃	CDCl ₃	1.20	1.19 ^b	0.015
IIIc	2,3-benzo	CDCl ₃	1.17 ^b		0
IIId	2,4,6-(CH ₃) ₃	CDCl ₃	1.25	1.14 ^b	0.112
IIId	2,4,6-(CH ₃) ₃	C ₆ D ₆	0.87	0.99 ^c	0.12
IIId	2,4,6-(CH ₃) ₃	acetone-d ₆	1.19	1.05 ^c	0.14
IIId	2,4,6-(CH ₃) ₃	DMSO-d ₆	1.11	0.96 ^c	0.15
			$\delta(\text{CH}_2)$ ^d (ppm, vs TMS)		
IIIe		CDCl ₃	5.03	5.01 ^b	0.021

^a $^3J(\text{H-H})$ 6 Hz. ^b Bruker WM-300. ^c JEOL FS 100. ^d $^2J(\text{H-H})$ 12 Hz.

The influence of *ortho*-substitution (R) in the other aryl group of the bismuthine on the diastereotopic effect is remarkable. For the methyl protons of the isopropoxy group of IIIb ($R = 2\text{-CH}_3$) this effect is small ($\Delta\delta$ 0.015 ppm in CDCl₃), and in IIc ($R = 2,3\text{-benzo}$) it could not even be observed. In IIId ($R = 2,4,6\text{-(CH}_3)_3$), however, the effect is significant ($\Delta\delta$ 0.11 ppm in CDCl₃, 0.15 ppm in DMSO-d₆).

The ^{13}C spectrum of IIId also exhibits a diastereotopic effect for the methyl carbons of the isopropoxy group ($\Delta\delta$ 0.27 ppm). For the methylene protons of IIIe a diastereotopic effect of 0.021 ppm and a geminal coupling constant of 12 Hz is found (in CDCl₃).

*This is not a general method. In some cases (e.g. $X = \text{Cl}$, $R = 4\text{-CH}_3$, $R' = 4\text{-CH}_3\text{O}$ or $4\text{-C}_2\text{H}_5$) this reaction leads to inseparable mixtures, consisting of at least three different triaryl bismuthines, as was shown by 300 MHz PMR. Possibly redistribution reactions occur in these cases (e.g. $2 \text{Ar}_2\text{BiX} \rightarrow \text{Ar}_3\text{Bi} + \text{ArBiX}_2$ or $\text{Ar}_2\text{Ar}'\text{Bi} + \text{Ar}_2\text{BiX} \rightarrow \text{ArAr}'\text{BiX} + \text{Ar}_3\text{Bi}$).

These data strongly suggest that these chiral bismuthines possess a stable, pyramidal configuration.

More detailed information on the synthesis and the spectral properties of this novel class of organobismuth compounds will be published separately.

References

- 1 M. Gielen, *Acc. Chem. Research*, (1973) 198.
- 2 H.O. van der Kooi and J. Wolters, *Third Int. Conf. Organometal. Coord. Chem. Ge, Sn, Pb, Dortmund* (1980) A29; J. Wolters and D. de Vos, *J. Organometal Chem. Library*, 13 (1982) 647.
- 3 M. Wieber, *Gmelin Handbuch der Anorganischen Chemie*, Bd 47, *Bismutorganische Verbindungen*, Springer Verlag, Berlin 1977.
- 4 R.O.C. Norman and R. Taylor, *Electrophilic substitution in benzenoid compounds*, Elsevier Publishing Company, Amsterdam 1965.