Synthesis of planar chiral tricarbonylcyclopentadienylmanganese complexes using a Mn(CO)₃⁺ transfer reaction[†]

Seung Uk Son,^{*a*} Kang Hyun Park,^{*a*} Seung Jung Lee,^{*a*} Young Keun Chung^{**a*} and Dwight A. Sweigart^{**b*}

^a School of Chemistry and Center for Molecular Catalysis, Seoul National University, Seoul 151-747, Korea. E-mail: ykchung@plaza.snu.ac

^b Department of Chemistry, Brown University, Providence, RI 02912, USA

Received (in Cambridge, UK) 22nd March 2001, Accepted 30th May 2001 First published as an Advance Article on the web 27th June 2001

Planar chiral ligand transfer reactions can be carried out by the reaction of $[(naphthalene)Mn(CO)_3]BF_4$ with planar chiral ferrocene derivatives.

Ligand transfer from one metal to another is a useful reaction when the complex cannot be easily prepared by conventional methods and is popular in organometallic chemistry.¹ Among these, cyclopentadienyl transfer reactions may be of synthetic interest,^{2,3} in particular if cheap substrates such as commercially available ferrocene derivatives can be used as starting materials. However, there have been no reports of ligand transfer reactions from ferrocene derivatives to a manganese moiety to prepare cyclopentadienyl manganese complexes. Such complexes have many notable features such as high thermal stability, lack of substantial air or water sensitivity, kinetic inertness toward substitution reactions, and stereochemical robustness. These characteristics will be useful properties when manganese complexes are used as auxiliaries in other reactions.

Herein we report the synthesis of planar chiral cyclopentadienyltricarbonylmanganese complexes from the reaction of planar chiral ferrocenes with $[(\eta^6-naphthalene)Mn(CO)_3]BF_4^4$ and their application in asymmetric reactions. Planar chiral chelates of CpMn(CO)_3 derivatives are quite rare and hence their use in asymmetric reactions has not been developed. There has been a report⁵ of enantioselective allylic substitution of cyclic substrates by catalysis with palladium complexes of *P*,*N*chelate ligands attached to a CpMn(CO)_3 unit.

Refluxing equimolar amounts of a planar chiral ferrocene derivative **1a** and $[(\eta^6-naphthalene)Mn(CO)_3]BF_4$ in the presence of 4 Å molecular sieves in CH₂Cl₂, followed by standard work-up procedures, led to a 46% yield of the air-stable, planar chiral tricarbonylcyclopentadienylmanganese complex **2a** [eqn. (1)].[‡]



As far as we are aware, this is the first use of a ligand transfer reaction in the synthesis of a planar chiral tricarbonylcyclopentadienylmanganese complex. Interestingly, formation of CpMn(CO)₃ was not observed in the reaction products. Birch *et al.*⁶ first tested the idea of chirality transfer in the reaction between a chiral donor complex and an unsymmetrical diene ligand. Since then, there have been many reports on the synthesis of chiral organometallic compounds.⁷ However, there have been no known reactions on the synthesis of tricarbonylcyclopentadienylmanganese derivatives. Surprisingly, during the transfer reaction, a *complete inversion* of the absolute configuration occurred [eqn. (1)]. This suggests that a bimetallic species [Cp–Fe–Cp–Mn(CO)₃]⁺ **A**, in which one of the Cp rings

www.rsc.org/chemcomm



Acid hydrolysis of 2a yielded the known compound 3 [eqn. (2)];¹⁰ according to the ¹H NMR spectrum of 2a and the optical rotation of 3, compound 3 is optically pure.

2

This journal is © The Royal Society of Chemistry 2001

$$a \xrightarrow{H^*} \overbrace{\underset{(CO)_3}{Mn}}^{Kn} CHO$$
(2)

We investigated whether the triol group in 2 could be used as a directing group in the functionalization of the cyclopentadienyl ring. Deprotonation of 2a by Bu⁴Li followed by the addition of PPh₂Cl yielded 4 in 67% yield [eqn. (3)].

$$2a \xrightarrow{Bu'Li, Ph_2PCI} \underbrace{We}_{Mn PPh_2} O \xrightarrow{O}_{MeO} (3)$$

The structure of **4** was confirmed by an X-ray crystal structure determination (Fig. 1).¹¹ Thus, the diphenylphosphino group was successfully introduced with the aid of the triol group.

We next investigated the use of the ligand transfer reaction in the synthesis of *P*,*N*-ligands and studied palladium-catalyzed asymmetric allylic alkylation. The use of ligands derived from (arene)Cr(CO)₃ as a chiral chelating ligand is quite popular.¹² However, the use of CpMn(CO)₃ as a chiral chelating ligand is quite rare.⁵

[†] Electronic supplementary information (ESI) available: characterization of **2b**, **2c**, **4**, **6**, **7**, **8** and **9**, and a typical procedure for Pd-catalyzed allylic alkylation. See http://www.rsc.org/suppdata/cc/b1/b102693n/

The planar chiral P,N-ligands **9** were synthesized from **5** [eqn. (4)].



Ligand transfer reaction of **5** afforded **6** in 74% yield. Deprotonation of **6** followed by the addition of PPh₂Cl and subsequent reaction with acid gave **8** in 75% yield. The condensation reaction of **8** with *tert*-butyl amine gave a *P*,*N*ligand **9a** in 96% yield. The de of **7** prepared from **6** was 75%. However, after recrystallization, the de of **7** was >99.5%.

The palladium-catalyzed asymmetric allylic alkylation of *rac*-1,3-diphenylprop-2-en-1-yl acetate with dimethyl malonate was successfully carried out [eqn. (5)].

Reaction of *rac*-1,3-diphenylprop-2-en-1-yl acetate with dimethyl malonate in the presence of 1 mol% Pd catalyst at 15 °C for 4 h gave the allylic alkylation product methyl-2-carbomethoxy-3,5-diphenylpent-4-enolate in high yields and high ee values (95% yield and 89% ee in CH₂Cl₂; 96% yield and 92% ee in DMSO). Lowering the reaction temperature to -20 °C in CH₂Cl₂ led to an increase of the ee value to 94% with a 74% yield.

In conclusion, we have conducted the first demonstration that a planar chiral ligand transfer reaction can be carried out by the reaction of $[(naphthalene)Mn(CO)_3]BF_4$ with planar chiral ferrocene derivatives and we have shown that the *P*,*N*-ligand prepared in this study can be employed in palladium-catalyzed asymmetric allylic alkylation.



Fig. 1 X-Ray structure of 4.

Notes and references

 $\ddagger Compounds~(\eta^6-C_{10}H_8)Mn(CO)_3]BF_4,~1$ and 5 were synthesized as previously described 4a,13,14

Synthesis of **2a**: a Schlenk flask containing 0.50 g of molecular sieves was flame-dried. After the flask was cooled to r.t., **1a** (0.35 g, 1.06 mmol), $[(\eta^{c}-C_{10}H_8)Mn(CO)_3]BF_4$ (0.40 g, 1.13 mmol), and 15 mL of CH₂Cl₂ were added to the flask. The resulting solution was heated at reflux for 18 h. After the solution was cooled to r.t., the solution was filtered over a pad of Celite. The filtrate was evaporated to dryness and chromatographed on a silica gel column eluting with hexane and diethyl ether (10:1 v/v). Removal of the solvent gave **2a** (oil, 46%, 0.17 g). IR v(CO) 2012, 1920 cm⁻¹; $\delta_{H}(300 \text{ MHz}, d_6\text{-benzene}, TMS) 5.21 (br s, 1 H), 5.03 (br s, 1 H), 4.15 (br s, 1 H), 4.07 (dd, 3.7, 11.0 Hz, 1 H), 3.85 (m, 1 H), 3.60 (m, 2 H), 3.56 (m, 1 H), 3.55 (s, 3 H), 2.05 (s. 3 H), 1.87 (m, 1 H), 1.26 (d, 13.0 Hz, 1 H); <math>\delta_{C}(75 \text{ MHz}, d_6\text{-benzene}) 225.7, 102.8, 99.5, 96.4, 82.6, 81.1, 79.8, 76.1, 75.3, 66.6, 59.1, 28.0, 12.1; HRMS calc. for C₁₅H₁₆MnO₆:$ *m*/z 349.0484; obs: 349.0483. [a)²D² = -36 (c 0.42 in CH₂Cl₂).

- A. Z. Rubezhov and S. P. Gubin, Adv. Organomet. Chem., 1972, 10, 347; A. Efraty, J. Organomet. Chem., 1973, 57, 1; P. E. Garrou, Adv. Organomet. Chem., 1984, 23, 95.
- 2 S. Top, C. Lescop, J.-S. Lehm and G. Jaouen, J. Organomet. Chem., 2000, **593–594**, 167; R. L. Halterman, Chem. Rev., 1992, **92**, 965.
- 3 T. W. Spradau and J. A. Katzenellenbogen, Organometallics, 1998, 17, 2009; M. Wenzel, J. Labelled Compd. Radiopharm., 1992, 31, 641.
- 4 (a) S. Sun, L. K. Yeung, D. A. Sweigart, T.-Y. Lee, S. S. Lee, Y. K. Chung, S. R. Switzer and R. D. Pike, *Organometallics*, 1995, 14, 2613;
 (b) J. E. Kim, J. S. U. Son, S. S. Lee and Y. K. Chung, *Inorg. Chim. Acta*, 1998, 281, 229;
 (c) I. S. Lee, H. M. Seo and Y. K. Chung, *Organometallics*, 1999, 18, 1091.
- 5 S. Kudis and G. Helmchen, Angew. Chem., Int. Ed., 1998, 37, 3047.
- 6 A. J. Birch, W. D. Raverty and G. R. Stephenson, *Tetraheron Lett.*, 1980, 197.
- 7 V. I. Sokolov, Chirality and Optical Activity of Organometallics Compounds, Gordon and Breach, New York, 1990; A. Togni, Metallocenes, ed. A. Togni and R. L. Halterman, Wiley-VCH: Weinheim, Germany, 1998, vol. 2.
- 8 E. J. Watson, K. L. Virkaitis, H. Li, A. J. Nowak, J. S. D'Acchioli, K. Yu, G. B. Carpenter, Y. K. Chung and D. A. Sweigart, *Chem. Commun.*, 2001, 457.
- 9 G. E. Herberich, U. Englert, F. Marken and P. Hofmann, *Organometallics*, 1993, **12**, 4039.
- 10 N. M. Loim, M. A. Kondratenkp and V. I. Sokolov, J. Org. Chem., 1994, 59, 7485; Y. Yamazaki, J. Chromatogr., 1991, 542, 129.
- 11 *Crystal data*: for **4**: monoclinic, space group *P*2₁; *a* = 8.0816(7), *b* = 15.0390(4), *c* = 10.9049(9) Å, β = 90.422(3)°, *V* = 1325.34(16) Å³, *Z* = 2, *D_c* = 1.334 Mg m⁻³; -10 < *h* < 10, -17 < *k* < 17, -14 < *l* < 13, *R*1 = 0.0379, *wR*2 = 0.10791. For **7**: orthorhombic, space group *P*2₁₂₁₂₁; *a* = 7.6659(2), *b* = 17.7774(8), *c* = 18.6479(10) Å; *V* = 2541.33(19) Å³, *Z* = 4; *D_c* = 1.355 Mg m⁻³; -9 < *h* < 9, -23 < *k* < 22, -24 < *l* < 24; *R*1 = 0.0526, *wR*2 = 0.1398. CCDC reference numbers 157495 and 157495. See http://www.rsc.org/supdata/cc/b1/b102693n/ for crystallographic data in CIF or other electronic format.
- 12 M. Uemura, R. Miyake and Y. Hayashi, J. Chem. Soc., Chem. Commun., 1991, 1696; S. B. Heaton and G. B. Jones, *Tetrahedron Lett.*, 1993, **33**, 1693; Y. Hayashi, H. Sakai, N. Kaneta and M. Uemura, J. Organomet. Chem., 1995, **503**, 143; H.-Y. Jang, H. Seo, J. W. Han and Y. K. Chung, *Tetrahedron Lett.*, 2000, **41**, 5083; C. Bolm and K. Muniz, Chem. Soc. Rev., 1999, **28**, 51.
- 13 O. Riant, O. Samuel, T. Flessner, S. Tandien and H. Kagan, J. Org. Chem., 1997, 62, 6733.
- 14 G. Hftime, J.-C. Daran, E. Manoury and G. G. A. Balavoine, J. Organomet. Chem., 1998, 565, 115.