


## Transition Metal Derivatives of Aryl Diazonium Ions. Part II.<sup>1</sup> Arylazo Derivatives of Substituted Iron Carbonyls

By William E. Carroll and Fergus J. Lalor,\* Department of Chemistry, University College, Cork, Ireland

Diazonium salts react with phosphine-substituted iron carbonyls,  $L_2Fe(CO)_3$ , to give stable orange arylazo derivatives of formula  $\{ArN_2Fe(CO)_2L_2\}^+BF_4^-$ . I.r., electronic, n.m.r. ( $^1H$ ,  $^{19}F$ , and  $^{31}P$ ), and Mössbauer spectroscopic data have been recorded for selected complexes. The  $\pi$ -acceptor character of the  $ArN_2$  ligand is discussed in the context of preliminary results of a structural determination.

ALTHOUGH the reaction of diazonium salts with transition metal carbonyls was first studied ten years ago,<sup>2</sup> it is only recently that stable arylazo derivatives of the transition metals have been isolated, but no derivative of a first-row transition metal has yet been described. Recent work in these laboratories has resulted in the development of useful synthetic routes to neutral arylazo complexes of cobalt and both neutral and ionic arylazo derivatives of iron. The present paper reports the synthesis and properties of one such class of compounds.



<sup>1</sup> Part I, F. J. Lalor and P. L. Pauson, *J. Organometallic Chem.*, 1970, **25**, C51.

<sup>2</sup> G. R. Schrauzer, *Chem. Ber.*, 1961, **94**, 1891.

Disubstituted derivatives of pentacarbonyliron such as bis(triphenylphosphine)tricarbonyliron react with nitrosyl chloride<sup>3</sup> or nitrosonium hexafluorophosphate<sup>4</sup> to give salts of a nitrosyl-cation (Ia) and (Ib). Since the nitrosonium and diazonium cations are isoelectronic we felt that arylazo complexes of iron might be accessible *via* the reactions of  $L_2Fe(CO)_3$  with diazonium tetrafluoroborates. This reaction does proceed smoothly and the major products are orange crystalline solids for which microanalytical data suggest the empirical formulae (IIa—f).

In the case of complex (IIe) where L is 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane, (etpb), we had difficulty in obtaining the pure precursor  $L_2Fe(CO)_3$  by


<sup>3</sup> G. R. Crooks and B. F. G. Johnson, *J. Chem. Soc. (A)*, 1968, 1238.

<sup>4</sup> B. F. G. Johnson and J. A. Segal, *J.C.S. Dalton*, 1972, 1268.

standard procedures.<sup>5</sup> Crooks and Johnson<sup>3</sup> had found that the nitrosyl complex (Ib) could also be prepared in reasonable yield from  $(\text{Ph}_3\text{P})\text{Fe}(\text{CO})_4$  and we successfully synthesised complex (IIe) starting with  $(\text{etpb})\text{Fe}(\text{CO})_4$  or a crude mixture of the mono- and bis-etpb iron carbonyls. Similarly the only product isolated from the reaction of  $(\text{Ph}_3\text{P})\text{Fe}(\text{CO})_4$  and benzenediazonium salts was the bis-phosphine substituted arylazo complex (IIa). However the related tetracarbonyl  $(o\text{-CH}_3\text{C}_6\text{H}_4)_3\text{PFe}(\text{CO})_4$  failed to react. We have shown elsewhere<sup>6</sup> that this latter complex has a sterically crowded iron atom and it seems likely that steric reasons are responsible for the failure to react with diazonium ion. Trofimenko<sup>7</sup> has shown that the nitrosonium ion is less sensitive to steric effects than the diazonium cation, and in agreement with this  $(o\text{-CH}_3\text{C}_6\text{H}_4)_3\text{PFe}(\text{CO})_4$  does react<sup>8</sup> with  $\text{NOPF}_6$  to yield  $\{(o\text{-CH}_3\text{C}_6\text{H}_4)_3\text{PFe}(\text{CO})_3\text{NO}\}^+\text{PF}_6^-$ .

All the new compounds are stable, both in solution in polar solvents and as solids. Complex (IIa) has been exposed to normal laboratory atmospheres for periods in excess of one year without appreciable deterioration. The analogous nitrosyl-cations decompose after *ca.* three weeks.<sup>3</sup> Conductance values in acetone solution are in the region expected<sup>9</sup> for 1 : 1 electrolytes and the complexes are diamagnetic (n.m.r. method<sup>10</sup>).

A yellow air-stable cationic nitrogen-free iron complex (III) inevitably accompanies the formation of complexes (IIa–f) in these reactions. Interestingly a similar (and probably isostructural) nitrogen-free complex (IV) is the sole product from the reaction of diazonium salts and  $\text{Ru}(\text{CO})_3(\text{PPh}_3)_2$ . Johnson and Segal<sup>4</sup> have reported that whereas  $\text{L}_2\text{Fe}(\text{CO})_3$  reacts with  $\text{NOPF}_6$  to give the nitrosyl complex (Ib),  $\text{L}_2\text{Ru}(\text{CO})_3$  gives instead  $\{\text{L}_2\text{Ru}(\text{CO})_3\text{H}\}^+$ . On the basis of their stability in air and the absence of any n.m.r. evidence for a hydridic hydrogen it appears that neither complex (III) or (IV) is a hydrido-cation. These compounds are receiving further study.



Following the work of Ibers and others<sup>11</sup> on nitrosyl complexes, two limiting structures (i) and (iii) may be considered for the co-ordinated arylazo ligand. Structure (i) involves a formal  $\text{ArNN}^+$  ligand (analogous to  $\text{NO}^+$ ) behaving as  $\sigma$ -donor to the metal, with some degree

of reverse  $\pi$ -bonding from metal to ligand  $\pi^*$  orbitals. Both N atoms of the azo-structure are  $sp$  hybridised and the  $\text{ArNN}$  ligand is linear. Structure (iii) is related to (ii) by a formal transfer of two electrons from metal to ligand so that the ligand now behaves as  $\text{ArNN}^-$  (analogous to  $\text{NO}^-$ ). The ligand-metal interaction is purely  $\sigma$  in nature, both N atoms are  $sp^2$  hybridised and the metal-arylazo structure has a bent azobenzene-like geometry. A third possibility (ii) may be recognised for the  $\text{ArN}_2$  ligand and, unlike structures (i) and (iii), structure (ii) has no geometric equivalent in nitrosyl chemistry. Here  $\text{M} \rightarrow \text{L}$  electron transfer may be considered to have taken place to such an extent as to involve a full formal M–L double bond, but falling short of the complete transfer envisaged for (iii). The NN linkage is now a double bond with  $sp$  hybridisation at the metal-bonded nitrogen atom and  $sp^2$  hybridisation at the nitrogen bonded to the aryl group. This implies linear MNN and angular  $\text{ArNN}$  structures. Of course structures (i)–(iii) are limiting only and a broad continuum of geometries between (i) and (iii) may be expected with deviation from linearity becoming more pronounced as  $\text{M} \rightarrow \text{L}$  electron-transfer increases.

Preliminary structural data<sup>12</sup> for complex (IIa) indicates that the overall geometry is trigonal-bipyramidal with apical phosphines, the equatorial positions being occupied by the carbonyl groups and the arylazo ligand. A similar geometry has recently been confirmed<sup>13</sup> for the related complex  $\{(\text{NO})\text{Os}(\text{CO})_2(\text{Ph}_3\text{P})_2\}^+\text{ClO}_4^-$  and has been inferred on the basis of i.r. spectroscopic data for the iron analogue (Ib). While the data for complex (IIa) is in an early state of refinement it is clear that the  $\text{FeNN}$  unit is linear with an N–N bond length of *ca.* 1.19 Å. The  $\text{PhNN}$  unit on the other hand is distinctly bent with a C–N–N angle *ca.* 124°. The structure thus approximates to bonding condition (ii) above.

The new complexes have been characterised by their  $^1\text{H}$ , n.m.r., i.r., and u.v.–visible spectra. In addition the  $^{19}\text{F}$  and  $^{31}\text{P}$  n.m.r. spectra and  $^{57}\text{Fe}$  Mössbauer spectra have been recorded for selected compounds. These results are summarised in the Table.

The  $^1\text{H}$  n.m.r. spectra for complexes (IIa–d) and (IIf) are uninformative as the proton resonances of the phosphine or arsine ligands overlap with those of the aryl ring of the ligand. In complex (IIe) the protons of the cyclic phosphite ligand and the arylazo ligand may be distinguished. As expected the  $-\text{OCH}_2-$  protons of the phosphite show the triplet pattern characteristic of virtual coupling.<sup>14</sup> The value of  $J(\text{POCH}_2)$  is almost unchanged from that observed<sup>5</sup> for the related complex  $\text{Fe}(\text{CO})_3\{\text{P}(\text{OCH}_2)_3\text{CCH}_3\}_2$ . The  $^{31}\text{P}$  spectrum of complex (IIa) displays a single resonance as expected for a structure with two equivalent phosphorus atoms.

<sup>5</sup> J. G. Verkade, R. E. McCarley, D. G. Hendricker, and R. W. King, *Inorg. Chem.*, 1965, **4**, 228.

<sup>6</sup> W. E. Carroll, F. A. Deeney, J. A. Delaney, and F. J. Lalor, *J.C.S. Dalton*, 1973, 718.

<sup>7</sup> S. Trofimenko, *Inorg. Chem.*, 1971, **10**, 504.

<sup>8</sup> W. E. Carroll and F. J. Lalor, *J. Organometallic Chem.*, submitted for publication.

<sup>9</sup> R. B. King, *Organometallic Synth.*, 1965, **1**, 52.

<sup>10</sup> D. F. Evans, *J. Chem. Soc.*, 1959, 2003.

<sup>11</sup> D. M. P. Mingos and J. A. Ibers, *Inorg. Chem.*, 1971, **10**, 1479 and references therein.

<sup>12</sup> J. A. Ibers, personal communication.

<sup>13</sup> G. R. Clark, K. R. Grundy, W. R. Roper, J. M. Waters, and K. R. Whittle, *J.C.S. Chem. Comm.*, 1972, 119.

<sup>14</sup> W. E. Stancil and D. G. Hendricker, *Inorg. Chem.*, 1968, **7**, 1242 and references therein.

Complexes (IIa—f) owe their orange colour to strong end-absorption from the u.v. region. No maximum was observed in the region 412—600 nm assigned by previous workers to the  $n \rightarrow \pi^*$  transition of the coordinated azo group.<sup>1,15-19</sup> At higher energies a weak inflection occurs in the region 349—398 nm which we tentatively assign to  $\lambda_{n-\pi^*}$ . Following the conclusions of previous workers<sup>18</sup> a shift to high energy of  $\lambda_{n-\pi^*}$  implies that delocalisation of electron density from the iron atom along the arylazo chain is not extensive.

Some degree of confirmation for this suggestion is provided by the i.r. spectra of complexes (IIa—f) (see Table) and the <sup>19</sup>F n.m.r. spectra of complexes (IIb and c). Of the three strong bands in the 2100—1700 cm<sup>-1</sup> region the two of highest wavenumber are assigned to  $\nu(\text{CO})$ . The third band is confirmed by an isotopic labelling experiment to be due to  $\nu(\text{NN})$ . Assignment of  $\nu(\text{NN})$  in arylazo complexes to absorptions in the range 1440—1620 cm<sup>-1</sup> (neutral compounds)<sup>1,15-17,19-22</sup> and 1622—1700 cm<sup>-1</sup> (cationic complexes)<sup>18,23,24</sup> has been made by previous workers. Confirmation of assignment by isotopic substitution is only available<sup>17</sup> for the narrower range of 1440—1505 cm<sup>-1</sup>. Our work considerably extends the verified range of  $\nu(\text{NN})$ . The value of  $\nu(\text{NN})$  in compounds (IIa—f) is the highest yet recorded and suggests, with the u.v. evidence, a lower degree of metal-arylazo back bonding than previously observed for complexes of this type. The two  $\nu(\text{CO})$  absorptions in compounds (IIa—f) lie at lower wavenumber than those in the analogous nitrosyl (I)<sup>4</sup> suggesting that the arylazo group is moreover a weaker  $\pi$ -acceptor than is NO.

Because the *p*-fluorine atom of complex (IIb) is deshielded relative to the *m*-fluorine atom of complex (IIc) by —4.8 to —4.9 p.p.m. (internal and external C<sub>6</sub>H<sub>5</sub>F standard respectively), there is significant conjugative withdrawal of electrons from the aromatic ring by the attached  $-\{\text{N}_2\text{Fe}^+(\text{CO})_2\text{L}_2\}$  structure, hence transfer of electron density from metal to ligand by back-bonding cannot be extensive.

In view of the probable relationship between  $\text{M} \rightarrow \text{L}$  electron transfer and the geometry of the ArNN ligand it is interesting to compare the foregoing results with those reported<sup>19</sup> for  $[\text{Mo}(\text{CO})_2\{\text{HB}(\text{C}_3\text{H}_3\text{N}_2)\}(\text{PhNN})]$  in which the geometry of the ArNN ligand is structurally similar<sup>25</sup> to that in complexes (IIa—f). For complexes of this type the <sup>19</sup>F n.m.r. results\* indicate a very much smaller deshielding of the *p*-F atom (*ca.* —0.04 p.p.m.)

\* Data from ref. 19 measured with respect to external C<sub>6</sub>H<sub>5</sub>F standard. For this complex <sup>19</sup>F n.m.r. spectra are complicated by specific solute-standard interactions when the standard is internal. Our measurements on (IIb—c) were not similarly affected.

<sup>15</sup> G. W. Parshall, *J. Amer. Chem. Soc.*, 1965, **87**, 2133.

<sup>16</sup> C. D. Cook and G. S. Jauhal, *J. Amer. Chem. Soc.*, 1968, **90**, 1464.

<sup>17</sup> (a) G. W. Parshall, *J. Amer. Chem. Soc.*, 1967, **89**, 1822; (b) A. B. Gilchrist, G. W. Rayner-Canham, and D. Sutton, *Nature*, 1972, **235**, 42.

<sup>18</sup> S. Cenini, R. Ugo, and G. LaMonica, *J. Chem. Soc. (A)*, 1971, 3441.

and therefore significantly more back-bonding from the metal to the arylazo ligand. The close similarity of structures is unexpected. It appears that the geometry of the ArNN ligand is more sensitive to the effects of population of the ligand  $\pi^*$  orbitals than might have been predicted; even the small degree of back-bonding evident in (IIa—f) being sufficient to significantly distort the structure from linearity.

Finally we have determined the <sup>57</sup>Fe Mössbauer spectra of arylazo complexes (IIa and d) and the nitrosyl analogue (Ib). Replacing the NO ligand by the ArNN structure produces a marked increase in the isomer shift ( $\delta$ ) of the iron nucleus and decreases the quadrupole splitting ( $|\Delta E_q|$ ). Thus the effective *s*-electron density at the <sup>57</sup>Fe nucleus is greater in the nitrosyl than in the arylazo complex. This information is certainly consistent with our tentative conclusion from i.r. data that ArN<sub>2</sub> is the poorer  $\pi$ -acceptor of the two. However a similar effect would be produced if ArN<sub>2</sub> were a weaker  $\sigma$ -donor than NO. More information will be necessary to separate the two effects. Complex (Ia) may be described as formally  $d^8\text{-Fe}^0$  or  $d^7\text{-Fe}^I$ . In either case the non-spherically symmetric *d* shell renders it difficult to interpret changes in  $|\Delta E_q|$  values in terms of changes in metal to ligand  $\pi$ -bonding. While there are some grounds for suggesting that in complexes of this type increasing  $\pi$ -bonding produces a decrease in  $|\Delta E_q|$ , the evidence is not conclusive.<sup>6</sup> Here too more work is needed. It may be pointed out, however, that complex (IIa) has the smallest value of  $|\Delta E_q|$  yet recorded for complexes of unambiguously trigonal-bipyramidal geometry. In future greater care will be necessary before the geometry of a co-ordinate sphere can be assigned simply on the basis of  $|\Delta E_q|$  value. We are presently carrying out an extensive study of the transmission of electronic effects along the PhNNFe chain using Mössbauer and other spectroscopic techniques as well as investigating the chemistry of these new and very stable iron complexes.

#### EXPERIMENTAL

Substituted anilines were available commercially and were converted to diazonium tetrafluoroborates or hexafluorophosphates by standard procedures.<sup>26</sup> Sodium [<sup>15</sup>N]-nitrite (99%) was purchased from the Isomet Corporation, 103 Bauer Drive, Oakland, New Jersey, and was used to synthesise <sup>15</sup>N-labelled benzenediazonium tetrafluoroborate. Triphenylphosphine, triphenylarsine, and 4-ethyl-2,6,7-trioxa-1-phosphabicyclo[2.2.2]octane were commercially available and were used as received. The L<sub>2</sub>Fe(CO)<sub>3</sub>

<sup>19</sup> S. Trofimenko, *Inorg. Chem.*, 1969, **8**, 2675.

<sup>20</sup> R. B. King and M. B. Bisnette, *Inorg. Chem.*, 1966, **5**, 300.

<sup>21</sup> M. C. Baird and G. Wilkinson, *J. Chem. Soc. (A)*, 1967, 865.

<sup>22</sup> F. W. B. Einstein, A. B. Gilchrist, G. W. Rayner-Canham, and D. Sutton, *J. Amer. Chem. Soc.*, 1972, **94**, 645.

<sup>23</sup> D. Cashman and F. J. Lalor, *J. Organometallic Chem.*, 1970, **24**, C29; 1971, **32**, 351.

<sup>24</sup> J. A. McCleverty and R. N. Whiteley, *Chem. Comm.*, 1971, 1159.

<sup>25</sup> G. Avitabile, P. Ganis, and M. Nemiroff, *Acta Cryst.*, 1971, **B27**, 725.

<sup>26</sup> A. Roe, *Org. Reactions*, 1949, **5**, 193.

compounds were synthesised by literature methods.<sup>5,27</sup> Microanalytical and molecular weight data were carried out in the Analytical Laboratory of this department and by the Pascher Microanalytical Laboratory, Buchstrasse 54, Bonn, W. Germany.

I.r. spectra were recorded on a Perkin-Elmer 257 grating spectrophotometer. Electronic spectra were measured using a Perkin-Elmer 402 u.v.-visible instrument. <sup>1</sup>H and <sup>19</sup>F n.m.r. spectra were obtained on a Perkin-Elmer-Hitachi R 20A spectrometer at 60.0 and 56.456 MHz respectively. Field-sweep mode was used in both cases. The <sup>31</sup>P spectrum of complex (IIa) was measured on a Bruker HFX spectrometer with complete proton decoupling. We are indebted to Dr. W. R. Jackson of Queen's University, Belfast, and Dr. B. E. Mann of the University of Leeds for this data. Conductivity measurements were made with a Radiometer CDM2e direct-reading conductivity meter using a CDC104

product. Orange micro-crystals of analytical purity were obtained by recrystallisation from acetone-ether or methylene chloride-ether. The yield of pure material averaged 50% of theory. Further work-up of the original reaction mixture yielded a second cationic product which is currently being investigated (see text). Microanalytical, m.p., and i.r. spectroscopic data for the new complexes are collected in the Table.

**Nuclear Magnetic Resonance Data.**—<sup>1</sup>H Spectrum of complex (IIe) determined in [<sup>2</sup>H<sub>6</sub>]acetone (internal Me<sub>4</sub>Si):  $\tau$  ca. 2.4(m) (aromatic protons), 5.35(t) (—OCH<sub>2</sub>—protons), ca. 8.65(m) (exocyclic methylene group), and 9.15(m) (methyl group).  $J(\text{POCH}_2)$  (taken as the separation of the outer parts of the endocyclic methylene triplet) was 5 Hz.

<sup>31</sup>P Spectra. After full proton decoupling, complex (IIa) (in solution in methylene chloride) showed a single <sup>31</sup>P resonance at —59.6 p.p.m. relative to 85% H<sub>3</sub>PO<sub>4</sub>.

#### Analytical, spectroscopic, and other data

Complex	R	L	X	Analytical data (%) <sup>a</sup>				Infrared <sup>b</sup>			Electronic spectra <sup>c</sup>			Mössbauer spectra <sup>e</sup>		
				C	H	N	M.p./°C	$\nu(\text{CO})$	$\nu(\text{NN})$		$\Delta\epsilon$ <sup>d</sup>	$\delta/\text{mm s}^{-1}$	$ AEq /\text{mm s}^{-1}$			
(IIa)	H	PPh <sub>3</sub>	BF <sub>4</sub>	64.0(63.8)	4.6(4.3)	3.5(3.4)	204—205	2030	1778	1723 <sup>g</sup>	349(4.23)	293(45.0)	232(61.5)	143 ± 3	0.092 ± 0.01	0.97 ± 0.01
(IIb)	<i>p</i> -F	PPh <sub>3</sub>	BF <sub>4</sub>	62.6(62.4)	4.2(4.0)	3.6(3.3)	210—211	2027	1776	1720	346(5.55)	296(43.1)	232(45.0)	140 ± 2		
(IIc)	<i>m</i> -F	PPh <sub>3</sub>	BF <sub>4</sub>	62.8(62.4)	4.2(4.0)	3.4(3.3)	187	2030	1981	1714	349(3.75)	295(44.6)	231(51.2)			
(IId)	<i>p</i> -NO <sub>2</sub>	PPh <sub>3</sub>	BF <sub>4</sub>	60.4(60.5)	4.0(3.9)	4.7(4.8)	166—168	2036	1984	1703	398(2.74)	308(37.4)	231(49.8)	132 ± 2	0.092 ± 0.01	1.08 ± 0.01
(IIe)	<i>p</i> -F	etpb	PF <sub>6</sub>	33.9(34.1)	3.7(3.7)	4.2(4.0)	170	2070	2014	1763	350(1.60)	276(33.0)	212(22.5)			
(II <sub>f</sub> )	H	AsPh <sub>3</sub>	BF <sub>4</sub>	57.7(57.6)	3.9(3.8)	3.3(3.0)	173—174	2028	1978	1720	370(3.15)	299(47.3)	217(59.0)			
(Ib)								2051 <sup>h</sup>	1995 <sup>h</sup>	1792 <sup>h,i</sup>				0.075 ± 0.01	2.05 ± 0.01	

<sup>a</sup> Calculated values in parentheses. <sup>b</sup> In cm<sup>-1</sup>, CH<sub>2</sub>CH<sub>2</sub> solution, 2100—1700 cm<sup>-1</sup> region. <sup>c</sup> Methanol solution, in nm, extinction coefficients  $\times 10^{-3}$  in parentheses. <sup>d</sup> Acetone solution, concentration  $1.0 \times 10^{-3}$ — $2.0 \times 10^{-4}$  mol l<sup>-1</sup>, in  $\Omega^{-1}$  cm<sup>2</sup> mol<sup>-1</sup>. <sup>e</sup> Measured with respect to natural iron. <sup>f</sup> Iron analysis, 6.8(6.7)%. <sup>g</sup>  $\nu(^{14}\text{N}^{15}\text{N})$  occurs at 1690 cm<sup>-1</sup>. <sup>h</sup> Data from ref. 4. <sup>i</sup>  $\nu(\text{NO})$ .

electrode. Mössbauer spectra were run at room temperature using a commercial drive unit and transducer, TMC model 305/306 with a TMC Gammascope II multichannel analyser. The source was <sup>57</sup>Co in Pd and was of nominal strength 5 mCi. Absorbers were made up in Al foil and were ca. 50 mg cm<sup>-1</sup> in thickness. A non-linear curve-fitting programme was used and the data processed on an 1130 IBM computer. We are grateful to Dr. F. A. Deeney of the Experimental Physics Department of this College for the Mössbauer data.

**Preparation of the Complexes** [ArN<sub>2</sub>Fe(CO)<sub>2</sub>L<sub>2</sub>]<sup>+</sup>X<sup>-</sup> (IIa—f).—A stirred solution or suspension of L<sub>2</sub>Fe(CO)<sub>3</sub> [or LFe(CO)<sub>4</sub> where L is etpb] in acetone or methylene chloride was treated under nitrogen with a slight excess of solid diazonium salt. Vigorous effervescence occurred and the yellow solution developed a transient green tinge, changing rapidly to deep red. After 30 min stirring at room temperature [40 °C in the case of (etpb)Fe(CO)<sub>4</sub>] the solution was filtered. Addition of ether precipitated the crude

<sup>19</sup>F Chemical shifts were determined in methylene chloride solution (fluorobenzene standard). A shift to higher frequency is taken as being positive in sign. Complex (IIb); *p*-fluorine; —8.34 p.p.m. (internal standard) —8.22 p.p.m. (external standard). BF<sub>4</sub><sup>-</sup> ion; +38.72 p.p.m. (internal standard) and +39.02 p.p.m. (external standard). Complex (IIc): *m*-fluorine; —3.51 p.p.m. (internal standard) and —3.28 p.p.m. (external standard). BF<sub>4</sub><sup>-</sup> ion; +39.16 p.p.m. (internal standard) and +39.16 p.p.m. (external standard).  $\delta_{p-F} - \delta_{m-F} = -4.83$  p.p.m. (internal standard) and —4.94 p.p.m. (external standard).

This work was carried out during the tenure by one of us (W. E. C.) of a Department of Education maintenance grant for postgraduate research.

[2/1903 Received, 10th August, 1972]

<sup>27</sup> A. F. Clifford and A. K. Mukherjee, *Inorg. Chem.*, 1963, **2**, 151.