

## Germanium Chemistry

**Germacarboxylic Acid: An Organic-Acid Analogue Based on a Heavier Group 14 Element\*\****Leslie W. Pineda, Vojtech Jancik, Herbert W. Roesky,\* and Regine Herbst-Irmer**Dedicated to Professor Jean-Marie Lehn on the occasion of his 65th birthday*


Carbon multiple-bonded species, especially those containing carbonyl groups (aldehydes, amide esters, ketones) are widely known and useful systems in organic chemistry.<sup>[1]</sup> As far as the heavier congeners of Group 14 are concerned, a steady and remarkable development has been experienced, which includes the synthesis of unsaturated species. Additionally a myriad of mixed unsaturated compounds has been prepared containing elements of Groups 14–16.<sup>[2]</sup> However, owing to the high reactivity and tendency to polymerize these species have to be thermodynamically and kinetically stabilized.

In the case of germanium–chalcogen double-bonded species, a few thio-, seleno-, and telluroketones were prepared.<sup>[3]</sup> We have already reported the synthesis and structure of  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{Ge}(\text{S})\text{X}]^{[4]}$  ( $\text{X} = \text{Cl}, \text{F}$ ,  $\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2 = 2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{NC}(\text{CH}_3)\text{CH}-\text{C}(\text{CH}_3)\text{NC}_6\text{H}_3\text{-}2,6\text{-}i\text{Pr}_2$ ) with Group 14 and 16 elements

[\*] Dipl.-Chem. L. W. Pineda, Dipl.-Chem. V. Jancik, Prof. Dr. H. W. Roesky, Dr. R. Herbst-Irmer  
Institut für Anorganische Chemie  
Universität Göttingen  
Tammannstrasse 4, 37077 Göttingen (Germany)  
Fax: (+49) 551-39-3373  
E-mail: hroesky@gwdg.de


[\*\*] This work was supported by the Deutsche Forschungsgemeinschaft, the Göttinger Akademie der Wissenschaften, and the Fonds der Chemischen Industrie. L.W.P. thanks the Deutscher Akademischer Austauschdienst (DAAD) for a predoctoral fellowship.

bearing a halide. Furthermore, the synthesis and structures of the selenium analogue  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{Ge}(\text{Se})\text{X}]$  ( $\text{X} = \text{Cl}, \text{F}$ ), as well as the functionalized derivative  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{Ge}(\text{E})\text{R}]$  ( $\text{E} = \text{S}, \text{Se}$ ;  $\text{R} = \text{Me}$  or  $n\text{Bu}$ ) were reported.<sup>[5]</sup> Recently, we succeeded in the isolation and structural characterization of the first terminal hydroxide based on germanium(II)  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{GeOH}]$  (**1**).<sup>[6]</sup> As a stable precursor compound **1** is quite intriguing for the generation of new functional groups. To our knowledge the co-existence and fast tautomeric equilibrium for thiolcarboxylic acid **2** and thionocarboxylic acid **3** is known, but the latter group does not exist in the free state (Scheme 1).<sup>[7]</sup> Herein we report the first successful isolation and full characterization of a germanium thionoacid  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{Ge}(\text{S})\text{OH}]$  (**4**), which has no isolated precedent in the carbon system (see Scheme 2).



**Scheme 1.** Tautomeric equilibrium for the thiol- and thionocarboxylic acid.

The reaction of **1** in the presence of equivalent amounts of elemental sulfur at room temperature in toluene leads after three days to the white compound **4** in moderate yield. (Scheme 2).




**Scheme 2.** Ar = 2,6-*i*Pr<sub>2</sub>C<sub>6</sub>H<sub>3</sub>.

Compound **4** is soluble in benzene, THF, and hexane, but insoluble in pentane and shows no decomposition on exposure to air. Compound **4** was fully characterized by IR, <sup>1</sup>H, and <sup>13</sup>C NMR spectroscopy, EI mass spectrometry, elemental analysis, and X-ray structural analysis. By comparison of the IR spectrum of **1**, which exhibits a sharp OH stretching vibration at 3571 cm<sup>-1</sup><sup>[6]</sup> with the corresponding frequency of **4** (3238 cm<sup>-1</sup>) a significant shift to lower wave numbers is observed. Such behavior may be due to the formation of hydrogen bonds. Interestingly, the <sup>1</sup>H NMR spectrum of **4** exhibits a resonance signal at  $\delta = 2.30$  ppm for the hydroxy hydrogen, which by comparison with that of **1** ( $\delta = 1.54$  ppm) clearly shows a downfield shift. Again intermolecular hydrogen interaction and a change of the oxidation state of the germanium atom are plausible explanations for the observed

shift. Thus, this resonance shift indicates a fairly acidic nature of the terminal OH proton. Furthermore no evidence was found for any tautomeric equilibrium of **4**. The most abundant ion peak in the EI mass spectrum appeared at  $m/z$  525  $[\text{M}-\text{Me}]^+$ , and the signal at  $m/z$  540 (40%) was assigned to the molecular ion  $[\text{M}]^+$  (correct isotopic pattern).

Maintaining a toluene solution of **4** for two weeks at  $-20^\circ\text{C}$ , resulted in colorless single crystals suitable for X-ray structural analysis.<sup>[8]</sup> Compound **4** crystallizes in the monoclinic space group  $C2/c$ , with one monomer and one molecule of toluene in the asymmetric unit. Intermolecular interaction of the hydroxy group with the sulfur atom results in the formation of the hydrogen-bonding array (O–H $\cdots$ S) leading to dimers (Figure 1). The hydrogen bonded donor–acceptor



**Figure 1.** Thermal ellipsoid plot of **4** (thermal ellipsoids set at 50% probability). H atoms, except for the OH group, and interstitial toluene molecules, are omitted for clarity. Selected bond lengths [Å] and angles [°]: Ge(1)–O(1) 1.751(2), Ge(1)–N(2) 1.911(2), Ge(1)–N(1) 1.916(2), Ge(1)–S(1) 2.077(1); O(1)–Ge(1)–N(2) 99.6(1), O(1)–Ge(1)–N(1) 102.2(1), N(2)–Ge(1)–N(1) 95.6(1), O(1)–Ge(1)–S(1) 121.4(1), N(2)–Ge(1)–S(1) 118.8(1), N(1)–Ge(1)–S(1) 114.9(1).

separations (H $\cdots$ S, 2.537 Å and O $\cdots$ S 3.234 Å) follow the same trend as those reported in literature.<sup>[9]</sup> The coordination environment around the germanium atom comprises two nitrogen atoms from the supporting ligand, one hydroxy group, and one sulfur atom, and has a distorted tetrahedral geometry.

The Ge–O bond length (1.751(2) Å) in **4** is significantly shorter than that in **1** (1.828(1) Å), as a result of the smaller atomic radius of Ge<sup>IV</sup> compared with that of Ge<sup>II</sup>. Indeed, similar Ge–O bond lengths for Ge<sup>IV</sup> species have been described, (*t*Bu<sub>2</sub>Ge(OH)<sub>2</sub> (1.781(4) and 1.779(2) Å)<sup>[10]</sup> and 1.779(5) Å in  $[(\text{FcN})_3\text{GeOH}]$  (Fc = CpFe( $\eta^5$ -C<sub>5</sub>H<sub>4</sub>)).<sup>[11]</sup> A shorter Ge–N bond length and wider N–Ge–N angle are expected (av. 1.914(2) Å and 95.6(1)°) than in **1**. A comparison of the Ge–S bond length in  $[\{\eta^3\text{-}[(\mu\text{-}t\text{BuN})_2(\text{SiMe}_2\text{Bu})_2]\text{GeS}\}]^{[12]}$  (2.063(3) Å), and in  $[\text{HC}\{(\text{CMe})(2,6\text{-}i\text{Pr}_2\text{C}_6\text{H}_3\text{N})\}_2\text{Ge}(\text{S})\text{X}]^{[4]}$  ( $\text{X} = \text{Cl}, \text{F}$ ), (2.053(6) and 2.050(9) Å), with that in **4** (2.077(1) Å) shows a good agreement. Likewise, the Ge–O bond length in  $[(\text{dppe})\text{Pd}(\mu\text{-S})(\mu\text{-})$

$\text{CH}_2\text{O}\text{Ge}\{\text{N}(\text{SiMe}_3)_2\}_2$  (dppe = bis(diphenylphosphanyl)-ethene)<sup>[13]</sup> (1.785(6) Å), a compound which has almost the same coordination environment and geometry at the germanium center as **4**, correlates well with that in **4**.

In summary the reaction of **1** and elemental sulfur resulted in the formation of the title compound **4** which represents a new class of “carbon-free” carbonic acid analogues based on germanium. The stability of **4** against oxygen and water at room temperature makes these systems quite interesting for biological investigations.

## Experimental Section

All manipulations were performed under a dry and oxygen-free atmosphere ( $\text{N}_2$  or Ar) by using Schlenk-line and glove-box techniques. Solvents were purified prior to use by distillation over appropriate drying agents in a nitrogen atmosphere.

**4**: A solution of **1** (1.56 g, 3.07 mmol) in toluene (30 mL) was slowly added to a suspension of elemental sulfur (0.09 g, 3.07 mmol) in toluene (15 mL) by cannula at room temperature. After 3 days under constant stirring at ambient temperature the yellow solution turned slightly green. After removal of all volatiles the remaining crude product was rinsed with pentane ( $3 \times 10$  mL) and dried under reduced pressure to yield pure **4**. Yield: 1.10 g (66%); m.p. 300 °C (decomp); IR (KBr):  $\tilde{\nu} = 3238, 3063, 2965, 2867, 1638, 1539, 1442, 1388, 1322, 1257, 1175, 1102, 1023, 933, 875, 797, 712, 501$   $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{C}_6\text{D}_6$ , 25 °C, TMS):  $\delta = 7.09\text{--}7.16$  (m, 6H, *m*-, *p*-Ar-H), 4.83 (s, 1H,  $\gamma$ -CH), 3.62 (sept,  $^3J(\text{H,H}) = 6.8$  Hz, 2H,  $\text{CH}(\text{CH}_3)_2$ ), 3.35 (sept,  $^3J(\text{H,H}) = 6.8$  Hz, 2H,  $\text{CH}(\text{CH}_3)_2$ ), 2.30 (s, 1H, OH), 1.57 (d,  $^3J(\text{H,H}) = 6.8$  Hz, 6H,  $\text{CH}(\text{CH}_3)_2$ ), 1.47 (s, 6H,  $\text{CH}_3$ ), 1.26 (d,  $^3J(\text{H,H}) = 6.8$  Hz, 6H,  $\text{CH}(\text{CH}_3)_2$ ), 1.16 (d,  $^3J(\text{H,H}) = 6.8$  Hz, 6H,  $\text{CH}(\text{CH}_3)_2$ ), 1.05 ppm (d,  $^3J(\text{H,H}) = 6.8$  Hz, 6H,  $\text{CH}(\text{CH}_3)_2$ );  $^{13}\text{C}$  NMR (125.8 MHz,  $\text{C}_6\text{D}_6$ , 25 °C, TMS):  $\delta = 169.9$  (C=N), 145.9, 144.9, 137.2, 128.9, 124.8, 124.6 (*i*-, *o*-, *m*-, *p*-, Ar), 98.5 ( $\gamma$ -CH), 29.5 ( $\text{CH}_3$ ), 27.9 ( $\text{CH}(\text{CH}_3)_2$ ), 26.3 ( $\text{CH}(\text{CH}_3)_2$ ), 24.7 ( $\text{CH}(\text{CH}_3)_2$ ), 24.6 ( $\text{CH}(\text{CH}_3)_2$ ), 23.8 ( $\text{CH}(\text{CH}_3)_2$ ), 23.7 ppm ( $\text{CH}(\text{CH}_3)_2$ ); EI-MS (70 eV):  $m/z$  (%): 540 (40)  $[\text{M}]^+$ , 525 (100)  $[\text{M}-\text{CH}_3]^+$ ; elemental analysis (%) calcd for  $\text{C}_{29}\text{H}_{42}\text{GeN}_2\text{OS}$  (539.32): C 64.59, H 7.85, N 5.19; found: C 64.20, H 7.57, N 5.12.

Received: June 4, 2004

**Keywords:** acids · germanium · hydrogen bonds · oxidative addition · sulfur

- [1] J. McMurry, *Organic Chemistry*, Brooks-Cole, CA, **1992**, pp. 695–705.  
 [2] P. P. Power, *Chem. Rev.* **1999**, *99*, 3463–3504.  
 [3] a) N. Tokitoh, R. Okazaki, *Adv. Organomet. Chem.* **2001**, *47*, 121–166; b) N. Tokitoh, T. Matsumoto, R. Okazaki, *Bull. Chem. Soc. Jpn.* **1999**, *72*, 1665–1684; c) T. Matsumoto, N. Tokitoh, R. Okazaki, *J. Am. Chem. Soc.* **1999**, *121*, 8811–8824.

- [4] Y. Ding, Q. Ma, H. W. Roesky, R. Herbst-Irmer, I. Usón, M. Noltemeyer, H.-G. Schmidt, *Organometallics* **2002**, *21*, 5216–5220.  
 [5] Y. Ding, Q. Ma, H. W. Roesky, I. Usón, M. Noltemeyer, H.-G. Schmidt, *J. Chem. Soc. Dalton Trans.* **2003**, 1094–1098.  
 [6] L. W. Pineda, V. Jancik, H. W. Roesky, D. Neculai, A. M. Neculai, *Angew. Chem.* **2004**, *116*, 1443–1445; *Angew. Chem. Int. Ed.* **2004**, *43*, 1419–1421.  
 [7] a) F. Duus in *Comprehensive Organic Chemistry, Vol. 3* (Eds.: D. Barton, W. D. Ollis), Pergamon, Oxford, **1979**, pp. 420–421; b) J. Hine, *Physical Organic Chemistry*, McGraw-Hill, **1962**, p. 238.  
 [8] Crystal data for **4**-toluene:  $\text{C}_{36}\text{H}_{50}\text{GeN}_2\text{OS}$ ,  $M_r = 631.43$ , monoclinic, space group  $C2/c$ ,  $a = 26.021(1)$ ,  $b = 16.045(1)$ ,  $c = 18.006(2)$  Å,  $\beta = 114.79(1)^\circ$ ,  $V = 6825(1)$  Å<sup>3</sup>,  $Z = 8$ ,  $\rho_{\text{calcd}} = 1.229$   $\text{g cm}^{-3}$ ,  $F(000) = 2688$ ,  $\lambda = 1.54178$  Å,  $T = 100(2)$  K,  $\mu(\text{Cu}_{\text{K}\alpha}) = 2.002$   $\text{mm}^{-1}$ . Data for the structure were collected on a Bruker three-circle diffractometer equipped with a SMART 6000 CCD detector. Intensity measurements were performed on a rapidly cooled crystal ( $0.20 \times 0.10 \times 0.10$   $\text{mm}^3$ ) in the range  $6.66 \leq 2\theta \leq 118.08^\circ$ . Of the 21 505 measured reflections, 4748 were independent [ $R(\text{int}) = 0.0363$ ]. The structure was solved by direct methods (SHELXS-97)<sup>[14]</sup> and refined with all data by full-matrix least squares on  $F^2$ .<sup>[15]</sup> The hydrogen atoms of C–H bonds were placed in idealized positions, whereas the hydrogen atom from the OH moiety was localized from the difference electron-density map and refined isotropically. The final refinements converged at  $R1 = 0.0283$  for  $I > 2\sigma(I)$ ,  $wR2 = 0.0730$  for all data. The final difference Fourier synthesis gave a min/max residual electron density  $-0.279/+0.355$   $\text{e} \text{Å}^{-3}$ . CCDC-240065 (**4**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via [www.ccdc.cam.ac.uk/conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html) (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).  
 [9] a) N. N. Greenwood, A. Earnshaw, *Chemistry of the Elements*, Butterworth-Heinemann, Oxford, **1997**, p. 60; b) A. F. Wells, *Structural Inorganic Chemistry*, Clarendon, Oxford, **1984**, p. 357.  
 [10] H. Puff, S. Franken, W. Schuh, W. Schwab, *J. Organomet. Chem.* **1983**, *254*, 33–41.  
 [11] A. Fischer, K. Jacob, F. T. Edelmann, *Z. Anorg. Allg. Chem.* **2003**, *629*, 963–967.  
 [12] M. Veith, S. Becker, V. Huch, *Angew. Chem.* **1989**, *101*, 1287–1289; *Angew. Chem. Int. Ed. Engl.* **1989**, *28*, 1237–1238. For comparison, see also M. C. Kuchta, G. Parkin, *J. Chem. Soc. Chem. Commun.* **1994**, 1351–1352; M. Veith, A. Rammo, *Z. Anorg. Allg. Chem.* **1997**, *623*, 861–872; I. Saur, G. Rima, H. Gornitzka, K. Miqueu, J. Barrau, *Organometallics* **2003**, *22*, 1106–1109.  
 [13] Z. T. Cygan, J. W. Kampf, M. M. Banaszak Holl, *Organometallics* **2004**, *23*, 2370–2375.  
 [14] “SHELXS-97, Program for Structure Solution”: G. M. Sheldrick, *Acta Crystallogr. Sect. A* **1990**, *46*, 467–473.  
 [15] G. M. Sheldrick, SHELXL-97, Program for Crystal Structure Refinement, University of Göttingen, Göttingen (Germany), **1997**.