

Expanding Coefficient: A Parameter To Assess the Stability of Induced-Fit Complexes

Carmen Talotta,* Gerardo Concilio, Margherita De Rosa, Annunziata Soriente, Carmine Gaeta, Antonio Rescifina,* Pablo Ballester,* and Placido Neri*



Cite This: *Org. Lett.* 2021, 23, 1804–1808



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: Here we propose a new parameter, the Expanding Coefficient (EC), that can be correlated with the thermodynamic stability of supramolecular complexes governed by weak secondary interactions and obeying the induced-fit model. The EC values show a good linear relationship with the log K_{app} of the respective pseudorotaxane complexes investigated. According to Cram's Principle of Preorganization, the EC can be considered an approximate mechanical measure of the host's reorganization energy cost upon adopting the final bound geometry.



Molecular recognition is a fundamental phenomenon at the ground of every biological function.¹ Its fundamental relevance was recognized since the early days of chemical and biological sciences,² and afterward, many studies have been devoted to comprehending its underlying principles and explaining the origin of the amazing selectivity³ or affinity⁴ observed in some instances. Two main models are currently adopted to describe a molecular recognition complexation: (i) the lock-and-key model⁵ and (ii) the induced-fit complexation.⁶ In the first one, the degree of geometrical (steric) fitting between a rigid, undeformable receptor and substrate is evaluated, while, in the second one, the best match of flexible counterparts is considered after their mutual adaptation.

In the field of synthetic receptors, a single value parameter, the Packing Coefficient (PC), defined as the ratio between the van der Waals volume of the hosted guest and the volume of the host cavity,⁷ was introduced by Rebek and Mecozzi to simplify the assessment of the complex stability for rigid, lock-and-key-like systems. It was found that the optimal PC value, the fraction of occupied volume, is 0.55 ± 0.09 (55%)⁷ in the liquid state when weak intermolecular interactions (dispersion forces, van der Waals interactions) are present. The PC can increase (up to 70–84%) when stronger interactions (hydrogen bonds, solvation effects) come into play^{8a,b} or can decrease (down to 40%) when gaseous molecules are involved.^{8c,d} This simple rule has been validated in many synthetic host–guest systems,⁹ has been used to explain reactivity results,¹⁰ and has also been extended to biological receptors.¹¹

From the above definitions, it is evident that the PC parameter cannot be applied to induced-fit systems because the void receptor cavity volume tends to adapt to that of the guest to give the best fitting with it.¹² Therefore, it is expected that the PC value of induced-fit complexes calculated for the final adapted geometry will always overcome the 55% rule (vide

infra), independently from their actual stability, thus vanishing any comparison.


It is evident that another useful and straightforward rule is necessary to assess the stability of induced-fit complexes in order to predict the ideal host–guest couple. We propose here a new single-value parameter to address such point.

The induced-fit system chosen for this work is the pseudorotaxane complex formed by hexamethoxy-*p*-*tert*-butylcalix[6]arene¹³ **1** and alkylbenzylammonium axles **2a–k**⁺ (Scheme 1). In previous studies,¹⁴ we have demonstrated that this kind of dialkylammonium axles can thread the calixarene cavity when coupled to the weakly interacting BARF[–] (or TFPB[–]) “superweak anion” to give the **2**⁺⊂**1** pseudorotaxanes (Scheme 1). In addition, we have found that the aromatic cavity mostly prefers to host the alkyl portion of the axle with respect to the benzyl one (the so-called “endo-alkyl rule”).^{14a–e}


The aromatic walls of receptor **1** are freely rotating through the macrocyclic annulus, and thus they adapt their relative orientation to give the best interactional fitting with guest **2**. As a result, a less-symmetrical induced-fit complex is obtained in which each aromatic ring is differently inclined toward the cavity (*vide infra*). Since the PhCH₂NH₂⁺ moiety is common to all the **2a–k**⁺ axles and since it is external to the aromatic calix-cavity, we can assume that the stability constant of **2**⁺⊂**1** pseudorotaxanes could be directly linked to the best interac-

Received: January 20, 2021

Published: February 16, 2021



Scheme 1. Threading of Calix[6]arene **1** with Alkylbenzylammonium Axles **2a–k**⁺



tional fitting of the alkyl moiety of **2** inside the aromatic cavity of **1**.

Based on these considerations, the question arises as to whether the shape and dimension of the hosted alkyl moiety can influence the effectiveness of complexation: is there any cavity-filling effect? Is there a maximum or an optimal filling? Is there any quantitative parameter in suitable agreement with the experimental results?

The threading abilities of alkylbenzylammonium cations **2a–k**⁺ toward calix[6]arene **1** (Scheme 1) were studied by ¹H NMR titration experiments (CDCl₃, 298 K) by mixing equimolar quantities of **1** and **2** in CDCl₃ (3.8 mM solution) (Figures S24–S45).^{14a} With all the axles **2a–k**⁺, the pseudo[2]rotaxane stereoisomer with *endo*-alkyl stereochemistry was preferentially formed (see the Supporting Information for further details), following the *endo*-alkyl rule^{14c} previously reported. The apparent association constants (Table 1) for the formation of pseudo[2]rotaxanes **2a–k**⁺**1** were determined by integration of the slowly exchanging ¹H NMR signals for

both free and complexed hosts (see Supporting Information (SI)).¹⁵ In some instances, competition experiments with known pseudo[2]rotaxanes were also used (see SI).

At this point, to test the initial hypothesis, the log *K*_{app} data reported in Table 1 were correlated with the PC of the calix-cavity in pseudo[2]rotaxanes **2a–k**⁺**1** obtained using the DFT-optimized structures at the B3LYP-D3/6-31G(d,p) level of theory (D3 stands for Grimme's dispersion correction energy term¹⁶ and has been already used for DFT calculations in calixarene-based pseudorotaxane structures¹⁷). As expected, close inspection of the data reported in Table 1 revealed that the PC of the calix-cavity in pseudo[2]rotaxane **2a–k**⁺**1** structures overcomes the above-mentioned 55% rule (PC range = 72–100%) and the correlation coefficient of the linear fitting is low (*R*² = 0.38, Figure S46), indicating that there is a low correlation between the two parameters.

In Figure 1a–c, the superimposed calix[6]arene-wheels of the optimized structures **2**⁺**1** are reported (see also Figure




Figure 1. DFT-optimized structures, at B3LYP-D3/6-31G(d,p) level of theory, of the following: (a) superimposed calix[6]arene-wheels of **2**⁺**1** pseudorotaxanes (global minimum); (b) only **2a**⁺**1**, **2b**⁺**1**, and **2j**⁺**1**; (c) only **2d**⁺**1** and **2k**⁺**1**.

S47). Inspection of Figure 1a reveals that the calix[6]arene-wheels adopt two main conformations upon complexation: the cone-1,3,5-out (**2a–c**⁺**1** and **2e–j**⁺**1**; see Figure 1b) and the cone-1,4-out (**2d**⁺**1** and **2k**⁺**1**; see Figure 1c). Moreover, it is evident that, during the induced-fit recognition process, the calix-cavity deforms, allowing the change of the void volume of the receptor.

Close inspection of Figure 1b,c reveals a remarkable change of inclination of all the aromatic rings (−21° from yellow to coral, and +23° from yellow to purple, on average, respectively) of **1**, in pseudo[2]rotaxanes **2a**⁺**1**, **2b**⁺**1**, and **2k**⁺**1**. Thus, the conformational freedom of **1** ensures the

Table 1. Apparent Association Constants of **2**⁺**1** Pseudorotaxanes and Their PC, CC, and EC Parameters

Axle	<i>K</i> _{app} (M ^{−1}) ^a	log <i>K</i> _{app}	PC (%)	CC (%)	EC	Δ <i>G</i> _{Reorg} (kcal/mol)
2a ⁺ 1	(1.1 ± 0.2) × 10 ⁶	6.04	72	70	5.5	19.2
2b ⁺ 1	(4.8 ± 0.8) × 10 ³	3.68	100	72	7.2	28.0
2c ⁺ 1	(6.5 ± 0.9) × 10 ⁴	4.81	88	74	8.0	28.8
2d ⁺ 1	(2.4 ± 0.6) × 10 ³	3.38	94	71	10.1	32.0
2e ⁺ 1	(4.2 ± 0.6) × 10 ⁴	4.62	74	71	7.2	25.5
2f ⁺ 1	(3.6 ± 0.5) × 10 ²	2.56	82	77	9.5	36.9
2g ⁺ 1	(5.1 ± 0.6) × 10 ³	3.71	88	74	9.5	37.6
2h ⁺ 1	(1.7 ± 0.6) × 10 ³	3.23	86	75	10.8	31.9
2i ⁺ 1	(6.9 ± 0.8) × 10 ³	3.84	89	75	8.4	31.6
2j ⁺ 1	(2.9 ± 0.5) × 10 ³	3.46	87	76	10.5	35.3
2k ⁺ 1	(2.7 ± 0.4) × 10 ²	2.43	97	80	11.6	41.1

^aThe apparent association constant values were determined by mixing equimolar quantities of host and guest in CDCl₃ (3.8 mM solution each, 298 K) by using the following methods: (i) ¹H NMR competition experiment (**2a**⁺, **2b**⁺, **2c**⁺, and **2e**⁺); (ii) quantitative ¹H NMR experiment using 1,1,2,2-tetrachloroethane as the internal standard (**2k**⁺); (iii) integration of free and complexed ¹H NMR signals of host or guest (**2d**⁺, **2f**⁺, **2g**⁺, **2h**⁺, **2i**⁺, and **2j**⁺).

best fit around the alkyl portion of guest $2a^+$, $2b^+$, or $2k^+$ to establish an extended area of contact between them. In other words, the aromatic walls of calixarene host **1** move to wrap the guest and maximize the secondary interactions with it.

From the above data, it is evident that another useful and straightforward rule is necessary to assess the stability of these induced-fit complexes in order to predict the ideal cavity-filling effect. Initially, we reasoned that the maximization of weak secondary interactions should be parallel to the maximization of the contact surface between host and guest; therefore, we studied a new surface-based single-value parameter to address such point. With this aim, we considered the *Contacting Coefficient* (CC, eq 1) defined as the ratio between the molecular surface of the guest in close contact with the cavity surface (S_{Contact}) of the host, and the total surface of the guest (S_{Guest}).

$$\text{CC}(\%) = \frac{S_{\text{Contact}}}{S_{\text{Guest}}} \times 100 \quad (1)$$

This new parameter does not consider the host cavity volume and could be applied to host–guest processes that follow the induced-fitting mechanism. In addition, the CC parameter should be more directly related to the thermodynamic stability of the complex because it considers the host–guest contacting surface, which should be related to the extension of van der Waals and C–H $\cdots\pi$ interactions between them.

Starting from the complexes' DFT-optimized structures, the molecular surfaces of the guest inside the cavity (S_{Guest}) were computed by YASARA software, which also permits the direct measure of the contact surface between guest and host (S_{Contact}). From the ratio of those surfaces, CC (%) values of 70, 74, and 80 were calculated, through eq 1, for $2a^+C1$, $2c^+C1$, and $2k^+C1$ pseudo[2]rotaxanes (Table 1), respectively. The S_{Contact} is represented in red in Figure S48, while the S_{Guest} indicates the total molecular surface of the guest. Close inspection of Figure S48 reveals that, in addition to the S_{Contact} (in red), there are free portions of the guest's molecular surface not in contact with the calixarene cavity (in yellow).

Unfortunately, the CC of the whole set has only a discrete correlation coefficient ($R^2 = 0.54$, Figure S49).

At this point, we decided to evaluate another single-value geometrical parameter, which could take more directly into account the energy cost associated with the host reorganization upon induced-fit complexation. Therefore, we considered the *Expanding Coefficient* (EC, eq 2) defined as the ratio between the final and the initial cavity volumes of the host, i.e., the volume of the host cavity after the complexation ($V_{\text{complexed_Host}}$ at the global minimum) and that of the host cavity before the complexation ($V_{\text{free_Host}}$ at the global minimum).

$$\text{EC} = \frac{V_{\text{complexed_Host}}}{V_{\text{free_Host}}} \quad (2)$$

The actual values of $V_{\text{complexed_Host}}$ and $V_{\text{free_Host}}$ were measured by using the DFT-optimized structures of the separated host and guest for all the 2^+C1 complexes (see the SI)¹⁸ with the Caver software. From these values, the corresponding ECs were then calculated (Table 1), and a linear regression analysis was performed with the pertinent $\log K_{\text{app}}$ data. As shown in Figure 2, a good correlation coefficient ($R^2 = 0.74$) was obtained, demonstrating good linearity between the new EC parameter and the complex's thermodynamic stability. It is




Figure 2. Linear regression analysis of ECs vs $\log K_{\text{app}}$ values for 2^+C1 alkylbenzylammonium-based pseudorotaxane complexes.

evident that the EC parameter is now less affected than CC by the structural differences of the variously branched alkyl chains of $2a^+$.

The good correlation performance of the geometrical EC parameter induced us to consider its possible physical meaning. In fact, this EC can be considered an approximate geometrical measure of the energy cost paid by the host when it reorganizes itself from the initial lowest-energy conformation to the final geometry adopted in the complex. A higher EC value implies a higher deformation of the host, which in turn implies a higher energetical cost. From another point of view, the EC can be considered an approximate indirect inverse measurement of the preorganization of the host for the complexation of the given guest. The importance of this reorganization cost was first recognized by Cram,^{3c} who stated that “preorganization is a central determinant of binding power” leading to the formalization of the *Principle of Preorganization*, which states that “the more highly hosts and guests are organized for binding, and the lower the solvation before their complexation, the more stable will be their complexes”.

To verify the correctness of this “reorganization” point of view, we decided to calculate the energy of the host **1** in its bound conformation by performing a single-point calculation on each 2^+C1 complex after taking away the 2^+ guest. The difference between this single-point bound conformation energy and the lowest energy of **1** gives a ΔG_{Reorg} value, which can be considered the theoretically computed energetical cost for the above-mentioned reorganization of the host, from the initial lowest-energy conformation to the final geometry adopted in the complex. The ΔG_{Reorg} values computed for all the $2a-k^+C1$ complexes are reported in Table 1. Interestingly, these data seem to be in accord with the “reorganization” point of view, and indeed a good linear correlation ($R^2 = 0.79$) was found by regression analysis between ΔE_{Reorg} vs $\log K_{\text{app}}$ values (Figure S50). In summary, this analysis indicates that, under the above assumption of weak intermolecular interactions (dispersion forces, van der Waals interactions), the primary determinant to the stability of induced-fit complexes will be the degree of deformation with respect to the ground conformation.

In conclusion, we have defined the EC new parameter that can be correlated with the thermodynamic stability of supramolecular complexes governed by weak secondary interactions that obey the induced-fit model. The EC values show a good linear relationship with the $\log K_{\text{app}}$ of the respective pseudorotaxane complexes. This EC can be considered an approximate mechanical measure of the

reorganization energy cost paid by the host upon changing from the initial free lowest-energy conformation to the final bound geometry in the complex. This conclusion is in accordance with the Principle of Preorganization, by which the reorganization cost is a central determinant of binding power. We believe that the EC parameter can be of general applicability in all those instances in which no new strong intermolecular interactions (e.g., H-bonds) are generated during the induced-fit process.¹⁹ We anticipate future studies to test the ECs applicability in different systems, including the biological ones.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.orglett.1c00165>.

Experimental details, characterization data, NMR spectra, calculations details, and Cartesian coordinates (PDF)

■ AUTHOR INFORMATION

Corresponding Authors

Carmen Talotta – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy; orcid.org/0000-0002-2142-6305; Email: ctalotta@unisa.it

Antonio Rescifina – Dipartimento di Scienze del Farmaco e della Salute, Università di Catania, I-95125 Catania, Italy; orcid.org/0000-0001-5039-2151; Email: a.rescifina@unict.it

Pablo Ballester – Institute of Chemical Research of Catalonia (ICIQ), The Barcelona Institute of Science and Technology (BIST), 43007 Tarragona, Spain; orcid.org/0000-0001-8377-6610; Email: pballester@iciq.es

Placido Neri – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy; orcid.org/0000-0003-4319-1727; Email: neri@unisa.it

Authors

Gerardo Concilio – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy

Margherita De Rosa – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy; orcid.org/0000-0001-7451-5523

Annunziata Soriente – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy; orcid.org/0000-0001-6937-8405

Carmin Gaeta – Dipartimento di Chimica e Biologia “A. Zambelli”, Università di Salerno, I-84084 Fisciano (Salerno), Italy; orcid.org/0000-0002-2160-8977

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acs.orglett.1c00165>

Notes

The authors declare no competing financial interest.

■ REFERENCES

- (1) (a) Berg, J. M.; Tymoczko, J. L.; Gatto, G. J., Jr.; Stryer, L. *Biochemistry*, 9th ed.; Macmillan: 2019. (b) Taylor, M. R.; Simon, E. J.; Dickey, J. L.; Hogan, K. A. *Campbell Biology: Concepts & Connections*, 10th ed.; Pearson: 2020.
- (2) (a) Ehrlich, P.; Bolduan, C. *Collected studies on immunity*, 1st ed.; J. Wiley & Sons: New York, 1906. (b) Persch, E.; Dumele, O.; Diederich, F. Molecular Recognition in Chemical and Biological Systems. *Angew. Chem., Int. Ed.* **2015**, *54*, 3290.
- (3) (a) Cram, D. J. The Design of Molecular Hosts, Guests, and Their Complexes (Nobel Lecture). *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1009. (b) Lehn, J.-M. Supramolecular Chemistry - Scope and Perspectives, Molecules, Supramolecules, and Molecular Devices (Nobel Lecture). *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 89. (c) Pedersen, C. J. The Discovery of Crown Ethers (Nobel Lecture). *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1021.
- (4) (a) Cao, L. P.; Sekutor, M.; Zavalij, P. Y.; Mlinaric-Majerski, K.; Glaser, R.; Isaacs, L. Cucurbit[7]uril. Guest Pair with an Attomolar Dissociation Constant. *Angew. Chem., Int. Ed.* **2014**, *53*, 988. (b) Shetty, D.; Khedkar, J. K.; Park, K. M.; Kim, K. Can we beat the biotin-avidin pair?: cucurbit[7]uril-based ultrahigh affinity host-guest complexes and their applications. *Chem. Soc. Rev.* **2015**, *44*, 8747.
- (5) (a) Fischer, E. Ueber die optischen Isomeren des Traubenzuckers, der Gluconsäure und der Zuckersäure. *Ber. Dtsch. Chem. Ges.* **1890**, *23*, 2611. (b) Fischer, E. Einfluss der Configuration auf die Wirkung der Enzyme. *Ber. Dtsch. Chem. Ges.* **1894**, *27*, 2985.
- (6) (a) Thoma, J. A.; Koshland, D. E., Jr. Stereochemistry of enzyme, substrate, and products during beta-amylase action. *J. Biol. Chem.* **1960**, *235*, 2511. (b) Koshland, D. E., Jr. The Key-Lock Theory and the Induced Fit Theory. *Angew. Chem., Int. Ed. Engl.* **1995**, *33*, 2375. The following EC parameter can be also applied to a third “Conformational Selection” model; see: (c) Yang, L.-P.; Zhang, L.; Quan, M.; Ward, J. S.; Ma, Y.-L.; Zhou, H.; Rissanen, K.; Jiang, W. A supramolecular system that strictly follows the binding mechanism of conformational selection. *Nat. Commun.* **2020**, *11*, 2740.
- (7) Mecozzi, S.; Rebek, J., Jr. The 55% solution: A formula for molecular recognition in the liquid state. *Chem. - Eur. J.* **1998**, *4*, 1016.
- (8) (a) Puttreddy, R.; Beyeh, N. K.; Kalenius, E.; Ras, R. H. A.; Rissanen, K. 2-Methylresorcinarene: a very high packing coefficient in a mono-anion based dimeric capsule and the X-ray crystal structure of the tetra-anion. *Chem. Commun.* **2016**, *52*, 8115. (b) Lazar, A. I.; Biedermann, F.; Mustafina, K. R.; Assaf, K. I.; Hennig, A.; Nau, W. M. Nanomolar Binding of Steroids to Cucurbit[n]urils: Selectivity and Applications. *J. Am. Chem. Soc.* **2016**, *138*, 13022. (c) Shivanyuk, A.; Scarso, A.; Rebek, J., Jr. Coencapsulation of large and small hydrocarbons. *Chem. Commun.* **2003**, 1230. (d) Leontiev, A. V.; Saleh, A. W.; Rudkevich, D. M. Hydrophobic encapsulation of hydrocarbon gases. *Org. Lett.* **2007**, *9*, 1753.
- (9) (a) Nakano, K.; Sada, K.; Kurozumi, Y.; Miyata, M. Importance of Packing Coefficients of Host Cavities in the Isomerization of Open Host Frameworks: Guest-Size-Dependent Isomerization in Cholic Acid Inclusion Crystals with Monosubstituted Benzenes. *Chem. - Eur. J.* **2001**, *7*, 209. (b) Hayashida, O.; Sebo, L.; Rebek, J., Jr. Molecular discrimination of N-protected amino acid esters by a self-assembled cylindrical capsule: Spectroscopic and computational studies. *J. Org. Chem.* **2002**, *67*, 8291. (c) Scarso, A.; Trembleau, L.; Rebek, J., Jr. Encapsulation induces helical folding of alkanes. *Angew. Chem., Int. Ed.* **2003**, *42*, 5499. (d) Ajami, D.; Rebek, J., Jr. Coiled molecules in spring loaded devices. *J. Am. Chem. Soc.* **2006**, *128*, 15038. (e) Pirondini, L.; Bonifazi, D.; Cantadori, B.; Braiuca, P.; Campagnolo, M.; De Zorzi, R.; Geremia, S.; Diederich, F.; Dalcanale, E. Inclusion of methano[60]fullerene derivatives in cavitand-based coordination cages. *Tetrahedron* **2006**, *62*, 2008. (f) Purse, B. W.; Rebek, J., Jr. Self-fulfilling cavitands: Packing alkyl chains into small spaces. *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 2530. (g) Schramm, M. P.; Rebek, J., Jr. Moving targets: Recognition of alkyl groups. *Chem. - Eur. J.* **2006**, *12*, 5924. (h) Rebek, J., Jr. Contortions of encapsulated alkyl groups. *Chem. Commun.* **2007**, 2777. (i) Scarso, A.; Pellizzaro, L.; De Lucchi, O.; Linden, A.; Fabris, F. Gas hosting in enantiopure self-assembled oximes. *Angew. Chem., Int. Ed.* **2007**, *46*, 4972. (j) Gottschalk, T.; Jarowski, P. D.; Diederich,

F. Reversibly controllable guest binding in precisely defined cavities: selectivity, induced fit, and switching in novel resorcin[4]arene-based container molecules. *Tetrahedron* **2008**, *64*, 8307. (k) Liu, S.; Gibb, B. C. High-definition self-assemblies driven by the hydrophobic effect: synthesis and properties of a supramolecular nanocapsule. *Chem. Commun.* **2008**, 3709. (l) Raatikainen, K.; Huuskonen, J.; Kolehmainen, E.; Rissanen, K. Molecular pacman: Folding, inclusion, and X-ray structures of tri- and tetraamino piperazine cyclophanes. *Chem. - Eur. J.* **2008**, *14*, 3297. (m) Ménand, M.; Leroy, A.; Marrot, J.; Luhmer, M.; Jabin, I. Induced-Fit Encapsulation by a 1,3,5-Alternate Calix[6]arene. *Angew. Chem., Int. Ed.* **2009**, *48*, 5509. (n) Ajami, D.; Rebek, J., Jr. More Chemistry in Small Spaces. *Acc. Chem. Res.* **2013**, *46*, 990. (o) Brancatelli, G.; Gattuso, G.; Geremia, S.; Manganaro, N.; Notti, A.; Pappalardo, S.; Parisi, M. F.; Pisagatti, I. Encapsulation of biogenic polyamines by carboxylcalix[5]arenes: when solid-state design beats recognition in solution. *CrystEngComm* **2016**, *18*, 5012. (p) Zhang, W. Y.; Yang, D.; Zhao, J.; Hou, L. K.; Sessler, J. L.; Yang, X. J.; Wu, B. Controlling the Recognition and Reactivity of Alkyl Ammonium Guests Using an Anion Coordination-Based Tetrahedral Cage. *J. Am. Chem. Soc.* **2018**, *140*, 5248. (q) Taylor, C. G. P.; Argent, S. P.; Ludden, M. D.; Piper, J. R.; Mozaceanu, C.; Barnett, S. A.; Ward, M. D. One Guest or Two? A Crystallographic and Solution Study of Guest Binding in a Cubic Coordination Cage. *Chem. - Eur. J.* **2020**, *26*, 3054.

(10) Lee, T. C.; Kalenius, E.; Lazar, A. I.; Assaf, K. I.; Kuhnert, N.; Grun, C. H.; Janis, J.; Scherman, O. A.; Nau, W. M. Chemistry inside molecular containers in the gas phase. *Nat. Chem.* **2013**, *5*, 376.

(11) (a) Zurcher, M.; Gottschalk, T.; Meyer, S.; Bur, D.; Diederich, F. Exploring the flap pocket of the antimalarial target plasmepsin II: The "55% rule" applied to enzymes. *ChemMedChem* **2008**, *3*, 237. (b) Ehmke, V.; Kilchmann, F.; Heindl, C.; Cui, K. Q.; Huang, J.; Schirmeister, T.; Diederich, F. Peptidomimetic nitriles as selective inhibitors for the malarial cysteine protease falcipain-2. *MedChemComm* **2011**, *2*, 800.

(12) (a) Le Gac, S.; Marrot, J.; Reinaud, O.; Jabin, I. Allosterically coupled double induced fit for 1+1+1+1 self-assembly of a calix[6]trisamine, a calix[6]trisacid, and their guests. *Angew. Chem., Int. Ed.* **2006**, *45*, 3123. (b) Coquiere, D.; Marrot, J.; Reinaud, O. Spectacular induced-fit process for guest binding by a calix[6]arene Zn(II) funnel complex. *Org. Biomol. Chem.* **2008**, *6*, 3930.

(13) (a) Chang, S. K.; Cho, I. New Metal Cation-Selective Ionophores Derived from Calixarenes - Their Syntheses and Ion-Binding Properties. *J. Chem. Soc., Perkin Trans. 1* **1986**, 211. (b) Gutsche, C. D. *Calixarenes: an introduction*, 2nd ed.; RSC Publishing: Cambridge, 2008. (c) Neri, P.; Sessler, J. L.; Wang, M.-X. *Calixarenes and Beyond*; Springer International Publishing: Cham, 2016.

(14) (a) Gaeta, C.; Troisi, F.; Neri, P. endo-Cavity Complexation and Through-the-Annulus Threading of Large Calixarenes Induced by Very Loose Alkylammonium Ion Pairs. *Org. Lett.* **2010**, *12*, 2092. (b) Gaeta, C.; Talotta, C.; Farina, F.; Teixeira, F. A.; Marcos, P. M.; Ascenso, J. R.; Neri, P. Alkylammonium Cation Complexation into the Narrow Cavity of Dihomooxalix[4]arene Macrocycle. *J. Org. Chem.* **2012**, *77*, 10285. (c) Gaeta, C.; Talotta, C.; Margarucci, L.; Casapullo, A.; Neri, P. Through-the-Annulus Threading of the Larger Calix[8]arene Macrocycle. *J. Org. Chem.* **2013**, *78*, 7627. (d) Gaeta, C.; Talotta, C.; Mirra, S.; Margarucci, L.; Casapullo, A.; Neri, P. Catenation of Calixarene Annulus. *Org. Lett.* **2013**, *15*, 116. (e) Talotta, C.; Gaeta, C.; Qi, Z. H.; Schalley, C. A.; Neri, P. Pseudorotaxanes with Self-Sorted Sequence and Stereochemical Orientation. *Angew. Chem., Int. Ed.* **2013**, *52*, 7437. (f) Talotta, C.; De Simone, N. A.; Gaeta, C.; Neri, P. Calix[6]arene Threading with Weakly Interacting Tertiary Ammonium Axles: Generation of Chiral Pseudorotaxane Architectures. *Org. Lett.* **2015**, *17*, 1006.

(15) Schalley, C. A. *Analytical methods in supramolecular chemistry*, 2nd ed.; Wiley-VCH: Weinheim, 2012.

(16) Grimme, S. Semiempirical GGA-type density functional constructed with a long-range dispersion correction. *J. Comput. Chem.* **2006**, *27*, 1787.

(17) (a) Gaeta, C.; Talotta, C.; Neri, P. Pseudorotaxane orientational stereoisomerism driven by pi-electron density. *Chem. Commun.* **2014**, *50*, 9917. (b) Talotta, C.; Gaeta, C.; De Rosa, M.; Ascenso, J. R.; Marcos, P. M.; Neri, P. Alkylammonium Guest Induced-Fit Recognition by a Flexible Dihomooxalix[4]arene Derivative. *Eur. J. Org. Chem.* **2016**, 2016, 158.

(18) Particular attention was focused toward searching for the lowest-energy conformation of the very highly mobile host **1**, which can assume a series of almost degenerate conformations related to variously distorted 1,2,3-alternate geometries. These conformers gave very similar cavity volume values, which gave undistinguishable results in the EC calculations.

(19) As suggested by a Reviewer, we wish to make clear to the reader that for the general applicability of the proposed EC parameter further blind verification tests with unrelated data sets are necessary.