

## ISOLATION AND CHARACTERIZATION OF STABLE ALKYLIDENEPHOSPHINES


Masaaki YOSHIFUJI,\* Kozo TOYOTA, Katsuhiro SHIBAYAMA, and Naoki INAMOTO

Department of Chemistry, Faculty of Science, The University of  
Tokyo, Hongo, Tokyo 113

E and Z-P- $\alpha$ -(t-Butyldimethylsilyloxy)benzylidene(2,4,6-tri-t-butylphenyl)phosphines were isolated as stable compounds.

Primary phosphines are quite toxic and evil-smelling compounds. Generally they are very easily oxidized and, particularly, those containing the lower aliphatic moieties ignite spontaneously in air.<sup>1)</sup> Recently, we<sup>2)</sup> and others<sup>3)</sup> reported the preparation of a primary phosphine, 2,4,6-tri-t-butylphenylphosphine (1), which is quite stable to air and odorless. The phosphine 1 has turned out to be a useful starting material for another sterically protected phosphorus-containing unusual species.<sup>2-5)</sup> Phosphorus(III)-containing  $p_{\pi}-p_{\pi}$  double bond compounds have been of current interest, since some of those with  $-P=C<$ <sup>6-8)</sup>  $-P=N-$ ,<sup>9)</sup> and  $-P=P-$ <sup>10)</sup> were reported to be isolable materials if protected sterically.

We now report the preparation and characterization of alkylidenephosphines (6 and 7) starting from 1 via a silylphosphine 3 through a 1,3-silyl migration of an acylsilylphosphine intermediate 5 as shown below.



The phosphine 1 was metallated with butyllithium in tetrahydrofuran (THF) at room temperature under argon and the resulting phosphide 2 was quenched with *t*-butylchlorodimethylsilane to give *t*-butyldimethylsilyl(2,4,6-tri-*t*-butylphenyl)-phosphine (3) almost quantitatively after removal of THF, addition of pentane, and filtration of insoluble material. 3: oil;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ =7.21 (2H, d,  $^4J_{\text{PH}}=2.4$  Hz, arom.), 4.30 (1H, d,  $^1J_{\text{PH}}=211.1$  Hz, PH), 1.52 (18H, s, o-Bu<sup>t</sup>), 1.27 (9H, s, p-Bu<sup>t</sup>), 0.89 (9H, s, Si-Bu<sup>t</sup>), and -0.24 (6H, d,  $^3J_{\text{PH}}=8.8$  Hz, Si-Me); IR (KBr) 2375  $\text{cm}^{-1}$  (P-H); MS  $m/e$  392 ( $\text{M}^+$ ). The silylphosphine 3 was repeatedly lithiated with butyllithium in THF at room temperature under argon to give the corresponding phosphide 4 and allowed to react with benzoyl chloride to give *E-P*- $\alpha$ -(*t*-butyldimethylsilyloxy)benzylidene(2,4,6-tri-*t*-butylphenyl)phosphine (6) in 63% yield from 1 after purification by column chromatography ( $\text{SiO}_2$ ). 6: mp 121 - 124 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ =7.38 (2H, d,  $^4J_{\text{PH}}=1.5$  Hz, arom.), 7.0-6.4 (5H, m, Ph), 1.49 (18H, d,  $^5J_{\text{PH}}=0.7$  Hz, o-Bu<sup>t</sup>), 1.38 (9H, s, p-Bu<sup>t</sup>), 1.07 (9H, s, Si-Bu<sup>t</sup>), and 0.39 (6H, d,  $^3J_{\text{PH}}=1.8$  Hz, Si-Me); IR (KBr) 840  $\text{cm}^{-1}$  (P=C); MS  $m/e$  496 ( $\text{M}^+$ ); UV ( $\text{CH}_2\text{Cl}_2$ ) 247 ( $\epsilon$  14700) and 321 nm (10000). The *E*-isomer 6 was irradiated with a medium pressure mercury lamp for 2 h in benzene at 0 °C under argon to give the corresponding *Z*-isomer 7 almost quantitatively. 7: mp 102 - 104 °C;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$ =7.5 - 7.3 (5H, m, Ph), 7.25 (2H, d,  $^4J_{\text{PH}}=0.9$  Hz, arom.), 1.55 (18H, s, o-Bu<sup>t</sup>), 1.33 (9H, s, p-Bu<sup>t</sup>), 0.56 (9H, s, Si-Bu<sup>t</sup>), and -0.36 (6H, s, Si-Me); IR (KBr) 830  $\text{cm}^{-1}$  (P=C); MS (field desorption ionization)  $m/e$  496; UV ( $\text{CH}_2\text{Cl}_2$ ) 244 ( $\epsilon$  15800) and 306 nm (10600). Table 1 shows  $^{31}\text{P}$  NMR parameters of some phosphorus compounds obtained during the course of this work. Table 2 shows  $^{13}\text{C}\{^1\text{H}\}$  NMR data of 3, 6, and 7. It should be noted that the alkyls on the silicon atom in 6 interact more strongly with the phosphorus atom than those in 7<sup>8)</sup> in terms of spin-spin coupling constants  $J_{\text{PC}}$ .

The reaction might proceed via the benzoylsilylphosphine 5 first formed as an intermediate and the silyl group migrates in a 1,3-fashion to give the *E*-isomer 6. By considering the Dreiding models of 6 and 7, the *E*-isomer 6 is more crowded than the *Z*-isomer 7, therefore photoisomerization of *E* to *Z* is reasonably explained, however, 6 did not isomerize to 7 in refluxing benzene for several hours. A preliminary measurement of spin-lattice relaxation times ( $T_1$ ) by an inversion recovery FT method in  $^{31}\text{P}$  NMR (in  $\text{CDCl}_3$  at 30 °C) indicated that the more crowded *E*-isomer 6 has a shorter  $T_1$  (5.6 s) than that for the *Z*-isomer 7 (8.3 s), which is also consistent with this assignment of configurations.<sup>11)</sup> It should also be noted

TABLE 1.  $^{31}\text{P}$  NMR Parameters of Some Phosphorus Compounds

| Compound   |     | Solvent         | $\delta_{\text{P}}/\text{ppm}$ | $^1J_{\text{PH}}/\text{Hz}$ |
|--|-----|-----------------|--------------------------------|-----------------------------|
| $\text{ArPH}_2$  | (1) | $\text{CDCl}_3$ | -129.9                         | 210.6                       |
| $\text{ArP(H)Li}$                                      | (2) | THF             | -108.8                         | 170.9                       |
| $\text{ArP(H)SiMe}_3$                                  |     | THF             | -130.1                         | 212.4                       |
| $\text{ArP(SiMe}_3\text{)Li}$                          |     | THF             | -151.2                         | -----                       |
| $\text{ArP(H)SiMe}_2\text{Bu}^{\text{t}}$              | (3) | $\text{CDCl}_3$ | -136.2                         | 210.0                       |
| $\text{ArP(SiMe}_2\text{Bu}^{\text{t}}\text{)Li}$      | (4) | THF             | -156.0                         | -----                       |
| $\text{E-ArP=C(OSiMe}_2\text{Bu}^{\text{t}}\text{)Ph}$ | (6) | $\text{CDCl}_3$ | 160.3                          | -----                       |
| $\text{Z-ArP=C(OSiMe}_2\text{Bu}^{\text{t}}\text{)Ph}$ | (7) | $\text{CDCl}_3$ | 141.3                          | -----                       |

TABLE 2.  $^{13}\text{C}\{^1\text{H}\}$  NMR Data of Compounds 3, 6, and 7 in  $\text{CDCl}_3$ 

| Compound     | $\delta_{\text{C}}/\text{ppm}$ (assignment, multiplicity, $J_{\text{PC}}$ ) |   |
|--------------|---|---|
| <br><u>3</u> | 154.4 ( $\text{C}^2$ , d, $^2J=6.1$ Hz)                                     | 38.3 ( $\text{C}^5$ , s)                  |
|              | 146.9 ( $\text{C}^4$ , d, $^4J=1.2$ Hz)                                     | 34.5 ( $\text{C}^7$ , s)                  |
|              | 128.3 ( $\text{C}^1$ , d, $^1J=32.4$ Hz)                                    | 33.7 ( $\text{C}^6$ , d, $^4J=6.7$ Hz)    |
|              | 121.0 ( $\text{C}^3$ , d, $^3J=3.7$ Hz)                                     | 31.4 ( $\text{C}^8$ , s)                  |
|              |   | 27.4 ( $\text{C}^{10}$ , d, $^3J=2.4$ Hz) |
|              |   | 19.6 ( $\text{C}^9$ , d, $^2J=9.8$ Hz)    |
|              |   | -3.0 ( $\text{C}^{11}$ , d, $^2J=5.5$ Hz) |
|              |   |   |
| <br><u>6</u> | 192.1 ( $\text{C}^{12}$ , d, $^1J=42.7$ Hz)                                 | 38.2 ( $\text{C}^5$ , s)                  |
|              | 154.9 ( $\text{C}^2$ , d, $^2J=2.4$ Hz)                                     | 35.0 ( $\text{C}^7$ , s)                  |
|              | 150.6 ( $\text{C}^4$ , s)   | 32.6 ( $\text{C}^6$ , d, $^4J=6.7$ Hz)    |
|              | 140.6 ( $\text{C}^{13}$ , d, $^2J=14.0$ Hz)                                 | 31.5 ( $\text{C}^8$ , s)                  |
|              | 132.8 ( $\text{C}^1$ , d, $^1J=56.2$ Hz)                                    | 26.0 ( $\text{C}^{10}$ , d, $^5J=1.2$ Hz) |
|              | 127.3 ( $\text{C}^{16}$ , d, $^5J=4.3$ Hz)                                  | 18.5 ( $\text{C}^9$ , s)                  |
|              | 126.9 ( $\text{C}^{14}$ , d, $^3J=5.5$ Hz)                                  | -4.9 ( $\text{C}^{11}$ , d, $^4J=8.6$ Hz) |
|              | 126.7 ( $\text{C}^{15}$ , d, $^4J=8.5$ Hz)                                  |   |
|              | 122.2 ( $\text{C}^3$ , s)   |   |
| <br><u>7</u> | 188.5 ( $\text{C}^{12}$ , d, $^1J=51.9$ Hz)                                 | 38.1 ( $\text{C}^5$ , s)                  |
|              | 154.6 ( $\text{C}^2$ , d, $^2J=2.4$ Hz)                                     | 34.9 ( $\text{C}^7$ , s)                  |
|              | 149.0 ( $\text{C}^4$ , s)   | 32.9 ( $\text{C}^6$ , d, $^4J=6.7$ Hz)    |
|              | 142.6 ( $\text{C}^{13}$ , d, $^2J=32.3$ Hz)                                 | 31.5 ( $\text{C}^8$ , s)                  |
|              | 133.2 ( $\text{C}^1$ , d, $^1J=53.7$ Hz)                                    | 25.7 ( $\text{C}^{10}$ , s)               |
|              | 128.6 ( $\text{C}^{16}$ , d, $^5J=3.1$ Hz)                                  | 18.2 ( $\text{C}^9$ , s)                  |
|              | 128.0 ( $\text{C}^{14}$ , d, $^3J=10.4$ Hz)                                 | -3.1 ( $\text{C}^{11}$ , s)               |
|              | 127.5 ( $\text{C}^{15}$ , d, $^4J=14.0$ Hz)                                 |   |
|              | 121.6 ( $\text{C}^3$ , d, $^3J=1.2$ Hz)                                     |   |

that both 6 and 7 are stable for a long period of time in the pure state and neither isomerize nor polymerize.

Further studies on the reactions of 6 and 7 are in progress.<sup>12)</sup> We thank Mr. Hiroaki Shiraishi at the National Institute for Environmental Research, Japan, for obtaining FD-MS of 7 and Shin-Etsu Chemical Co., Ltd. for donation of the silyl halides used in this work. This work was also supported in part by the Scientific Research Grant-in-Aid (Nos. 543008, 57540276, and 58840023) from the Ministry of Education, Science and Culture of Japan.

#### References

- 1) L. Maier, "Organic Phosphorus Compounds," ed by G. M. Kosolapoff and L. Maier, Wiley-Interscience, New York (1971), Vol. I, p. 1.
- 2) M. Yoshifuji, K. Shibayama, N. Inamoto, T. Matsushita, and K. Nishimoto, J. Am. Chem. Soc., 105, 2459 (1983).
- 3) K. Issleib, H. Schmidt, and Chr. Wirkner, Z. Anorg. Allg. Chem., 488, 75 (1982); A. Zschunke, E. Bauer, H. Schmidt, and K. Issleib, *ibid.*, 495, 115 (1982).
- 4) M. Yoshifuji, K. Shibayama, K. Toyota, and N. Inamoto, Tetrahedron Lett., 24, in press.
- 5) A. H. Cowley, J. E. Kilduff, S. K. Mehrotra, N. C. Norman, and M. Pakulski, J. Chem. Soc., Chem. Commun., 1983, 528.
- 6) Th. Klebach, R. Lourens, and F. Bickelhaupt, J. Am. Chem. Soc., 100, 4886 (1978).
- 7) R. Appel, F. Knoll, and I. Ruppert, Angew. Chem., Int. Ed. Engl., 20, 731 (1981).
- 8) G. Becker, W. Becker, and O. Mundt, Phosphorus and Sulfur, 14, 267 (1983).
- 9) E. Niecke, R. Rüger, and W. W. Schoeller, Angew. Chem., Int. Ed. Engl., 20, 1034 (1981).
- 10) M. Yoshifuji, I. Shima, N. Inamoto, K. Hirotsu, and T. Higuchi, J. Am. Chem. Soc., 103, 4587 (1981); 104, 6167 (1982).
- 11) Ramarajan, M. D. Herd, and K. D. Berlin, Phosphorus and Sulfur, 11, 199 (1981).
- 12) Attempts to use chlorotrimethylsilane instead of t-butylchlorodimethylsilane have been unsuccessful because the silylphosphine obtained was unstable and the further lithiation was not efficient enough to give exclusively the corresponding phosphide ( $\delta_P$  -151.2 ppm in THF) and addition of butyllithium in excess amount cleaved the silyl group to give the starting 2 (see Table 1).

( Received August 12, 1983 )