

Stereoselective Synthesis of (\pm)-Ancistrofuran: Stereoselective Reduction of a γ -Hydroxyketone

Raymond Baker,* Paul D. Ravenscroft, and Christopher J. Swain

Department of Chemistry, The University, Southampton SO9 5NH, U.K.

A remarkably stereoselective reduction of a γ -hydroxyketone with two equiv. of lithium triethylborohydride has been observed and this effect has been suggested to originate from 'chelation control'; the reaction has been utilised in a short stereoselective synthesis of ancistrofuran.

The major soldiers of the West African termite, *Ancistrotermes cavithorax*, produce a defensive secretion which consists predominantly of ancistrofuran (**1**).¹ Although two syntheses of this compound have been reported^{2,3} neither route was stereoselective and both involved difficult separations of undesired isomers. We now report an efficient and stereoselective synthesis of ancistrofuran, together with a remarkably stereoselective reduction of a γ -hydroxyketone.


The starting material was the lactone (**2**) which is readily available *via* the mercury(II) ion initiated cyclisation of homogeranic acid.³ Reduction of the lactone (**2**) with diisobutylaluminium hydride yielded the hemiacetal (**3**)³ in 60% yield; subsequent reaction with two equiv. of 3-lithiofuran yielded a 1:1 mixture of two diastereoisomeric diols [(**4a**) and (**4b**)]. This mixture of diols was not normally separated but oxidised with MnO₂ in CH₂Cl₂ to yield the ketone (**5**)[†] in 90% yield.

The ketone (**5**) is also available in 60% yield in one step from the reaction of lactone (**2**) with one equiv. of 3-lithiofuran. The ketone (**5**), a white crystalline solid m.p. 77–78 °C (Et₂O), exists predominantly in the γ -hydroxyketone form there being no evidence of lactol formation, ν_{\max} (CCl₄), 3300

br.s, 1670s cm⁻¹; ¹H n.m.r. δ 8.05 (1H, m, H^{1‡}), 7.2 (1H, m, H^{3‡}), 6.73 (1H, m, H^{2‡}), 2.8 (2H, m, CH₂CO-), 1.07 (1H, m, -CH-), 1.1–1.8 (6H, m, 3 \times CH₂), 1.1 (3H, s, -CH₃), 0.82 (6H, s, gem 2 \times CH₃).

Reduction of ketone (**5**) with LiAlH₄, ZnBH₄, and NaBH₄ all yielded mixtures of the diols (**4a**) and (**4b**) (Table 1). However, reduction with two equiv. of lithium triethylborohydride in tetrahydrofuran proceeded in a highly stereoselective manner to yield the required diol (**4a**) as the only product in 88% yield as a white crystalline solid m.p. 85–87 °C (Et₂O). Treatment of the diol with one equiv. of toluene-*p*-sulphonyl chloride and two equiv. of pyridine in methylene chloride gave a quantitative yield of ancistrofuran§ (**1**) (Scheme 1) *via* esterification of the secondary alcohol and S_N2 displacement by the tertiary alcohol.

It is proposed that the basis of this remarkably high degree of stereoselectivity in the reduction is due to initial reaction of the reducing agent with the tertiary alcohol and subsequent chelation to the carbonyl to yield the cyclic structure (**6**). Attack of hydride from a second equiv. of the reducing agent can now only occur from the least hindered β -face. Stereoselective reduction of α -hydroxyketones⁴ and β -hydroxyketones⁵ have been reported and Kishi *et al.* have developed conditions for the reduction of some γ - and δ -epoxy ketones.⁶ This appears to be the first example of stereoselective reduction of a γ -hydroxyketone. The use of 'chelation




Scheme 1. Reagents and conditions: i, AlBu₃H, toluene, -78 °C; ii, 3-lithiofuran (2 equiv.), tetrahydrofuran (THF), -78 °C; iii, MnO₂, CH₂Cl₂, 12 h; iv, 3-lithiofuran (1 equiv.), THF, -78 °C; v, LiBHET₃ (2 equiv.), THF, -78 °C; vi, *p*-MeC₆H₄SO₂Cl (1 equiv.), pyridine (2 equiv.), CH₂Cl₂, room temp., 48 h.

† All new compounds gave satisfactory elemental microanalyses and/or accurate mass measurements.

Table 1. Reduction of γ -hydroxyketone (**5**).

Reducing Agent	(4a) ^a	(4b) ^a	Yield (%) ^b
LiAlH ₄	40	60	80
NaBH ₄	50	50	84
ZnBH ₄	60	40	66
LiBHET ₃ H	100	0	88

^a Ratio of products measured by g.l.c. (3 m \times 3 mm), 5% OV101 at 190 °C. ^b Refers to isolated yield.



‡ Furanoid protons, designated on structure (**5**).

§ Identical with the authentic sample.

control' in the reaction of Grignards⁷ and cuprates with carbonyl compounds⁸ is well documented and the present work represents a further example of stereocontrol in reactions of acyclic systems.

We acknowledge financial support from the S.E.R.C.

Received, 24th October 1983; Com. 1389

References

- 1 R. Baker, P. H. Briner, and D. A. Evans, *J. Chem. Soc., Chem. Commun.*, 1978, 410.
 - 2 R. Baker, P. H. Briner, and D. A. Evans, *J. Chem. Soc., Chem. Commun.*, 1978, 981.
 - 3 T. R. Hoye and A. J. Caruso, *J. Org. Chem.*, 1981, **46**, 1198.
 - 4 T. Nakata, T. Tanaka, and T. Oishi, *Tetrahedron Lett.*, 1983, **24**, 2053.
 - 5 T. Nakata, M. Fukui, M. Ohtsuka, and T. Oishi, *Tetrahedron Lett.*, 1983, **24**, 2661.
 - 6 T. Fukuyama, B. Vranesic, D. P. Negri, and Y. Kishi, *Tetrahedron Lett.*, 1978, **19**, 2741.
 - 7 D. B. Collum, J. H. McDonald, and W. C. Still, *J. Am. Chem. Soc.*, 1980, **102**, 2118.
 - 8 J. A. Schneider and W. C. Still, *Tetrahedron Lett.*, 1980, **21**, 1035.
-