# **Reaction of AlR<sub>3</sub> with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]: Facile Degradation of [B(C\_6F\_5)\_4]^- by Transient "[AlR<sub>2</sub>]<sup>+</sup>**"

Manfred Bochmann\* and Mark J. Sarsfield

School of Chemistry, University of Leeds, Leeds LS2 9JT, U.K.

Received May 19, 1998

Trimethylaluminum reacts with  $[CPh_3][B(C_6F_5)_4]$  at elevated temperatures to give a mixture of  $AlMe_{3-x}(C_6F_5)_x$  compounds, depending on the Al/B ratio.  $AlBu_{3}^{i}$  reacts significantly faster under  $\beta$ -hydride abstraction. The Al-C<sub>6</sub>F<sub>5</sub> species rapidly react with Cp<sub>2</sub>ZrMe<sub>2</sub> or  $[Cp_2ZrMe]^+$  under C<sub>6</sub>F<sub>5</sub> transfer to give poorly active Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>); this reaction may have implications for the deactivation of  $Cp_2ZrMe_2/AlR_3/[CPh_3][B(C_6F_5)_4]$  olefin polymerization catalysts.

## Introduction

As is now well established, highly active olefin polymerization catalysts can be generated by reacting metallocene dialkyls  $Cp_2MR_2$  (M = Ti, Zr, Hf) with cation-generating agents, notably B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, [HNMe<sub>2</sub>Ph]- $[B(C_6F_5)_4]$ , or  $[CPh_3][B(C_6F_5)_4]$ , to give complexes in which the highly electrophilic active species [Cp<sub>2</sub>MR]<sup>+</sup> is stabilized by weakly coordinating but chemically very robust counteranions such as  $[MeB(C_6F_5)_3]^-$  and  $[B(C_6F_5)_4]^{-.1}$  Because of the high sensitivity of these catalyst systems toward trace impurities the addition of aluminum alkyls is usually beneficial. For example, we have shown previously that the addition of 1 equiv of AlMe<sub>3</sub> stabilizes the 14-electron cation [Cp<sub>2</sub>ZrMe]<sup>+</sup> by formation of a heterobinuclear complex,  $[Cp_2Zr(\mu -$ Me)<sub>2</sub>AlMe<sub>2</sub>]<sup>+,2</sup> and Chien et al. have developed highly active catalysts based on mixtures of metallocene dichlorides and  $[CPh_3][B(C_6F_5)_4]$  in the presence of an excess of AlMe<sub>3</sub> or AlEt<sub>3</sub>.<sup>3</sup>

The success of these catalysts depends on the selective reaction of  $B(C_6F_5)_3$  or  $[CPh_3][B(C_6F_5)_4]$  with the transition metal alkyl complex in preference to reaction with the main group metal alkyl additive. This may seem at first glance surprising since it is not obvious why reactive electrophiles such as  $B(C_6F_5)_3$  or  $CPh_3^+$  should fail to react with a strong alkylating reagent such as AlMe<sub>3</sub>. Indeed, we have shown earlier<sup>4</sup> that the reaction of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> with Cp<sub>2</sub>AlMe to give  $[AlCp_2]^+[MeB(C_6F_5)_3]^$ is fast and quantitative even at low temperatures, and Jordan et al.<sup>5</sup> recently reported the facile methyl anion abstraction from (benzamidinato)AlMe<sub>2</sub> compounds to give cationic aluminum methyl complexes.

Two reasons attracted our interest in analogous reactions of AlMe<sub>3</sub>. First, although the Cp<sub>2</sub>MR<sub>2</sub>/[CPh<sub>3</sub>]- $[B(C_6F_5)_4]/AIR_3$  catalysts mentioned above<sup>2,3</sup> show very high initial activities, there is typically a significant reduction of activity with time. Little is known about the chemistry that underlies such catalyst deactivation processes,<sup>6</sup> and we speculated that a decay of the pentafluorophenylborate anion, possibly with the participation of AlR<sub>3</sub>, might be important. Second, while there are numerous cationic aluminum alkyls [AlR<sub>2</sub>- $(L)_2$ <sup>+</sup> stabilized by N- and O-ligands,<sup>7</sup> with the exception of the metallocenes  $[AlCp_2]^+$  and  $[AlCp_2]^+$ <sup>8</sup> cationic  $[AIR_2]^+$  species free of donor ligands are not known, and we were intrigued by the possibility of stabilizing such species with weakly coordinating anions such as  $[B(C_6F_5)_4]^-$ .

### **Results**

A solution of  $[CPh_3][B(C_6F_5)_4]$  in toluene- $d_8$  was treated with a 10-fold molar excess of AlMe<sub>3</sub> and

<sup>(1)</sup> Reviews: Bochmann, M. J. Chem. Soc., Dalton Trans. 1996, 255. Brintzinger, H. H.; Fischer, D.; Mülhaupt, R.; Rieger, B.; Waymouth, R. M. Angew. Chem., Int. Ed. Engl. 1995, 34, 1143. Guram, A. S.; Jordan, R. F. In Comprehensive Organometallic Chemistry II; Abel, E. W., Stone, F. G. A., Wilkinson, G., Eds.; Elsevier: Oxford, 1995; Chapter 12.

<sup>(2)</sup> Bochmann, M.; Lancaster, S. J. Angew. Chem., Int. Ed. Engl. 1994, 33, 1634. Bochmann, M.; Lancaster, S. J. J. Organomet. Chem. **1995**. *497*. 55.

<sup>(3)</sup> Chien, J. C. W.; Xu, B. Makromol. Chem., Rapid Commun. 1993, 14, 109. Chien, J. C. W.; Tsai, W. M. Makromol. Chem., Macromol. Symp. 1993, 66, 141. Chien, J. C. W.; Song, W.; Rausch, M. D. J. Polym. Sci., Part A: Polym. Chem. 1994, 32, 2387.
(4) Dawson, D. M.; Bochmann, M. Angew. Chem. 1996, 108, 2371;

Angew. Chem., Int. Ed. Engl. 1996, 35, 2226.

<sup>(5)</sup> Coles, M. P.; Jordan, R. F. J. Am. Chem. Soc. 1997, 119, 8125. Ihara, E.; Young, V. G.; Jordan, R. F. J. Am. Chem. Soc. 1998, 120, 8277. Radzewich, C. E.; Coles, M. P.; Jordan, R. F. J. Am. Chem. Soc. 1998, 120, 9384.

<sup>(6)</sup> For examples of reactions likely to be involved in catalyst deactivation see: Kaminsky, W.; Sinn, H. Liebigs Ann. 1975, 424. Kaminsky, W.; Vollmer, H. J. Liebigs Ann. 1975, 438. Kaminsky, W.; Kopf, J.; Sinn, H.; Vollmer, H. J. Angew. Chem. **1976**, 88, 688; Angew. Chem., Int. Ed. Engl. **1976**, 15, 629. Takahashi, T.; Kasai, K.; Suzuki, N.; Nakajima, K.; Negishi, E. *Organometallics* **1994**, *13*, 3413. Shapiro, P. J.; Cotter, W. D.; Schaefer, W. P.; Labinger, J. A.; Bercaw, J. E. J. Am. Chem. Soc. 1994. 116, 4623. Bochmann, M.; Cuenca, T.; Hardy, D. T. J. Organomet. Chem. 1994, 484, C10. Lancaster, S. J.; Robinson, O. B.; Bochmann, M.; Coles, S. J.; Hursthouse, M. B. Organometallics 1995, 14, 2456. Jiménez Pindado, G.; Thornton-Pett, M.; Bowkamp, M.; Meetsma, A.; Hessen, B.; Bochmann, M. Angew. Chem., Int. Ed. Engl. 1997, 36, 2358. Jiménez Pindado, G.; Thornton-Pett, M.; Bochmann, M. J. Chem. Soc., Dalton Trans. 1997, 3115.

<sup>(7)</sup> Tobias, R. S. Organomet. Chem. Rev. 1966, 1, 93. Bott, S. B.; Alvanipour, A.; Morley, S. D.; Atwood, D. A.; Means, C. M.; Coleman, Arvanpour, A.; Morley, S. D.; Atwood, D. A.; Means, C. M.; Coleman, A. W.; Atwood, J. L. Angew. Chem., Int. Ed. Engl. 1987, 26, 485. Engelhardt, L. M.; Kynast, U.; Raston, C. L.; White, A. H. Angew. Chem., Int. Ed. Engl. 1987, 26, 681. Self, M. F.; Pennington, W. T.; Laske, J. A.; Robinson, G. H. Organometallics 1991, 10, 36. Uhl, W.; Wagnen, L. Eangle, D.; Bourn, C. Z. Anger Allg. Cham. 1909, 612 Wagner, J.; Fenske, D.; Baum, G. Z. Anorg. Allg. Chem. 1992, 612,
 25. Atwood, D. A.; Jones, R. A.; Cowley, A. H.; Bott, S. G.; Atwood, J.
 L. J. Organomet. Chem. 1992, 425, C1. Atwood, D. A.; Jegier, J. Chem. Commun. 1996, 1507.

<sup>(8)</sup> Dohmeier, C.; Schnöckel, H.; Robl, C.; Schneider, U.; Ahlrichs, R. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1655.

warmed to 60 °C. The reaction was monitored by <sup>1</sup>H, <sup>11</sup>B, and <sup>19</sup>F NMR spectroscopy over 4.5 h. A slow reaction takes place; formation of MeCPh<sub>3</sub> is observed, along with BMe<sub>3</sub> [<sup>1</sup>H NMR (toluene- $d_8$ , 60 °C)  $\delta$  0.74; <sup>11</sup>B NMR  $\delta$  86.8], and a new set of signals appears in the <sup>19</sup>F NMR spectrum, at  $\delta$  –121.5 (*o*-F), –152.2 (*p*-F), and –160.8 (*m*-F), in addition to unreacted [CPh<sub>3</sub>]-[B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]. In the presence of excess AlMe<sub>3</sub> the reaction is first-order in [CPh<sub>3</sub><sup>+</sup>], with  $k_{obs} = 5 \times 10^{-5} \text{ s}^{-1}$  (at 60 °C). In the absence of AlMe<sub>3</sub> toluene solutions of [CPh<sub>3</sub>]-[B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] are stable under these conditions.

This observation demonstrates that methyl abstraction from AlMe<sub>3</sub> by CPh<sub>3</sub><sup>+</sup> is very much less favorable than with transition metal methyl complexes but occurs slowly at elevated temperatures to give a transient species "[AlMe<sub>2</sub>]<sup>+</sup>[B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup>", which immediately decomposes to B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>). Evidently cations such as [AlMe<sub>2</sub>]<sup>+</sup> (or possibly higher homologues, e.g., [Al<sub>2</sub>Me<sub>5</sub>]<sup>+</sup> or [Al<sub>3</sub>Me<sub>8</sub>]<sup>+</sup>, which might be envisaged in the presence of excess AlMe<sub>3</sub>) are too electrophilic to be stabilized even by [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> and immediately react with abstraction of C<sub>6</sub>F<sub>5</sub><sup>-</sup>. The difference in reactivity between [AlMe<sub>2</sub>]<sup>+</sup> and the thermally stable complexes [AlCp<sub>2</sub>]<sup>+</sup> and [(benzamidinate)AlMe]<sup>+</sup> is most probably due to the absence of any form of  $\pi$ -donor stabilization in the dialkyl species.

Further ligand exchange between  $AIMe_{3-x}(C_6F_5)_x$  and  $B(C_6F_5)_3$  can eventually lead to the formation of  $AI(C_6F_5)_3$  and  $BMe_3$ , which in an open system escapes into the gas phase, thereby shifting the equilibrium:

AlMe $(C_6F_5)_2$  + BMe $(C_6F_5)_2$   $\rightarrow$  Al $(C_6F_5)_3$  + BMe<sub>2</sub> $(C_6F_5)$ 

$$BMe_2(C_6F_5) + AIMe_3 \rightarrow AIMe_2(C_6F_5) + BMe_3$$

The reaction of  $B(C_6F_5)_3$  with AlMe<sub>3</sub> in toluene to afford  $Al(C_6F_5)_3$ •0.5toluene and  $BMe_3$  via a similar sequence of ligand redistribution processes has recently been described in a patent application by Biagini et al. as a convenient synthetic method for  $Al(C_6F_5)_3$ .<sup>9,10</sup>

Under the conditions of the NMR experiment in the presence of an excess of AlMe<sub>3</sub> the major product of the reaction between  $[CPh_3][B(C_6F_5)_4]$  and AlMe<sub>3</sub> is  $[AlMe_2-(C_6F_5)]_2$ . The identity of this species was confirmed by its independent preparation from a 2:1 mixture of AlMe<sub>3</sub> and Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene. Treating 2 equiv of AlMe<sub>3</sub> with Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene at 20 °C in toluene-*d*<sub>8</sub> in an NMR tube causes an immediate shift of the methyl resonance in the <sup>1</sup>H NMR spectrum from  $\delta$  -0.37 to -0.08 ppm. Upon cooling the sample to -60 °C the methyl signal is



split into seven observable peaks (although additional smaller methyl signals appear obscured by more intense peaks), while three sets of  $C_6F_5$  signals are observed in the <sup>19</sup>F NMR spectrum. None of the signals are attributable to Al<sub>2</sub>Me<sub>6</sub>.<sup>11</sup> Warming to -20 °C results in broadening of all the signals in the <sup>1</sup>H and <sup>19</sup>F NMR spectra, and at 20 °C the single methyl chemical shift value observed is the averaged chemical shift of the methyl groups of a number of isomers that are in rapid equilibrium with each other (Scheme 1). Similarly, 1:2 mixtures of AlMe<sub>3</sub> and Al( $C_6F_5$ )<sub>3</sub>·0.5toluene give predominantly [AlMe( $C_6F_5$ )<sub>2</sub>]<sub>2</sub>. The pertinent NMR data of the  $C_6F_5$  species reported here are collected in Table 1.

Mixing Al( $C_6F_5$ )<sub>3</sub>·0.5toluene with AlMe<sub>3</sub> in a ratio of 1:10 at room temperature in toluene on a preparative scale, followed by removal of the solvent in vacuo with mild heating, leads to the recovery of Al( $C_6F_5$ )<sub>3</sub>·0.5toluene. Isolation of the initial product, AlMe<sub>2</sub>( $C_6F_5$ ), was not possible due to the evidently very facile ligand exchange and quantitative removal of excess AlMe<sub>3</sub> even under these mild conditions. In related recent work, Roesky et al. reported the synthesis of AlMe<sub>2</sub>( $C_6F_5$ ) by the reaction of AlMe<sub>2</sub>Cl with LiC<sub>6</sub>F<sub>5</sub> as an uncharacterized intermediate which was converted to Al( $C_6F_5$ )<sub>3</sub> by heating to 180 °C in vacuo.<sup>12</sup> Considering the facile ligand exchange in this system (Scheme 1) we find that such high temperatures are not required to shift the equilibrium toward Al( $C_6F_5$ )<sub>3</sub>.

The <sup>19</sup>F NMR chemical shifts of all three aluminum  $C_6F_5$  species AlMe<sub>3-x</sub>( $C_6F_5$ )<sub>x</sub> (x = 1-3) are remarkably similar, with changes in the  $\delta$  value of *para*-F being most indicative, though identification is possibly by <sup>1</sup>H NMR. More pronounced changes are seen when the exchange process is suppressed by adduct formation with diethyl ether. The addition of excess Et<sub>2</sub>O to NMR samples of AlMe<sub>2</sub>( $C_6F_5$ ) in toluene-d<sub>8</sub> at 20 °C results in the formation of  $AlMe_2(C_6F_5)$ ·Et<sub>2</sub>O as the major product, together with small amounts of  $AIMe(C_6F_5)_2$ .  $Et_2O$  and  $AlMe_3 \cdot Et_2O$ . Likewise, addition of  $Et_2O$  to AlMe $(C_6F_5)_2$  gives AlMe $(C_6F_5)_2$ ·Et<sub>2</sub>O as the major product, with small amounts of  $AlMe_2(C_6F_5) \cdot Et_2O$  and AlMe<sub>3</sub>·Et<sub>2</sub>O also being detectable. The adducts were characterized by <sup>1</sup>H, <sup>19</sup>F, and <sup>13</sup>C{<sup>1</sup>H} NMR spectroscopy (Table 1). Surprisingly, although Roesky et al.<sup>12</sup> isolated a crystalline THF adduct of  $Al(C_6F_5)_3$ , the compound appears not to form a complex with Et<sub>2</sub>O in toluene solution, possibly for steric reasons.

To test the suitability of the reaction of AlMe<sub>3</sub> with  $[CPh_3][B(C_6F_5)_4]$  as a synthetic route for Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, a 10:1 mixture of the two reactants was heated at 65 °C over 14 h. Triphenylethane was produced, and the color of the solution changed from orange to pale yellow.

<sup>(9)</sup> Biagini, P.; Lugli, G.; Abis, L.; Andreussi, P. Eur. Patent Appl. EP 0 694 548 A1, 1996 (Enichem Elastomeri S.r.l.).

<sup>(10)</sup> Ligand redistribution processes in the borate  $[BBu_n(C_6F_5)_{4-n}]^-$  have been observed: Dioumaev, V. K.; Harrod, J. F. *Organometallics* **1997**, *16*, 2798.

<sup>(11)</sup> Williams, K. C.; Brown, T. L. J. Am. Chem. Soc. 1966, 88, 5460.
(12) Belgardt, T.; Storre, J.; Roesky, H. W.; Noltemeyer, M.; Schmidt, H. G. Inorg. Chem. 1995, 34, 3821.

Table 1 I and - I twik Data of Cars Compound	Table 1.	<sup>1</sup> H and	<sup>19</sup> F NMR	Data of	C <sub>6</sub> F <sub>5</sub>	Compound
--	----------	--------------------	---------------------	---------	-------------------------------	----------

			$^{19}$ F NMR ( <i>J</i> <sub>FF</sub> , Hz)		
compound	<sup>1</sup> H NMR	<i>o</i> -F	<i>p</i> -F	m-F <sup>b</sup>	
$AlMe_2(C_6F_5)^a$	-0.08 (Me)	-122.5 (16.9)	-151.9 (br)	-161.0	
$AlMe(C_6F_5)_2^a$	0.04 (Me)	-122.4 (16.9)	-151.4	-161.0	
Al(C <sub>6</sub> F <sub>5</sub> ) <sub>3</sub> •0.5toluene		$-122.7 (20.9)^{a}$	$-151.3 (21.2)^{a}$	$-161.1^{a}$	
		$-121.9(21)^{c}$	$-149.6 (21)^{c}$	$-160.4^{\circ}$	
$AlMe_2(C_6F_5) \cdot OEt_2^a$	-0.36 (Me)	-122.5 (28.2)	-155.1 (19.8)	-161.8	
	0.69, 3.31 (Et <sub>2</sub> O)				
$AlMe(C_6F_5)_2 \cdot OEt_2^a$	-0.18 (Me)	-123.1 (28.2)	-152.8 (19.8)	-161.0	
	0.88, 3.32 (Et <sub>2</sub> O)				
$Cp_2ZrMe(\mu-Me)Al(C_6F_5)_3$ <sup>c</sup>	-0.26 (µ-Me)	-124.0 (16.9)	-155.2 (19.8)	-162.6	
	0.51 (Zr-Me)				
	6.44 (Cp)				
$Cp_2ZrMe(C_6F_5)^{a}$	0.31 (Me)	-115.2 (25.4)	-156.7 (19.8)	-162.2	
-	5.66 (Cp)				

<sup>a</sup> Toluene-d<sub>8</sub>, 20 °C. <sup>b</sup> Multiplets or broadened signals. <sup>c</sup> CD<sub>2</sub>Cl<sub>2</sub>, -60 °C.

#### Scheme 2

 $A|Bu_{3}^{i} + [CPh_{3}][B(C_{6}F_{5})_{4}] \longrightarrow Ph_{3}CH + "[A|Bu_{2}^{i}]^{+}[B(C_{6}F_{5})_{4}]^{-n} + \sum$ 

 $AIBu_{3-x}^{i}(C_{6}F_{5})_{x} + BBu_{x}^{i}(C_{6}F_{5})_{3-x}$ 

Removal of the solvent and drying for 4 h in vacuo produced a light brown oily solid. The <sup>19</sup>F NMR spectrum shows a major product in approximately 50% yield with chemical shifts identical to an authentic sample of  $Al(C_6F_5)_3$ . 0.5 toluene. However, the remaining 50% of the signals were due to other  $C_6F_5$  species which could not be identified, and the reaction is evidently less clean than the reaction of  $AlMe_3$  with  $B(C_6F_5)_3$ .

In contrast to the rather slow reaction between AlMe<sub>3</sub> and [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] which is initiated by *alkyl* abstraction, the reaction of AlBu<sup>i</sup><sub>3</sub> with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] involves *hydride* abstraction<sup>13</sup> and isobutene elimination (Scheme 2). This reaction proceeds very much faster: at 20 °C a toluene solution of [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] discolors immediately on addition of AlBu<sup>i</sup><sub>3</sub>, and the <sup>19</sup>F NMR shows that all [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> has been consumed, with formation of BBu<sup>i</sup><sub>3</sub>, BBu<sup>i</sup><sub>2</sub>(C<sub>6</sub>F<sub>5</sub>), and BBu<sup>i</sup>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> (but not, under these conditions, B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>).

**Implications for the Deactivation Process of Metallocene Catalysts.** The foregoing results show that the reaction of aluminum alkyls with  $[CPh_3]$ - $[B(C_6F_5)_4]$  or  $B(C_6F_5)_3$  can lead to aluminum pentafluorophenyl compounds, with borate degradation in the presence of aluminum alkyls bearing  $\beta$ -hydrogens being particularly facile. As pointed out in the Introduction, catalyst deactivation processes are as yet little understood, and the possible contribution of  $Al-C_6F_5$  species to catalyst deactivation was therefore of interest.

Treatment of Cp<sub>2</sub>ZrMe<sub>2</sub> with an equimolar amount of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene in CD<sub>2</sub>Cl<sub>2</sub> at -60 °C leads immediately to the quantitative formation of Cp<sub>2</sub>ZrMe-( $\mu$ -Me)Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, according to <sup>1</sup>H, <sup>13</sup>C, and <sup>19</sup>F NMR spectroscopy (Scheme 3). The <sup>1</sup>H NMR spectrum shows Cp, Zr-Me, and Zr-Me-Al resonances at  $\delta$  6.44, 0.51, and -0.26 ppm, respectively. In the <sup>13</sup>C{<sup>1</sup>H} spectrum the Cp and Zr-Me signals are present at  $\delta$  113.9 and 40.5 ppm, the  $\mu$ -Me signal is broad and centered at  $\delta$ 7.9 ppm. Only one set of C<sub>6</sub>F<sub>5</sub> signals in the <sup>19</sup>F spectrum are observed (Table 1) which are entirely





different from Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene in CD<sub>2</sub>Cl<sub>2</sub> at -60 °C (cf. Table 1). Attempts to isolate this product resulted only in the formation of a number of Cp complexes, one of which is identified as Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>). We conclude that Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> interacts with Cp<sub>2</sub>ZrMe<sub>2</sub> in a way similar to B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub><sup>14</sup> to give initially zwitterionic methyl-bridged complexes, but that redistribution of C<sub>6</sub>F<sub>5</sub> ligands from aluminum to zirconium proceeds much more readily than in the case of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> (Scheme 2), as observed for example by Marks et al., who noted the slow decomposition of Cp<sup>"</sup><sub>2</sub>ZrMe( $\mu$ -Me)B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> to Cp<sup>"</sup><sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) by Me/C<sub>6</sub>F<sub>5</sub> exchange with a half-life of 10 h at room temperature [Cp<sup>"</sup> = C<sub>5</sub>H<sub>3</sub>(SiMe<sub>3</sub>)<sub>2</sub>].<sup>15</sup>

At 20 °C a benzene- $d_6$  solution of Cp<sub>2</sub>ZrMe<sub>2</sub> reacts immediately with Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> in a 1:1 molar ratio with formation of two major Cp compounds, one of which was identified as Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>), while the other could not be assigned. The formation of Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub><sup>16</sup> could not be detected, even in the presence of an excess of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>. By contrast, the reaction of [Cp<sub>2</sub>Zr( $\mu$ -H)-(C<sub>6</sub>F<sub>5</sub>)]<sub>2</sub> with [CPh<sub>3</sub>][BBu<sub>n</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>4-n</sub>] has been reported to proceed rapidly to give Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> and other decomposition products.<sup>10</sup> On a preparative scale Al-(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>•0.5toluene reacts with Cp<sub>2</sub>ZrMe<sub>2</sub> in a 1:3 ratio to give Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) in 54% yield. The same product results from a 1:1 mixture of Cp<sub>2</sub>ZrMe<sub>2</sub> and AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) in toluene- $d_8$  at 20 °C; the reaction is quantitative.

<sup>(13)</sup> For examples of  $\beta$ -H abstraction of main group alkyls by CPh<sub>3</sub><sup>+</sup> see: Jerkunica, J. M.; Traylor, T. G. *J. Am. Chem. Soc.* **1971**, *93*, 6278. Hannon, J. S.: Traylor, T. G. *J. Org. Chem.* **1981**, *46*, 3645.

<sup>(14)</sup> Yang, X.; Stern, C. L.; Marks, T. J. J. Am. Chem. Soc. 1991, 113, 3623. Deck, P. A.; Marks, T. J. J. Am. Chem. Soc. 1995, 117, 6128.
Siedle, A. R.; Newmark, R. A. J. Organomet. Chem. 1995, 497, 119.
(15) Yang, X.; Stern, C. L.; Marks, T. J. J. Am. Chem. Soc. 1994, 116, 10015. Gomez, R.; Green, M. L. H.; Haggit, J. L. J. Chem. Soc.

Dalton Trans. 1996, 939. (16) Chaudhari, M. A.; Stone, F. G. A. J. Chem. Soc. A 1966, 838.

The above discussion shows that under certain circumstances, and particularly if AlBu<sup>i</sup><sub>3</sub>/[CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] mixtures are used in catalyst preparation, the transfer of  $C_6F_5$  to zirconium may be facile. The deactivating effect of  $C_6F_5$  was assessed by reacting  $Cp_2ZrMe(C_6F_5)$ with  $[CPh_3][B(C_6F_5)_4]$  and  $B(C_6F_5)_3$ , to give  $[\{Cp_2Zr (C_6F_5)_2(\mu-Me)$  [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] and Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)( $\mu-Me$ )B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>, respectively.<sup>17</sup> Although at comparatively high zirconium concentrations (1 mM) some ethylene polymerization was observed, the activities are very poor compared to Cp<sub>2</sub>ZrMe<sub>2</sub>/activator mixtures. On the other hand, if conditions that can give rise to "[AlR<sub>2</sub>]+" (such as premixing  $CPh_3^+$  with AlR<sub>3</sub>) are avoided, the system  $Cp_2ZrMe_2/AlR_3/[CPh_3][B(C_6F_5)_4]$  is thermally stable. For example, heating equimolar mixtures of Cp<sub>2</sub>ZrMe<sub>2</sub> and  $[CPh_3][B(C_6F_5)_4]$  in toluene- $d_8$  for 1 h at 60 °C, followed by heating for a further 2 h in the presence of AlMe<sub>3</sub> (Al/Zr = 4), led to a darkening of the samples, but no degradation of the  $[B(C_6F_5)_4]^-$  anion was observed by <sup>19</sup>F NMR.

## Conclusion

Aluminum alkyl cocatalysts in MAO-free metallocene systems are capable of reacting with the catalyst activators B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> and [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] at elevated temperatures with formation of  $AlMe_{3-x}(C_6F_5)_x$  compounds. The degradation of  $[B(C_6F_5)_4]^-$  is initiated by a highly electrophilic transient " $[AIR_2]^+$ " species. The value of *x* depends on the ratio of C<sub>6</sub>F<sub>5</sub>:Me ligands in the system. Aluminum alkyls carrying  $\beta$ -hydrogens such as AlBu<sup>i</sup><sub>3</sub> react with CPh<sub>3</sub><sup>+</sup> under hydride abstraction and alkene elimination; this reaction leads to the rapid consumption of  $[B(C_6F_5)_4]^-$  even at ambient temperatures. Al $(C_6F_5)_3$ reacts with  $Cp_2ZrMe_2$  to give the adduct  $Cp_2ZrMe(\mu$ -Me)Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> which is stable only at low temperatures and on warming reacts under C<sub>6</sub>F<sub>5</sub> transfer from aluminum to zirconium. At room temperature and above  $Cp_2ZrMe_2$  reacts rapidly with  $AlMe_{3-x}(C_6F_5)_x$  to give  $Cp_2ZrMe(C_6F_5)$ . The ethene polymerization activity of  $Cp_2ZrMe(C_6F_5)/B(C_6F_5)_3$ , mixtures is about 3 orders of magnitude lower that that of  $Cp_2ZrMe_2/B(C_6F_5)_3$ , an illustration of the deactivating consequence of  $C_6F_5$ transfer reactions.

## **Experimental Section**

All manipulations were carried out with the use of Schlenk, vacuum-line, and glovebox techniques. Light petroleum (bp 40-60 °C), toluene, and diethyl ether were distilled under nitrogen from sodium-potassium alloy. Deuterated solvents were dried over Linde 4 Å molecular sieves and deoxygenated via several freeze-thaw cycles. AlMe<sub>3</sub> (97%, Aldrich) was used as purchased. The compounds Cp<sub>2</sub>ZrMe<sub>2</sub>,<sup>18</sup> B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>,<sup>19</sup> and  $[CPh_3][B(C_6F_5)_4]^{20}$  were prepared by literature methods. The <sup>1</sup>H, <sup>13</sup>C, <sup>11</sup>B, and <sup>19</sup>F NMR spectra were recorded on a Bruker DPX300 instrument. <sup>1</sup>H and <sup>13</sup>C NMR spectra were referenced to residual solvent peaks relative to TMS; <sup>11</sup>B and <sup>19</sup>F spectra were referenced externally to BF3·Et2O and CFCl3, respectively.

Synthesis of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene. This compound was prepared by modification of a literature procedure.<sup>9</sup> CAU-**TION:** Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> has been reported to detonate on attempted sublimation at elevated temperatures.<sup>21</sup> A solution of B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> (3.55 g, 6.94 mmol) in toluene (20 mL) was added to a solution of AlMe<sub>3</sub> (0.50 g, 6.94 mmol) in toluene (40 mL) at room temperature. A precipitate formed after 0.5 h, and the reaction was left to stir for 16 h. The solid was isolated and dried under vacuum for 3 h to give  $Al(C_6F_5)_3$ .0.5toluene as a white microcrystalline solid (1.8 g). The filtrate was concentrated and placed in the freezer overnight and provides another 1.5 g. combined yield 3.3 g (5.74 mmol, 83%)

Reaction of AlMe<sub>3</sub> with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]. For monitoring the reaction by NMR, a solution of  $[CPh_3][B(C_6F_5)_4]$  (0.014 g, 15  $\mu$ mol) in toluene- $d_8$  (0.4 mL) was treated with a solution of AlMe<sub>3</sub> (0.6 mL, 0.11 M, 66  $\mu$ mol) in toluene- $d_8$  and immediately placed in the spectrometer heated to 60 °C. The reaction was monitored via <sup>1</sup>H and <sup>19</sup>F NMR over a period of 4.5 h. Signals for AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) grew in and became the major species after 4.5 h. No other products were seen. The reaction is first-order in triphenylmethyl salt,  $k_{obs} = 5 \times 10^{-5} \text{ s}^{-1}$ .

In a separate experiment, a toluene solution (5 mL) of [CPh<sub>3</sub>]- $[B(C_6F_5)_4]$  (0.202 g, 0.22 mmol) was treated with a toluene solution of AlMe<sub>3</sub> (3 mL, 1.5 M, 4.50 mmol). The reagents were heated at 65 °C for 14 h before the solvent was removed, and the light brown oily residue was dried under vacuum for 4 h. A sample of the crude product contained  $Al(C_6F_5)_3$  as the major product in 50% yield by <sup>19</sup>F NMR spectroscopy. NMR <sup>19</sup>F (toluene- $d_8$ )  $\delta - 122.7$  (d, J = 20.9 Hz,  $\bar{6}F$ , o-F), -151.3 (t, J =21.2 Hz, 3F, p-F), -161.1 (m, 6F, m-F).

Reaction of AlBu<sup>i</sup><sub>3</sub> with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]. An NMR tube containing [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (0.012 mmol) was reacted with a solution of AlBu<sup>i</sup><sub>3</sub> (0.070 mmol) in toluene-d<sub>8</sub> at 20 °C. The solution turned from orange to colorless *immediately*, and the <sup>19</sup>F NMR spectrum contained no signals due to  $[B(C_6F_5)_4]^-$ . Apart from the reaction products isobutene and triphenylmethane, signals were observed for AlBui<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) [<sup>1</sup>H NMR  $\delta$ 1.99 (m, 1H, CH), 0.97 (d, 6H, J = 6.5 Hz, CH<sub>3</sub>), 0.28 (d, 2H, J = 7.2 Hz, CH<sub>2</sub>); <sup>19</sup>F NMR  $\delta$  -120.7 (m, 2F), -152.5 (t, 1F, J = 19.8 Hz), -161.6 (m, 2F)], BBu<sup>i</sup>(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub> [-130.9 (m, 2F), -147.4 (t, 2F, J = 21.2 Hz), -161.1 (m, 2F)], and BBu<sup>i</sup><sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) [-134.3 (m, 2F), -153.3 (t, 1F, J = 19.7 Hz), -161.8 (m, 2F)]but not  $B(C_6F_5)_3$ . The <sup>11</sup>B and <sup>1</sup>H NMR spectra show the presence of BBu<sup>i</sup><sub>3</sub> [<sup>1</sup>H NMR  $\delta$  1.56 (m, 1H, CH), 1.16 (d, 2H, J = 7.0 Hz, CH<sub>2</sub>), 0.88 (d, 6H, J = 6.5 Hz, CH<sub>3</sub>); <sup>11</sup>B NMR  $\delta$ 88.1].22

Reaction of 2 Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene with AlMe<sub>3.</sub> A solution of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene (43 mg, 76  $\mu$ mol) in toluene- $d_8$  (0.5 mL) was treated with a solution of AlMe<sub>3</sub> (0.1 mL, 0.38 M, 38  $\mu$ mol) in toluene- $d_8$  at 20 °C to form [AlMe(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>]<sub>2</sub>. The sample was cooled to -60 °C, and three major and three minor methyl signals could be seen in the <sup>1</sup>H NMR spectrum at  $\delta$  (rel int) 0.34(1.2), 0.26(8.9), 0.09(7.2), -0.26(10.0), -0.34(4.5), -0.51(3.8), and -0.57 (3.1). Three sets of peaks for  $C_6F_5$  groups are seen in the <sup>19</sup>F NMR spectrum, at  $\delta$  (rel int) –122.4 (40.0), -122.7 (12.6), -123.1 (3.2) (o-F), -150.1 (1.6), -151.2 (20.0), -152.4 (6.3) (p-F), and -160.1 (3.2), -160.4 (40.0), -160.9 (12.6) (*m*-F). At 20 °C [AlMe( $C_6F_5$ )<sub>2</sub>]<sub>2</sub> is characterized by the following NMR parameters: <sup>1</sup>H NMR (toluene- $d_8$ , 20 °C)  $\delta$  0.04 (s, br, Me);  $^{13}\mathrm{C}\{^{1}\mathrm{H}\}$  NMR (toluene- $d_{8}$ , 20 °C)  $\delta$  150.0 (dm,  $J_{\mathrm{CF}}$ = 236 Hz, o-CF), 142.3 (dm,  $J_{CF}$  = 254 Hz, p-CF), 137.2 (dm,  $J_{\rm CF} = 254$  Hz, *m*-CF), -6.9 (Me); <sup>19</sup>F NMR (toluene- $d_8$ , 20 °C)  $\delta$  -122.4 (d, J = 16.9 Hz, 6F, o-F), -151.4 (s, br, 3F, p-F), -161.0 (s, br, 6F, m-F).

Formation of AlMe(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>·Et<sub>2</sub>O. An excess of diethyl ether was added to the above solution at 20 °C. <sup>1</sup>H NMR

<sup>(17)</sup> A mixture of 50  $\mu$ mol Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) and 50  $\mu$ mol B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> in (17) A mixture of 50 μmol Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) and 50 μmol B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> in toluene (50 mL) at 50 °C under 1 bar ethene gave 0.2 g of PE after 1 h, corresponding to a productivity of 4 kg PE (mol Zr)<sup>-1</sup>·bar<sup>-1</sup>·h<sup>-1</sup>. Although recrystallized Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) was used, the possibility of the presence of traces of Cp<sub>2</sub>ZrMe<sub>2</sub> cannot be entirely discounted.
(18) Samuel, E.; Rausch, M. D. J. Am. Chem. Soc. 1973, 95, 6263.
(19) Massey, A. G.; Park, A. J. J. Organomet. Chem. 1992, 430.
(20) Bochmann, M.; Lancaster, S. J. J. Organomet. Chem. 1992, 434.
C1 Chian L.C. W. Pausch, M. D. J. Am. Chem. Soc. 1991, 1/3, 8570.

C1. Chien, J. C. W.; Rausch, M. D. J. Am. Chem. Soc. 1991, 113, 8570.

<sup>(21)</sup> Pohlmann, J. L. W.; Brinckmann, F. E. Z. Naturforsch. B 1965, 20b, 5. Chambers, R. D. Organomet. Chem. Rev. **1966**, *1*, 279. (22) Davies, A. G.; Hare, D. G.; White, R. F. M. J. Chem. Soc. **1960**,

<sup>1040.</sup> Mikhailov, B. M.; Kuimova, M. E. Bull. Acad. Sci. USSR, Div. Chem. Sci. 1979, 288.

(toluene- $d_8$ , 20 °C):  $\delta$  3.32 (q, J = 7.0 Hz, 4H, OC $H_2$ CH<sub>3</sub>), 0.88 (t, J = 7.0 Hz, 6H, OCH<sub>2</sub>CH<sub>3</sub>), -0.18 (t,  ${}^5J_{\rm HF}$  = 1.2 Hz, 3H, Me–Al).  ${}^{13}$ C{ $}^{1}$ H} (toluene- $d_8$ , 20 °C):  $\delta$  67.1 (OCH<sub>2</sub>CH<sub>3</sub>), 14.5 (OCH<sub>2</sub>CH<sub>3</sub>), -6.91 (s, br, Me–Al).  ${}^{19}$ F (toluene- $d_8$ , 20 °C):  $\delta$  -123.1 (d, J = 28.2 Hz, 6 F, o-F), -152.8 (t, J = 19.8 Hz, 3 F, p-F), -161.0 (m, 6 F, *m*-F). Small amounts of AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)· Et<sub>2</sub>O (see below) and AlMe<sub>3</sub>·Et<sub>2</sub>O are also present. AlMe<sub>3</sub>· Et<sub>2</sub>O:  ${}^{1}$ H (toluene- $d_8$ , 20 °C):  $\delta$  -0.51 (s, 9H, Me–Al);  ${}^{13}$ C{ $}^{1}$ H} (toluene- $d_8$ , 20 °C)  $\delta$  -7.7 (s, br, Me–Al). The diethyl ether signals quoted are a time-averaged value of all AlMe<sub>3-x</sub>(C<sub>6</sub>F<sub>5</sub>)<sub>x</sub>· Et<sub>2</sub>O species and free diethyl ether in equilibrium.

**Reaction of Al**( $C_6F_5$ )<sub>3</sub>·0.5toluene with 2AlMe<sub>3</sub>. A solution of Al( $C_6F_5$ )<sub>3</sub>·0.5toluene (43 mg, 76  $\mu$ mol) in toluene- $d_8$  (0.2 mL) was treated with a solution of AlMe<sub>3</sub> (0.4 mL, 0.38 M, 152  $\mu$ mol) in toluene- $d_8$  and placed in the spectrometer at 20 °C. The sample was cooled to -60 °C, and at least seven separate methyl signals could be seen, which coalesce to one singlet at 20 °C. [AlMe( $C_6F_5$ )<sub>2</sub>]<sub>2</sub>: <sup>1</sup>H NMR (toluene- $d_8$ , 20 °C)  $\delta$  -0.08 (s, br, 3H, Me); <sup>13</sup>C{<sup>1</sup>H} NMR (toluene- $d_8$ , 20 °C)  $\delta$  150.1 (dm,  $J_{CF} = 249$  Hz, o-C), 142.3 (dm,  $J_{CF} = 249$  Hz, p-CF), 137.2 (dm,  $J_{CF} = 249$  Hz, m-CF), -7.3 (Me); <sup>19</sup>F NMR (toluene- $d_8$ , 20 °C)  $\delta$  -122.5 (d, J = 16.9 Hz, 6F, o-F), -151.9 (s, br, 3F, p-F), -161.0 (s, br, 6F, m-F).

**Formation of AlMe**<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>)·Et<sub>2</sub>O. An excess of diethyl ether was added to the above solution. <sup>1</sup>H NMR (toluene-*d*<sub>8</sub>, 20 °C):  $\delta$  3.31 (q, J = 7.0 Hz, 4 H, OCH<sub>2</sub>CH<sub>3</sub>), 0.69 (t, J = 7.2 Hz, 6H, OCH<sub>2</sub>CH<sub>3</sub>), -0.36 (t, <sup>5</sup>*J*<sub>HF</sub> = 1.1 Hz, 6H, Me–Al). <sup>13</sup>C-{<sup>1</sup>H} (toluene-*d*<sub>8</sub>, 20 °C):  $\delta$  67.4 (OCH<sub>2</sub>CH<sub>3</sub>), 13.4 (OCH<sub>2</sub>CH<sub>3</sub>), -8.1 (s, br, Me–Al). <sup>19</sup>F (toluene-*d*<sub>8</sub>, 20 °C):  $\delta$  -122.5 (d, J = 28.2 Hz, 6F, o-F), -155.1 (t, J = 19.8 Hz, 3F, p-F), -161.8 (m, 6F, *m*-F). Small amounts of AlMe(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>·Et<sub>2</sub>O and AlMe<sub>3</sub>·Et<sub>2</sub>O are also present.

**Generation of Cp<sub>2</sub>ZrMe**( $\mu$ -**Me**)**Al**(**C**<sub>6</sub>**F**<sub>5</sub>)<sub>3</sub>. A solution of Cp<sub>2</sub>ZrMe<sub>2</sub> (12 mg, 48  $\mu$ mol) in CD<sub>2</sub>Cl<sub>2</sub> (0.6 mL) was treated with Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene (29.4 mg, 51  $\mu$ mol) in an NMR tube at -78 °C and then placed in the spectrometer cooled to -60 °C. An immediate reaction occurs providing a quantitative conversion to Cp<sub>2</sub>ZrMe( $\mu$ -Me)Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>, -60 °C):  $\delta$  6.44 (s, 10H, Cp), 0.51 (s, 3H, Zr-Me), -0.26 (s, 3H, Me-Al). <sup>13</sup>C{<sup>1</sup>H} NMR (CD<sub>2</sub>Cl<sub>2</sub>, -60 °C):  $\delta$  149.8 (dm, *J*<sub>CF</sub> = 234 Hz, *o*-CF), 141.0 (dm, *J*<sub>CF</sub> = 249 Hz, *p*-CF), 136.7 (dm, *J*<sub>CF</sub> = 226 Hz, *m*-CF), 113.9 (Cp), 40.5 (Zr-Me), 7.9 (br, Me-Al). <sup>19</sup>F NMR (CD<sub>2</sub>Cl<sub>2</sub>, -60 °C):  $\delta$  -124.0 (d, *J* = 16.9 Hz, 6H, *o*-F), -155.2 (t, *J* = 19.8 Hz, 3F, *p*-F), -162.6 (m, 6F, *m*-F).

**Generation of Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)**( $\mu$ -**Me**)**B**(C<sub>6</sub>F<sub>5</sub>)**.** A sample of Cp<sub>2</sub>ZrMeC<sub>6</sub>F<sub>5</sub> (19 mg, 0.053 mmol) in CD<sub>2</sub>Cl<sub>2</sub> (0.3 mL) was treated with B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> (28.5 mg, 0.056 mmol) in CD<sub>2</sub>Cl<sub>2</sub> (0.3 mL) at -78 °C and placed in the spectrometer at -50 °C. Immediate formation of Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)( $\mu$ -Me)B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> was observed. The Zr-C<sub>6</sub>F<sub>5</sub> moiety shows hindered rotation at low temperature. <sup>1</sup>H NMR (CD<sub>2</sub>Cl<sub>2</sub>):  $\delta$  0.56 (bs, 3H, B–Me), 6.60 (s, 10H, Cp). <sup>13</sup>C{<sup>1</sup>H} NMR:  $\delta$  116.1 (Cp), 33.8 (B–Me). <sup>11</sup>B NMR:  $\delta$  -11.1. <sup>19</sup>F NMR (-80 °C):  $\delta$  112.6 (d, 1F, *J* = 28.0 Hz, *o*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -114.1 (d, 1F, *J* = 28.0 Hz, *o*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -152.3 (t, 1F, *J* = 19.8 Hz, *p*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -158.2 (t, 3F, *J* = 21.2 Hz, *p*-F of B-C<sub>6</sub>F<sub>5</sub>), -159.4 (m, 1F, *m*-F of B-C<sub>6</sub>F<sub>5</sub>), -160.0 (m, 1F, *m*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -163.3 (m, 6F, *m*-F of B-C<sub>6</sub>F<sub>5</sub>).

**Generation of** [{**Cp**<sub>2</sub>**Zr**(**C**<sub>6</sub>**F**<sub>5</sub>)<sub>2</sub>( $\mu$ -**Me**)][**B**(**C**<sub>6</sub>**F**<sub>5</sub>)<sub>4</sub>]. A sample of Cp<sub>2</sub>ZrMeC<sub>6</sub>F<sub>5</sub> (16 mg, 0.045 mmol) in CD<sub>2</sub>Cl<sub>2</sub> (0.3 mL) was treated with [CPh<sub>3</sub>][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>] (0.024 mmol) in CD<sub>2</sub>Cl<sub>2</sub> (0.3 mL) at -78 °C and placed in the spectrometer at -50 °C. There was a slow reaction to give [{Cp<sub>2</sub>Zr(C<sub>6</sub>F<sub>5</sub>)<sub>2</sub>( $\mu$ -Me)][B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]. Complete conversion took 160 min at -30 °C. <sup>1</sup>H NMR (CD<sub>2</sub>-Cl<sub>2</sub>):  $\delta$  6.56 (s, 20H, Cp), -0.029 (bs, 3H,  $\mu$ -Me). <sup>13</sup>C{<sup>1</sup>H} NMR:  $\delta$  115.0 (Cp),  $\mu$ -Me not observed. <sup>11</sup>B NMR:  $\delta$  -14.5. <sup>19</sup>F NMR (-80 °C):  $\delta$  113.1 (d, 2F, J = 25.0 Hz,  $\rho$ -F of Zr-C<sub>6</sub>F<sub>5</sub>), -119.3 (d, 2F, J = 25.0 Hz,  $\rho$ -F of Zr-C<sub>6</sub>F<sub>5</sub>), -158.3 (m, 2F, *m*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -159.3 (m, 2F, *m*-F of Zr-C<sub>6</sub>F<sub>5</sub>), -160.9 (m, 4F, p-F of B-C<sub>6</sub>F<sub>5</sub>), -164.9 (m, 8F, *m*-F of B-C<sub>6</sub>F<sub>5</sub>).

**Reaction of Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene with Cp<sub>2</sub>ZrMe<sub>2</sub> at 20 °C.** A C<sub>6</sub>D<sub>6</sub> solution of Cp<sub>2</sub>ZrMe<sub>2</sub> (24 mg, 95  $\mu$ mol) was treated with Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene (18.5 mg, 32  $\mu$ mol) in an NMR tube at room temperature. An immediate reaction occurs providing a quantitative conversion to Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>).

In a preparative scale reaction two separate Schlenk tubes were charged with Cp<sub>2</sub>ZrMe<sub>2</sub> (0.641 g, 2.55 mmol) in one and  $Al(C_6F_5)_3$ ·0.5toluene (0.467 g, 0.85 mmol) in the other. Both were dissolved in toluene (ca. 10 mL). The Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> solution was added to Cp<sub>2</sub>ZrMe<sub>2</sub> at 20 °C. The reaction mixture was stirred for 10 min before filtering and cooling to -78 °C over 48 h. A pale yellow precipitate was isolated, washed twice with light petroleum, and dried in vacuo, to yield  $Cp_2ZrMe(C_6F_5)$ (0.55 g, 1.36 mmol, 54%). <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>, 20 °C):  $\delta$  5.66 (s, 10H, Cp), 0.31 (t,  ${}^{5}J_{\rm HF}$  = 4.1 Hz, 3H, Me).  ${}^{13}C{}^{1}H$  NMR (C<sub>6</sub>D<sub>6</sub>, 20 °C):  $\delta^{13}C{^1H}$  NMR (CD<sub>2</sub>Cl<sub>2</sub>, -60 °C):  $\delta^{145.4}$  (dm,  $J_{CF}$  = 226 Hz, o-CF), 139.5 (dm, J<sub>CF</sub> = 251 Hz, p-CF), 136.8 (dm, J<sub>CF</sub> = 255 Hz, *m*-CF), 111.4 (Cp), 45.4 (t,  ${}^{4}J_{CF}$  = 6.0 Hz, Me).  ${}^{19}F$ NMR (C<sub>6</sub>D<sub>6</sub>, 20 °C):  $\delta$  -115.2 (d, J = 25.4 Hz, 6H, o-F), -156.7 (t, J = 19.8 Hz, 3F, p-F), -162.2 (m, 6F, m-F). MS (EI): m/e(%) 402 (2) (M<sup>+</sup>), 387 (80) [M<sup>+</sup> - Me], 239 (85) [M<sup>+</sup> - C<sub>6</sub>F<sub>5</sub>], 220 (100) [Cp<sub>2</sub>Zr].

**Reaction of AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) with Cp<sub>2</sub>ZrMe<sub>2</sub>.** A sample of AlMe<sub>2</sub>(C<sub>6</sub>F<sub>5</sub>) (228  $\mu$ mol), prepared from Al(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>·0.5toluene (43 mg, 76  $\mu$ mol) and AlMe<sub>3</sub> (0.4 mL, 0.38 M, 152  $\mu$ mol) in toluened<sub>8</sub> as discussed above, was added to Cp<sub>2</sub>ZrMe<sub>2</sub> (57 mg, 228 mmol) at 20 °C. An immediate reaction takes place with quantitative conversion to Cp<sub>2</sub>ZrMe(C<sub>6</sub>F<sub>5</sub>) and AlMe<sub>3</sub>, identified by comparison with authentic samples.

**Thermal Stability of** [**C**<sub>P2</sub>**ZrMe**][**B**(**C**<sub>6</sub>**F**<sub>5</sub>)<sub>4</sub>]. In a control experiment, 13 mg (14  $\mu$ mol) of [CPh<sub>3</sub>][B(C<sub>6</sub>**F**<sub>5</sub>)<sub>4</sub>] and Cp<sub>2</sub>ZrMe<sub>2</sub> (14  $\mu$ mol) were mixed in an NMR tube at room temperature in 0.5 mL of toluene-*d*<sub>8</sub>. The <sup>19</sup>F NMR spectrum of the mixture was monitored at 60 °C for 1 h; there was no indication of anion degradation. Next, AlMe<sub>3</sub> (66  $\mu$ mol) was added and heating was continued for another 2 h. No change in the signals for [B(C<sub>6</sub>F<sub>5</sub>)<sub>4</sub>]<sup>-</sup> was observed.

**Acknowledgment.** This work was supported by BP Chemicals Ltd., Sunbury. We thank Drs. P. Brès and B. Dorer for helpful discussions.

OM980400J