

by the donors of the Petroleum Research Fund, administered by the American Chemical Society, and the NSF (CHE81-07629) are gratefully acknowledged.

Registry No. 3, 4646-69-9; 7, 92012-54-9; 8, 92012-55-0; 9, 286-85-1; 11, 92012-56-1; 12, 92012-57-2; 13, 92012-58-3; Ph₂CO, 119-61-9; PhC(O)CH₃, 98-86-2.

Synthesis of Dialkylruthenium Nitrosyls and Thermal Conversion to Ruthenium Oximate, Carboxamide, and Cyano Complexes

Mark D. Seidler and Robert G. Bergman*

Department of Chemistry, University of California
Berkeley, California 94720


Received June 8, 1984

Several years ago, in search of homogeneous models for the metal-catalyzed oxidation of propylene to acrylonitrile, a group at Du Pont¹ uncovered a reaction in which an allyl group and a nitric oxide ligand underwent conversion to an oxime π -complexed to nickel. This discovery presaged a growing number of processes in which coordinated nitric oxide can be used to form new carbon-nitrogen bonds in organic molecules.² We wish to report here some unusual ruthenium chemistry germane to the understanding of such processes, which includes the observation of (1) the transformation of coordinated nitric oxide into oxime functionality σ -bound to the metal, (2) in one system, the rearrangement to an isomeric amide structure, and (3) in another system, dehydration of the oxime, leading to coordinated cyanide.

Our results are summarized in Scheme I. Nitric oxide was introduced into the coordination sphere of ruthenium by treating Cp^{*}Ru(CO)₂Cl (Cp^{*} = η^5 -C₅Me₅)³ with NOCl⁴ in methylene chloride at room temperature. After chromatography on silica gel and recrystallization from boiling toluene, air-stable dark green crystals of Cp^{*}(NO)RuCl₂ (**1**) were obtained in 53% yield.⁵ After experimenting with a variety of alkylating agents, the transformation of **1** to the dialkyl complexes **2** was found to be most cleanly achieved with trialkylaluminum compounds. Thus treatment of a slurry of **1** in toluene with a dilute solution of AlMe₃ gave red-orange crystalline **2a** in 67% yield. The diethyl complex **2b** was prepared in a similar manner from AlEt₃ in 56% yield and also forms red-orange crystals. Both **2a** and **2b** are slightly air sensitive, in solution and as solids, and were purified by sublimation at 60 and 50 °C, respectively, at 10⁻⁴ torr.

In contrast to the closely related complex⁶ CpFe(NO)Me₂, which slowly decomposes at room temperature, complexes **2** are remarkably thermally stable. In the absence of phosphine, dimethyl complex **2a** decomposes slowly at 120 °C over 24 h, leading to intractable products. However, thermolysis in benzene at 150 °C for 20 h in the presence of trimethylphosphine produces a new material having an intense IR absorption at 2060 cm⁻¹ and resonances in the ¹H NMR spectrum at δ 1.66 (t, 15 H, $J = 1.4$ Hz) and 1.16 (virtual triplet, 18 H, $J + J = 8.4$ Hz). This, as well as other spectral evidence, is consistent with the formulation of this material as the metal cyanide Cp^{*}Ru(PMe₃)₂(CN) (**3**, Scheme I). Compound **3** is isolated as yellow crystals in 51% yield. Water and methane are also produced in this reaction (identified by GC/MS and quantified by GC and ¹H NMR). Complex **3** may be prepared independently by treatment of Cp^{*}Ru(PMe₃)₂Cl⁷

Scheme I



with potassium cyanide in methanol.⁸

Additional insight as to how this cyanide complex may be formed is provided by thermolysis of the corresponding diethyl complex **2b** in the presence of trimethylphosphine. This reaction proceeds, under somewhat milder conditions (85 °C in benzene), with extrusion of 0.82 equiv of ethane (as determined by GC/MS and quantified by GC; 0.09 equiv of ethylene is also formed); it gives as the organometallic product a yellow crystalline material analyzing correctly for the formula C₁₈H₃₇NOP₂Ru in 76% isolated yield. In addition to resonances for one Cp^{*} and two trimethylphosphine ligands in the ¹H NMR (δ 1.55 (t, 15 H, $J = 1.4$ Hz), 1.28 (virtual triplet, 18 H, $J + J = 8.2$ Hz)), it exhibits resonances characteristic of one vinylic hydrogen and methyl group (δ 6.43 (q, 1 H), 2.17 (d, 3 H)) weakly coupled to one another ($J = 5.2$ Hz). We therefore propose a ruthenium oximate (aza enolate) (Cp^{*}L₂Ru(ONCHCH₃)) formulation for this complex. That the carbon attached to nitrogen is in fact sp² hybridized is confirmed by the ¹³C-H coupling constant of 172.7 Hz measured in the gated carbon spectrum. This requires that the ruthenium oximate has the unusual⁹ (at least for a late-transition-metal complex) oxygen-metal bound structure **4** rather than the analogous carbon-metal bound structure **5**. Thermolysis of **4** at 75 °C for 4.5 h in benzene leads (35%) to an isomeric crystalline yellow material containing an exchangeable, rather than vinylic, hydrogen, which spectral data demonstrate to be the *N*-ruthenium amide complex **6**. Confirmation of this assignment was obtained by independent synthesis of **6** from Cp^{*}Ru(PMe₃)₂Cl and CH₃CONHLi in Me₂SO. The transformation of **4** to **6** is strongly inhibited by added phosphine. This explains why **6** is not formed at the higher temperature at which **4** is prepared. Furthermore, if dimethylphenylphosphine is added instead of trimethylphosphine, phosphine substitution is observed before isomerization.¹⁰

We propose the pathways outlined in Scheme I to explain these observations. The process is presumably initiated by migratory insertion of NO into one of the metal alkyl bonds,^{6,11} leading to nitrosoalkane intermediate **7**. Transfer of a hydrogen to the metal center from the carbon adjacent to nitrogen (this can be viewed either as β -elimination cyclometallation), reductive elimination

(1) (a) Clement, R. A. U.S. Pat. 3 652 620, 1972. (b) Clement, R. A.; Klabunde, U.; Parshall, G. W. *J. Mol. Catal.* **1978**, *4*, 87.

(2) Pandey, K. K. *Coord. Chem. Rev.* **1983**, *51*, 69.

(3) Heinekey, D. M. Ph.D. Dissertation, University of Alberta, Edmonton, Alberta, 1981.

(4) Nitrosyl chloride was prepared from thionyl chloride and isoamyl nitrite according to Stryker et al.; Stryker, J. M. Ph.D. Dissertation, Columbia University, New York, 1983.

(5) (η^5 -C₅H₅)Ru(NO)Cl₂ has been prepared in a similar manner. See: Connelly, N. G. *Inorg. Chim. Acta Rev.* **1972**, *6*, 47.

(6) Seidler, M. D.; Bergman, R. G. *Organometallics* **1983**, *2*, 1897.

(7) Tilley, T. D.; Grubbs, R. H.; Bercaw, J. E. *Organometallics* **1984**, *3*, 274.

(8) Related cyanoruthenium complexes have recently been reported and their chemistry investigated: (a) Bruce, M. I.; Wallis, R. C.; *Aust. J. Chem.* **1981**, *34*, 209. (b) Baird, G. J.; Davies, S. G. *J. Organomet. Chem.* **1984**, *262*, 215.

(9) King, R. B.; Chen, K. N.; *Inorg. Chem.* **1977**, *16*, 1164.

(10) For an example of amide complexes in the platinum series, see: Appleton, T. G.; Bennett, M. A. *J. Organomet. Chem.* **1980**, *199*, 119.

(11) (a) Weiner, W. P.; White, M. A.; Bergman, R. G. *J. Am. Chem. Soc.* **1981**, *103*, 3612. (b) Weiner, W. P.; Bergman, R. G. *Ibid.* **1983**, *105*, 3922.

of ethane, and trapping by phosphine leads to η^3 -oximate intermediate **8**. Reaction with an additional mole of phosphine may now force conversion of the oximate ligand to one of its η^1 isomers **4** or **5**; apparently formation of **4** is favored, and this complex is isolated.

Formation of the cyanide and *N*-ruthenium amide complexes **3** and **6** is more complex, and less well-precedented, processes. We suggest these reactions may be related by the mechanism outlined at the bottom of Scheme I. Reversible interconversion of O- and C-bound oximate complexes **4** and **5** could occur thermally on phosphine loss in **4**. As **5** is simply a ruthenium-substituted nitrosoalkane, it should undergo rapid prototropic isomerization to the corresponding oxime **9**, as is typical of this class of compounds. When R = H, elimination of water in this complex leads to isolated ruthenium cyanide **6**. When R = Me, this elimination is prevented, and we suggest that the ruthenium analogue of a Beckmann rearrangement converts oxime **9** to **10**. Complex **10** should quickly rearrange to its more stable, isolable carboxamide tautomer **6**.

Experiments are currently under way aimed at testing these mechanistic hypotheses and exploring further the chemistry of the O- and N-bound ruthenium(II) derivatives reported here.¹²

Acknowledgment. We are grateful for financial support of this work by the National Institutes of Health (Grant GM25459). We also appreciate helpful discussions with Prof. J. E. Bercaw and Rocco Paciello and their willingness to disclose data prior to publication. M.D.S. acknowledges a National Science Foundation Predoctoral Fellowship; R.G.B. acknowledges a Sherman Fairchild Scholarship from the California Institute of Technology (Jan-June, 1984).

Registry No. **1**, 90419-99-1; **2a**, 91993-47-4; **2b**, 91993-48-5; **3**, 91993-49-6; **4**, 91993-50-9; **6**, 91993-51-0; $\text{Cp}^*\text{Ru}(\text{CO})_2\text{Cl}$, 77488-50-7; AlMe_3 , 75-24-1; AlEt_3 , 97-93-8; NOCl , 2696-92-6.

Supplementary Material Available: Proton, carbon, and phosphorus NMR, infrared, mass spectra, elemental analysis, and melting point data for **1**, **2a**, **2b**, **3**, **4**, and **6** (4 pages). Ordering information is given on any current masthead page.

(12) $\text{Cp}^*\text{Ru}(\text{NO})\text{Cl}$, has recently been prepared independently: Efraty, A. *J. Organomet. Chem.* **1984**, 260, 321.

Stable Potassium Complex of a Pyridine-Fused Hexaaza[18]annulene. A New Rival for 18-Crown-6

Thomas W. Bell* and Frieda Guzzo


Department of Chemistry
State University of New York at Stony Brook
Stony Brook, New York 11794-3400

Received May 29, 1984

The crown ethers¹ are of increasing importance as complexing agents for the biologically relevant alkali metal ions, potassium and sodium. In recent years, considerable synthetic effort has been directed toward surpassing the ion affinities of crown ethers, notable achievements being cryptands² and spherands.³ Most simple structural modifications of 18-crown-6, such as benzanellation and heteroatom replacement, lead to decreased potassium complex stability.^{4,5} We report here a potassium complex of a nitrogen analogue of 18-crown-6 having a stability constant 1 order


of magnitude greater than that of 18-crown-6.

The dominant interaction between alkali metal ions and oxygen or nitrogen ligand sites is generally considered to be an ion/dipole electrostatic attraction.⁶ Reduced stability constants observed in saturated nitrogen analogues⁴ of crown ethers are consistent with this analysis, since the dipole moment of trimethylamine (0.61 D) is smaller than that of dimethyl ether (1.30 D).⁷ The sizeable dipole moment of (methylimino)methane (1.53 D) suggests the hexaaza[18]annulene **1**⁸ as the parent for a new family of alka-



li-metal complexing agents. Fusion of **1** with pyridine rings (2.19 D) should afford even better macrocyclic ligands, assuming a planar conformation. The dibenzo pyridido hexaaza[18]annulene **2**⁹ has been found in complexes with alkaline-earth and some transition-metal ions,¹⁰ although the free ligand has not been reported. An earlier report¹¹ of a metal-free synthesis of the tetramethyl derivative of **2** has been found in error.^{12,13}

Alkaline earth perchlorate complexes of tetramine **2** may be prepared by the metal-templated condensation¹⁰ shown in eq 1.



We have found that when the strontium complex $2 \cdot \text{Sr}(\text{ClO}_4)_2$ ¹⁰ is treated with potassium fluoride monohydrate in refluxing methanol, the corresponding potassium perchlorate complex is obtained. Recrystallization from acetonitrile afforded analytically pure (C, H, N, K) yellow needles (69%, mp 386–388 °C dec), exhibiting infrared absorptions¹⁴ consistent with the proposed structure ($2 \cdot \text{KClO}_4$). The proton NMR chemical shifts of this material in $\text{Me}_2\text{SO}-d_6$ are given in Table I (entry 3) in comparison with the chemical shifts of $2 \cdot \text{Sr}(\text{ClO}_4)_2$ ¹⁰ and the 2:1 sandwich complex of **2** with barium perchlorate¹⁰ (entries 1 and 2).

The chemical shifts of $2 \cdot \text{KClO}_4$ were found to be dependent on concentration and on the presence of added potassium complexing agents (Table I). The imine methine resonance (Hm), which occurs at δ 8.887 for a 2×10^{-2} M solution (entry 3), shifts to lower field with dilution, e.g., 8.962 ppm at 4×10^{-3} M (entry 6). The effect of added [2.2.2]cryptand is also dependent on the initial concentration of $2 \cdot \text{KClO}_4$. Thus, incremental addition of

(6) Pullman, A. In "Physical Chemistry of Transmembrane Ion Transport"; Spach, G., Ed.; Elsevier: New York, 1983; pp 181–206. See also: Izatt, R. M.; Christensen, J. J., Eds. "Progress in Macrocyclic Chemistry"; Wiley: New York, 1979; Vol. 1. (Especially Morf, W. E.; Amman, D.; Bissig, R.; Pretsch, E.; Simon, W., pp 1–61; Poonia, N. S., pp 115–155.)

(7) Nelson, R. D., Jr.; Lide, D. R.; Maryott, A. A. "Selected Values of Electric Dipole Moments for Molecules in the Gas Phase"; National Bureau of Standards: Washington, DC, NSRDS-NBS 10.

(8) Systematic name for **1**: 1,4,7,10,13,16-hexaazacyclooctadecine.

(9) Systematic name for **2**: 4,5,15,16-dibenzo-3,6,14,17,23,24-hexaazatricyclo[17.3.1.1^{8,12}]tetracos-1(23),4,8(24),9,11,15,19,21-octaene.

(10) Drew, M. G. B.; Cabral, J. de O.; Cabral, M. F.; Esho, F. S.; Nelson, S. M. *J. Chem. Soc., Chem. Commun.* **1979**, 1033–1035. Sodium and potassium complexes of the furan analogue of **2** were recently reported: Nelson, S. M.; Esho, F. S.; Drew, M. G. B. *J. Chem. Soc., Dalton Trans.* **1983**, 1857–1864.

(11) Stotz, R. W.; Stouffer, R. C. *J. Chem. Soc., Chem. Commun.* **1970**, 1682–1683.

(12) Cabral, J. de O.; Cabral, M. F.; Drew, M. G. B.; Esho, F. S.; Haas, O.; Nelson, S. M. *Ibid.* **1982**, 1066–1067.

(13) The reported failure of the tetramethyl derivative of **2** to extract alkali metal ions must also be disregarded: Ramdani, A.; Tarrago, G. *Tetrahedron* **1981**, 37, 991–1000.

(14) IR (KBr) 3050 (w), 3000 (w), 1620 (m), 1580 (s), 1480 (m), 1440 (m), 1365 (w), 1255 (m), 1080 (br, s), 800 (m), 725 (m) cm^{-1} .

(15) An equimolar ratio of the cryptand to **2** could not be obtained for 2×10^{-2} M solutions of $2 \cdot \text{KClO}_4$, due to low solubility.

(1) Pedersen, C. J. *J. Am. Chem. Soc.* **1967**, 89, 2495–2496. See also: Hiraoka, M. "Crown Compounds"; Elsevier: New York, 1982.

(2) Dietrich, B.; Lehn, J. M.; Sauvage, J. P. *Tetrahedron Lett.* **1969**, 2885–2888. Review: Lehn, J. M. *Pure Appl. Chem.* **1980**, 52, 2303–2319.

(3) Cram, D. J.; Kaneda, T.; Helgeson, R. C.; Lein, G. M. *J. Am. Chem. Soc.* **1979**, 101, 6752–6754. Review: Cram, D. J. *Science (Washington, D.C.)* **1983**, 219, 1177–1183.

(4) Frensdorff, H. K. *J. Am. Chem. Soc.* **1971**, 93, 600–606.

(5) Review: De Jong, F.; Reinhoudt, D. N. *Adv. Phys. Org. Chem.* **1981**, 17, 279–433.