

KINETICS OF THERMAL [1,7A]-SIGMATROPIC SHIFT OF HEXAFLUORO VITAMIN D₃ AND VITAMIN D₃ DERIVATIVES. EVALUATION OF CONFORMATIONS OF THE A RING AFFECTED BY 1-OH AND 3-OH GROUPS.

Jun-etsu Igarashi, Masahiko Ikeda, and Makoto Sunagawa*

*Development Research Laboratories I, Sumitomo Pharmaceuticals Research Center, 3-1-98 Kasugadenaka,
Konohanaku, Osaka 554, Japan.*

Abstract: The quantitative evaluation of the [1,7a]-sigmatropic rearrangement of vitamin D₃ and its analogs affected by the conformations of the A ring using the ¹H-NMR method was described. Although the side chain of the D ring had no effect on the hydrogen migration, the rearrangement was influenced by the hydroxy groups of the A ring. Copyright © 1996 Elsevier Science Ltd

It is well known that a key step in the primary metabolic pathway leading to physiologically active 1 α ,25-dihydroxyvitamin D₃ (1,25-(OH)₂D₃)¹⁾ is the transformation of previtamin D₃ (preD₃) to vitamin D₃ (D₃) (Figure 1). The kinetics and thermodynamics of this isomerization for the specific case of preD₃ has been studied in detail by Hanewald et al.^{2a)} and others.^{2b,c)} The equilibrium ratio of previtamin to vitamin is temperature dependent and the reaction follows reversible and first-order kinetics. Extensive work has been completed to evaluate the conformational features of preD₃ and D₃,³⁾ however, little attention has been paid to the relationship between the conformation of the A ring affected by the 1- and 3-OH groups and the kinetic study under the same reaction conditions. ST-630 (26,26,26,27,27,27-hexafluoro-1,25-dihydroxyvitamin D₃) is one of a few analogs that have greater biological activity than 1,25-(OH)₂D₃ *in vivo* in the vitamin D-deficient rat⁴⁾ and chick,⁵⁾ and its action is longer-lasting than that of 1,25(OH)₂D₃. ST-232 (26,26,26,27,27,27-hexafluoro-1,23(S),25-tri-hydroxyvitamin D₃) is the major metabolite of ST-630.⁶⁾ The reason for the enhanced biological activities⁴⁾ was explained in part by a decreased metabolic inactivation by way of 26- and 27-hydroxylations due to the

substituted fluoro groups at the 26- and 27-carbons.




Figure 1. Molecular formulas of hexafluorovitamin D₃ and vitamin D₃ derivatives with the numbering system used in this paper

It is the purpose of this article to describe the quantitative evaluation of the rearrangement of hexafluorovitamin D₃ and vitamin D₃ derivatives. This would provide the information about the effects of the hydroxyl substituents of the A ring, which relates the conformation of the A ring, during the [1,7a]-sigmatropic hydrogen shift.

The ¹H-NMR experiments were done using a JEOL A-500 (500 MHz) nmr spectrometer operating in the pulsed fourier transform mode with a DEC station 3200 computer and 32 K data points. Crystalline D₃ and 1 α -hydroxycholecalciferol (1-OH-D₃) were purchased from Solvay Duphar (Amsterdam, the Netherlands) and used without further purification. ST-630, ST-232 and 3-deoxyvitamin D₃ (3-deoxy-D₃) were synthesized⁷⁾ and then stored in the dark below -20 °C. Spectroscopic grade ethanol-d₆, which was obtained from E. Merck (Darmstadt, Germany), was used as the solvent with TMS serving as the internal reference. For the kinetic studies, solutions of ST-630, ST-232, D₃, 1-OH-D₃ and 3-deoxy-D₃ were dissolved in ethanol-d₆ such that the final concentration was approximately 5 mg/ml. These solutions were then cooled to -78 °C.

For a kinetic run, a sample was placed in the precalibrated ¹H-NMR probe, which was preset to a specific temperature. After thermal equilibration of the sample, the ¹H-NMR spectra were recorded at regular time intervals. For the isomerization of ST-630, ST-232, D₃, 1-OH-D₃ and 3-deoxy-D₃ to the thermodynamically less stable previtamin form, the rate of the reaction was monitored by following the disappearance of the H-6 signal (ST-630: δ 6.27, ST-232: δ 6.26, D₃: δ 6.25, 1-OH-D₃: δ 6.27, 3-deoxy-D₃: δ 6.16). As a cross check, the H-7 signal (ST-630: δ 6.08, D₃: δ 6.03, ST-232: δ 6.08, 1-OH-D₃: δ 6.09, 3-deoxy-D₃: δ 6.04) or the H-19 Z signal (ST-630: δ 5.29, ST-232: δ 5.27, D₃: δ 5.05, 1-OH-D₃: δ 5.27, 3-deoxy-D₃: δ 4.99) was also periodically monitored to measure the reliability of the integration data. Measurements were made at three different temperatures (45 °C, 60 °C, 75 °C).

The reversible first-order rate constants of the reaction between the previtamin form and vitamin form are defined^{2a)} by $\ln m/(m-x) = (k_1 + k_{-1})t$, where $m = (k_1 a - k_{-1} b) / (k_1 + k_{-1})$, and a and b are the concentrations of the previtamin form at t=0, respectively. x is the change in concentration. To calculate m, $m = (a - Kb) / (1 + K)$ is needed, where K is the equilibrium constant (k_1/k_{-1}). K, m, and x are obtained from the NMR results. In a plot of $\ln m/(m-x)$ versus time (s), the slope of the line is the sum of k_1 and k_{-1} . Thus k_1 and k_{-1} are obtained. The activation parameters were calculated from an Arrhenius plot of the natural logarithm of the rate constants for the previtamin form to vitamin form conversion (k_1) versus the reciprocal of the absolute temperature. A kinetic study of their [1,7a]-sigmatropic hydrogen shifts was carried out according to the ¹H-NMR analytical method developed by Okamura et al.⁸⁾ Integration of the H-6 or H-7 signal of the previtamin versus the H-6 or H-7 signal of the vitamin in the ¹H-NMR spectrum could be used to quantify the relative amounts of previtamin and vitamin. Assuming a reversible, first-order kinetic rate law and following the reaction to 7-15 half-lives, with separate determination of the equilibrium constants for the preD₃-D₃ interconversion over the same temperature range, the results summarized in Tables I and II were obtained.

The kinetic data and activation parameters for the transformation of the D₃ at 80 °C were essentially identical to the value previously reported by Hanewald et al.^{2a)} The rate constant for the [1,7a]-hydrogen migration of 1-OH-D₃ at 80 °C was calculated to be $5.65 \times 10^{-4} \text{ s}^{-1}$, which is comparable to the $5.63 \times 10^{-4} \text{ s}^{-1}$ value calculated for the isomerization of 1, 25-(OH)₂D₃.⁹⁾ The rate constants and activation parameters for the isomerization of ST-630 most closely resembled those of ST-232 and 1-OH-D₃ but different from those of D₃,

especially K_{eq} and the enthalpy of activation. Moreover, K_{eq} and the enthalpy of activation of D_3 were different from 3-deoxy- D_3 .

Table I. Kinetic Data for the Transformation of Vitamin D to Previtamin D^a

| Substrate | $k_1^b \times 10^4$ | $k_{-1}^b \times 10^4$ | K_{eq}^c |
|---------------------------------|---------------------|------------------------|-------------|
| 3-Deoxy-vitamin D_3 | 4.83 (0.11) | 1.58 (0.17) | 3.06 (0.31) |
| Vitamin D_3 | 5.04 (0.19) | 1.36 (0.23) | 3.71 (0.25) |
| 1- α -OH-Cholecalciferol | 5.65 (0.28) | 1.09 (0.13) | 5.18 (0.18) |
| ST-232 | 5.46 (0.33) | 1.07 (0.29) | 5.10 (0.35) |
| ST-630 | 5.46 (0.25) | 1.07 (0.22) | 5.10 (0.26) |

^a At 80 °C. Standard deviations are given in parentheses.

^b k_1 and k_{-1} is given in s^{-1} . ^c K_{eq} is defined as k_1/k_{-1} where the forward process is for the isomerization of previtamin D to vitamin D.

Table II. Activation Parameters for the Transformation of Vitamin D to Previtamin D^a

| Substrate | E_a^b | $\log A^c$ | ΔG^\ddagger^b | ΔH^\ddagger^b | ΔS^\ddagger^d |
|---------------------------------|------------|-------------|-----------------------|-----------------------|-----------------------|
| 3-Deoxy-vitamin D_3 | 23.5 (0.2) | 10.8 (0.1) | 26.9 (0.5) | 21.8 (0.2) | -14.4 (0.1) |
| Vitamin D_3 | 23.6 (0.1) | 10.7 (0.2) | 27.0 (0.6) | 22.9 (0.1) | -11.8 (0.4) |
| 1- α -OH-cholecalciferol | 24.5 (0.4) | 11.3 (0.09) | 27.2 (0.8) | 23.9 (0.4) | -9.3 (0.4) |
| ST-232 | 24.5 (0.1) | 11.2 (0.1) | 27.2 (0.7) | 23.8 (0.1) | -9.5 (0.3) |
| ST-630 | 24.5 (0.5) | 11.2 (0.3) | 27.2 (0.5) | 23.8 (0.3) | -9.5 (0.2) |

^a At 80 °C. Standard deviations are given in parentheses. ^b Units=kcal/mol. ^c A is given in s^{-1} .

^d Units=cal/mol K.

These data indicated that the 1- and 3-OH groups of the A-ring affected the [1,7a]-sigmatropic hydrogen migration and the nature of the side chain of the D-ring had essentially no effect on the isomerization.

These results agreed with the mechanism of the [1,7a]-sigmatropic migration because the A-, seco-B-, C- and D-ring are associated with the transition states of the isomerization and the side chain of the D-ring is far apart from the structure. The fact that the K_{eq} (or the free energy of activation) and the enthalpy of activation of ST-630, D₃, 3-deoxy-D₃ were decreased slightly in this order meant that the vitamin form of 3-deoxy-D₃ was more easily converted to its pre-form than ST-630 and D₃. No significant changes in the entropy of activation for this transformation were observed among ST-630, ST-232, D₃, 1-OH-D₃, and 3-deoxy-D₃.

A conformational equilibrium of the A ring between two chair forms must be occurring. Wing^{3a,b)} reported that the introduction of a hydroxyl group at 1 α in D₃ slightly shifts the conformational equilibrium to favor an equatorial 1 α -OH-D₃. We thought that the A ring of 3-deoxy-D₃ consists of an equilibrium mixture of almost an equal population of the two chair conformers, because it has no hydroxy groups at the A Ring. The [1,7a]-sigmatropic rearrangement would be influenced by the conformation of the A ring. We concluded that the shifts in the conformational equilibrium between the two chair forms of the A ring resulted in the difference of the equilibrium constants and the enthalpy of activation among the active type of D₃, D₃, and 3-deoxy-D₃. Although the presence of systematic experimental error can not be ruled out, the equilibrium constant and the enthalpy of activation of D₃ is bigger than 3-deoxy-D₃ and smaller than the active type of D₃ under the same reaction conditions. It was suggested that these data imply that the origin of the observed difference in the rate constant lies in a relative population of the active type of D₃, D₃, and 3-deoxy-D₃.

References

- 1) a) M. F. Holick, J. E. Frommer, S. C. McNeill, N. M. Richtand, J. W. Henley, J. T. Potts., Jr., *Biochem. Biophys. Res. Commun.*, **76**, 107 (1977). b) M. F. Holick, J. A. MacLaughlin, M. B. Clerk, S. A. Holick, J. T. Potts., Jr., R. R. Anderson, I. H. Blank, J. A. Parish, P. Elias, *Science*, **210**, 203 (1980). c) M. F. Holick, *J. Invest. Dermatol.*, **76**, 51 (1981).
- 2) a) K. H. Hanewald, M. P. Rappoldt, J. R. Roborgh, *Recl. Trav. Chim. Pays-Bas*, **80**, 1003 (1961). b) J. K. Yamamoto, R. F. Borch, *Biochemistry*, **24**, 3338 (1985). c) E. G. Cassis., Jr., R. G. Weiss,

Photochem. Photobiol. **35**, 439 (1982).

- 3) a) R. M. Wing, W. H. Okamura, A. Rego, M. R. Pirio, A. W. Norman, *J. Am. Chem. Soc.*, **97**, 4980 (1975). b) R. M. Wing, W. H. Okamura, M. R. Pirio, S. M. Sine, A. W. Norman, *Science*, **186**, 939 (1974). c) W. G. Dauben, D. J. H. Funhoff, *J. Org. Chem.*, **53**, 5070 (1988). d) W. G. Dauben, D. J. H. Funhoff, *ibid.*, **53**, 5376 (1988). e) V. Delafoff, P. Rathle, M. Legrand, *Bull. Soc. Chim. Fr.*, **1963** (1739). f) G. N. La Mar, D. L. Budd, *J. Am. Chem. Soc.*, **96**, 7317 (1974).
- 4) Y. Tanaka, H. F. DeLuca, Y. Kobayashi, N. Ikekawa, *Arch. Biochem. Biophys.*, **229**, 348 (1984).
- 5) T. Kiriya, S. Okamoto, H. Suzuki, A. Nagata, M. Izumi, H. Morii, S. Nagataki, *Acta Endocrinol.*, **121**, 520 (1989).
- 6) A. Nagata, N. Hamma, T. Katsumata, A. W. Norman (ed) *Vitamin D Molecular, Cellular and Clinical Endocrinology*. Water de Gruyter, Berlin, New York, pp 141-142 (1988).
- 7) a) B. Lythgoe, T. A. Moran, M. E. N. Nambudiry, J. Tindeswell, and P. W. Wright, J. C. S. Parkin I, **1978** (590). b) Y. Kobayashi, T. Taguchi, S. Mitsunashi, T. Eguchi, E. Ohshima, and N. Ikekawa, *Chem. Pharm. Bull.*, **30**, 4297 (1982). c) B. M. Trost, J. Dumas, and M. Villa, *J. Am. Chem. Soc.*, **114**, 9836 (1992). d) J. P. Sestelo, J. L. Mascarenas, L. Castedo, and A. Mouriro, *J. Org. Chem.*, **58**, 118 (1993).
- 8) M. L. Curtin, W. H. Okamura, *J. Org. Chem.*, **55**, 5278 (1990).
- 9) a) M. L. Curtin, W. H. Okamura, *J. Am. Chem. Soc.*, **113**, 6958 (1991). b) D. F. Maynard, W. G. Trankle, A. W. Norman, W. H. Okamura, *J. Med. Chem.*, **37**, 2387 (1994).

(Received in Japan 3 April 1996; accepted 17 May 1996)