

185. *SPELEANDS*. Macropolycyclic Receptor Cages Based on Binding and Shaping Sub-units. Synthesis and Properties of Macrocyclic-Cyclotrimeratrylene Combinations

Preliminary communication

by Josette Canceill^{a)}, André Collet^{a)}, Jacqueline Gabard^{a)}, Florence Kotzyba-Hibert^{b)} and Jean-Marie Lehn^{a, b)}

^{a)} Laboratoire de Chimie des Interactions Moléculaires, Collège de France, 11, Place Marcelin-Berthelot, F-75005 Paris¹⁾

^{b)} Institut Le Bel, Université Louis-Pasteur, 4, rue Blaise-Pascal, F-67000 Strasbourg²⁾

(16. VII. 82)

Summary

Combination of a receptor unit with a rigid shaping unit produces a new type of receptor molecules of the cryptand class, hollow macropolycyclic molecules termed *speleands*, capable of substrate inclusion. Two members of this category of compounds **1** and **2**, have been synthesized by connecting in a single step, a macrocyclic [18]-N₃O₃ binding unit with a rigid cyclotrimeratrylene unit *via* three bridges. Compound **1** binds the methylammonium cation forming both external and internal complexes; for the latter a '*speleate*' structure, schematically represented by **15**, may be proposed.

Cryptates are defined as inclusion complexes in which the substrate species is (are) contained in the intramolecular cavity of a macropolycyclic receptor molecule. Introduced originally for alkali and alkaline-earth complexes of macrobicyclic cryptands, this general concept has since been extended to inclusion complexes of a large variety of substrates (other inorganic cations, organic cations, as well as anionic species) with macropolycyclic molecules of various shapes and sizes, designed so as to display molecular recognition processes [1]. The molecular architecture of these macropolycycles is based on the combination of various sub-units possessing specific structural properties and containing defined binding sites, in order to achieve receptor-substrate complementarity for both geometry and interactions. Furthermore, ligand dynamics, *i.e.* the rigidity/flexibility balance, influence substrate binding and exchange rates.

Within this general class of cryptand molecules, the connection of polar *binding sub-units* to large, concave, more or less rigid and hydrophobic *shaping*

¹⁾ GR 20 of the CNRS.

²⁾ ERA 265 of the CNRS.

components, should produce a new type of hollow macropolycyclic receptors, capable of substrate inclusion. We propose the terms *speleands* (from $\sigma\pi\eta\lambda\alpha\iota\omicron\nu$ = cave) and *speleates* for individualizing this sub-class of cryptands and of inclusion complexes. The binding sub-units may be derived from chelating, tripodal or macrocyclic arrangements of binding sites (as for instance in polynucleating cryptands [2]). The shaping components may belong to different types of framework like phenol-aldehyde condensation products, cyclodextrins [3] or functionalized porphyrins and tetrapyrroles [4]. The former are of special interest since, in addition to known representatives (e.g. veratrole-formaldehyde cyclotrimeratrylene [5–10], phenol-formaldehyde [11] [12] calixarene [13] [14] or resorcinol-aldehyde [15–17] condensation products), many intriguing variations remain to be investigated. Furthermore, such substances form clathrate inclusion compounds with organic molecules in the solid state (cf. [14] [18]) a property which may confer additional binding potential to speleands, especially towards hydrophobic substrates.

We now report preliminary results on the synthesis and substrate-binding properties of two members of this family of macropolycycles (compounds **1** and **2**) which result from the connection of a macrocyclic binding sub-unit of type [18]-N₃O₃ (**3**, [19]) to the cyclotrimeratrylene (CTV) derivative **4** [9] [20] by three bridges. Triply bridged coreceptors incorporating two [18]-N₃O₃ rings and forming molecular cryptates have been described [21], as well as a molecular cage formed by two triply connected CTV units [20].


The racemic (C₃)-tricarboxylic acid ($\pm$) **5** [20] was converted into the trichloride **6** by treatment with thionyl chloride. Condensation of **6** with **3** in high dilution conditions [22] afforded the tricarboxamide **7** (m.p. $\approx 325^\circ$ (dec.)) in 35% yield after purification by chromatography over alumina. Reduction of **7** with diborane [22], gave the CTV-[18]-N₃O₃ molecular cage **1** (m.p. $\approx 230^\circ$) in 85% yield.

The CTV unit bearing longer branches was prepared as follows. Treatment of the sodium salt of vanillin with ICH₂CH₂OCH₂CO₂CH₃ (for the corresponding chloro-analog see [23]) gave **8** (m.p. 85° ; 85%) which was reduced to the alcohol **9** (m.p. 40° ; quantitative yield) with H₂/Raney nickel. Trimerization of **9** with perchloric acid in methyl orthoformate solution at room temperature for 20 h gave the CTV trimer **10** in 40% yield (m.p.: 98°). The crude triacid **11**, obtained as a glassy solid by basic hydrolysis of **10** (NaOH in methanol/water 95:5), was converted to the corresponding trichloride **12** by treatment with thionyl chloride. Following the same reaction sequence as for **1**, **12** was condensed with **3** to give the tricarboxamide **13** (glass; 40%), which was reduced to the larger CTV-[18]-N₃O₃ cage **2** (glass; 65%). All new compounds had physical properties in agreement with the proposed structures.


Compounds **1** and **2** represent a new type of macropolycyclic mesomolecules [1], *speleands*, containing both a receptor site and a rigid shaping unit. They combine the features of bis-[18]-N₃O₃ cylindrical macrotetracycles [21] and of the bis-CTV molecular cage [20].

The overall shape of **1** and **2** may be described as a circular component topped by the CTV unit in its usual 'crown'-type conformation (as shown by the characteristic AB-quadruplet of the methylene bridges at about 4.7 and 3.5 ppm, $J = 14$ Hz in the ¹H-NMR. spectrum [9] [20]).


Compounds **1** and **2** are *hollow molecules*, whose intramolecular cavity is maintained by the rigid CTV unit [9]. The flexible bridges permit adjustment of the cavity height along the C_3 -axis; the maximum size is clearly larger for **2** ($\approx 8 \text{ \AA}$) than for **1** ($\approx 5 \text{ \AA}$).




^{a)} The three $N\text{---}O\text{---}O$ bridges between the CTV unit and [18]- N_3O_3 stand for $N\text{---}COCH_2O$.



3 $R = H$
14 $R = CH_3$



4 $R = H$
5 $R = CH_2COOH$
6 $R = CH_2COCl$
10 $R = CH_2CH_2OCH_2COOCH_3$
11 $R = CH_2CH_2OCH_2COOH$
12 $R = CH_2CH_2OCH_2COCl$



$R = CH_2CH_2OCH_2COOCH_3$

8 $X = CHO$
9 $X = CH_2OH$

The [18]- N_3O_3 macrocyclic sub-unit of **1** and **2** may serve as receptor site for cationic species. Indeed, the tris(*N*-methyl) derivative **14** binds metal cations and is especially well-suited for selective complexation of primary ammonium cations [19]. The latter property is of particular interest in the present case, since binding of $R\text{---}NH_3^+$ by anchoring of the NH_3^+ head group into the [18]- N_3O_3 ring, may occur either inside or outside the central cavity depending on the nature of the *R* residue. Preliminary experiments have shown that **1** dissolves about 1 equivalent $CH_3NH_3^+$ picrate into $CDCl_3/CD_3OD$ 9:1 or into CD_2Cl_2 yielding a mixture (about 2:1) of two species giving $CH_3\text{---}NH_3^+$ NMR. signals around 2.3 ppm and -0.1 ppm (ill-resolved quadruplet, $J \approx 6 \text{ Hz}$ in CD_2Cl_2). They may be identified respectively as the *exo*- and the *endo*-complexes, if we accept that the large shift to high field is diagnostic of inside binding, thus placing the CH_3 group of the substrate into the shielding region of the aromatic rings, as observed for the diammonium cryptates of cylindrical coreceptors [21] [24]. A schematic representation of this cryptate-type inclusion complex, the *methylammonium speleate* of **1**, is given by structure **15**. $R\text{---}NH_3^+$ substrates with *R* larger than CH_3 apparently do not bind inside (experi-

ments in progress). Double-irradiation $^1\text{H-NMR}$ experiments indicate that *endo*- and *exo*-bound CH_3NH_3^+ species exchange slowly (saturation transfer), and that the protons of *exo*- and *endo*- NH_3^+ groups undergo respectively fast and slow exchange with the solvent. The binding properties of **2** are being investigated together with further studies of **1**, in particular by low-temperature NMR.

Developments of the chemistry of speleands may be concerned with: *i*) the incorporation of rigid rather than flexible bridges, as in **1** and **2**, or rigidification of the basic components by structural modifications (for a recent example on calixarenes see [25]), in order to enforce more strongly a preformed cavity; *ii*) the synthesis of optically active **1** and **2** since the basic CTV unit is chiral and has been resolved [9]; *iii*) the introduction of polar groups in order to confer solubility in water and to permit operation of hydrophobic effects. In a broader view, a great variety of other combinations of receptor and architectural units may be envisaged, as briefly mentioned above, adding another facet to the chemistry of molecular receptors.

REFERENCES

- [1] J. M. Lehn, *Structure and Bonding* 16, 1 (1973); *Acc. Chem. Res.* 11, 49 (1978); *Pure Appl. Chem.* 50, 871 (1978).
- [2] J. M. Lehn, *Pure Appl. Chem.* 52, 2441 (1980); in 'IUPAC - Frontiers of Chemical Research', Ed. K. J. Laidler, Pergamon Press, Oxford 1982, p. 265.
- [3] M. L. Bender & M. Komiyama, 'Cyclodextrin Chemistry', Springer-Verlag, Berlin 1978.
- [4] J. C. Lalloz, J. M. Lehn & A. K. Willard, unpublished results.
- [5] H. Erdman, F. Haglid & R. Ryhage, *Acta Chem. Scand.* 18, 1249 (1964).
- [6] A. S. Linsey, *J. Chem. Soc.* 1965, 1685.
- [7] R. C. Cookson, B. Halton & I. D. R. Stevens, *J. Chem. Soc. B* 1968, 767.
- [8] J. A. Hyatt, *J. Org. Chem.* 43, 1808 (1978); K. Frensch & F. Vögtle, *Justus Liebigs Ann. Chem.* 1979, 2121.
- [9] A. Collet & J. Jacques, *Tetrahedron Lett.* 1978, 1265; A. Collet & J. Gabard, *J. Org. Chem.* 45, 5400 (1980); A. Collet, J. Gabard, J. Jacques, M. Cesario, J. Guilhem & C. Pascard, *J. Chem. Soc., Perkin I* 1981, 1630.
- [10] F. M. Menger, M. Takeshita & J. F. Chow, *J. Am. Chem. Soc.* 103, 5938 (1981).
- [11] A. Zinke, R. Kretz, E. Leggewic & K. Hössinger, *Monatsh. Chem.* 83, 1213 (1952).
- [12] J. W. Cornforth, E. D. Morgan, K. T. Potts & R. J. W. Rees, *Tetrahedron* 29, 1659 (1973); J. W. Cornforth, P. D'Arcy Hart, G. A. Nicholls, R. J. W. Rees & J. A. Stock, *Brit. J. Pharmacol.* 10, 73 (1955).
- [13] C. D. Gutsche, B. Dhawan, K. H. No & R. Muthukrishnan, *J. Am. Chem. Soc.* 103, 3782 (1981) and ref. therein.
- [14] G. D. Andretti, R. Ungaro & A. Pochini, *J. Chem. Soc., Chem. Commun.* 1981, 533.
- [15] H. Erdman & S. Högberg, *Tetrahedron Lett.* 1968, 1679 and ref. therein.
- [16] A. G. S. Högberg, *J. Am. Chem. Soc.* 102, 6046 (1980).
- [17] J. B. Niederl & H. J. Vogel, *Ber. Dtsch. Chem. Ges.* 62, 2512 (1940).
- [18] V. Gaglioti, A. M. Lignori, N. Galo, E. Giglio & M. Scrocco, *J. Inorg. Nucl. Chem.* 8, 572 (1958).
- [19] J. M. Lehn & P. Vierling, *Tetrahedron Lett.* 1980, 1323.
- [20] J. Gabard & A. Collet, *J. Chem. Soc., Chem. Commun.* 1981, 1137.
- [21] F. Kotzyba-Hibert, J. M. Lehn & K. Saigo, *J. Am. Chem. Soc.* 103, 4266 (1981).
- [22] B. Dietrich, J. M. Lehn, J. P. Sauvage & J. Blanzat, *Tetrahedron* 29, 1629 (1973).
- [23] E. J. Salmi, R. Leimu & H. Kallio, *Suomen Kem. Lehti* 17B, 17 (1944); [*Chem. Abstr.* 40, 6491 (1946)].
- [24] F. Kotzyba-Hibert, J. M. Lehn & P. Vierling, *Tetrahedron Lett.* 1980, 941.
- [25] C. D. Gutsche & J. A. Levine, *J. Am. Chem. Soc.* 104, 2652 (1982).