A Regio- and Stereoselective ω-Transaminase/Monoamine Oxidase Cascade for the Synthesis of Chiral 2,5-Disubstituted Pyrrolidines**

Elaine O'Reilly, Cesar Iglesias, Diego Ghislieri, Jennifer Hopwood, James L. Galman, Richard C. Lloyd, and Nicholas J. Turner*

Abstract: Biocatalytic approaches to the synthesis of optically pure chiral amines, starting from simple achiral building blocks, are highly desirable because such motifs are present in a wide variety of important natural products and pharmaceutical compounds. Herein, a novel one-pot ω -transaminase (TA)/ monoamine oxidase (MAO-N) cascade process for the synthesis of chiral 2,5-disubstituted pyrrolidines is reported. The reactions proceeded with excellent enantio- and diastereoselectivity (>94% ee; >98% de) and can be performed on a preparative scale. This methodology exploits the complementary regio- and stereoselectivity displayed by both enzymes, which ensures that the stereogenic center established by the transaminase is not affected by the monoamine oxidase, and highlights the potential of this multienzyme cascade for the efficient synthesis of chiral building blocks.

he exquisite chemo-, regio- and stereoselectivity displayed by enzymes has led to their widespread application as catalysts for stereocontrolled organic synthesis.^[1] These properties, coupled with their ability to catalyze reactions under similar conditions, has enabled the development of elegant multienzyme cascade processes, in which the product formed by the action of the first enzyme becomes the starting material for the subsequent biotransformation.^[2] Such tandem processes alleviate the need for protecting-group manipulations and intermediate purification steps, thus providing cost-effective routes to target molecules.

Among the most synthetically useful biocatalysts for the synthesis of chiral amines are the ω -transaminase (TA) family and variants of monoamine oxidase from *Aspergillus niger* (MAO-N).^[3] TAs are capable of mediating the selective

[*] Dr. E. O'Reilly, Dr. D. Ghislieri, Dr. J. Hopwood, Dr. J. L. Galman, Prof. N. J. Turner
School of Chemistry, University of Manchester
Manchester Institute of Biotechnology
131 Princess Street, Manchester, M1 7DN (UK)
E-mail: nicholas.turner@manchester.ac.uk
C. Iglesias
Facultad de Química, Universidad de la República (Uruguay)
Dr. R. C. Lloyd
Chirotech Technology Ltd, 410 Cambridge Science Park
Milton Road, Cambridge, CB40PE (UK)
[**] The research leading to these results has received funding from the

European Union Seventh Framework Programme (FP7/2007-2013) under grant agreements n° 266025 (BIONEXGEN) and n° 245144 (AMBIOCAS). Funding also provided by BBSRC (CASE award to JH) and ANII MOV_CA_2012_1_7759 and CSIC-UdelaR-Uruguay.

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201309208.

Angew. Chem. Int. Ed. 2014, 53, 1-5

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Wiley Online Library

These are not the final page numbers!

reductive amination of prochiral ketones, thereby providing the corresponding chiral amines.^[3a-e] MAO-N catalyzes the oxygen-dependent conversion of amines into imines and is typically selective for the (*S*)-enantiomer.^[3f-I] Variants of MAO-N have been exploited for the deracemization of primary, secondary, and tertiary amines with diverse structural motifs.^[3a,f-j] The development of several chemoenzymatic routes^[3c,4] to industrially important target molecules by employing these two enzyme classes is testament to the advances in protein engineering^[1a,5] that have resulted in the development of biocatalysts with the desired substrate scope, selectivity, and stability.

2,5-Disubstituted pyrrolidines are important scaffolds in pharmaceutical compounds^[6] and natural products,^[7] and considerable efforts have been devoted to developing asymmetric routes to both *cis*- and *trans*-disubstituted derivatives that show moderate to good stereoselectivity.^[8] The lack of stereofacial bias induced by the preexisiting C2-stereocenter means that obtaining the *trans*-diastereomers through reduction of the corresponding imine in high *de* is not straightforward. Our approach (Scheme 1) features a highly selective



Scheme 1. A chemoenzymatic approach for the synthesis of 2,5disubstituted pyrrolidines by employing an ω -transaminase (TA) and a monoamine oxidase (MAO-N).

TA-mediated reductive amination of an achiral 1,4-diketone to generate an optically active pyrroline followed by diastereoselective chemoenzymatic conversion into the corresponding pyrrolidine by MAO-N/NH₃·BH₃.

Initially, we examined the ω -TA-mediated selective monoamination of commercially available 1,4-diketone **1**a, which bears a small methyl substituent and a large phenyl substituent (Scheme 2). The first example of the asymmetric bioamination of 1,5-diketones was recently reported with excellent regio- and stereoselectivity achieved.^[9] We found the commercially available (*S*)-selective transaminase ATA113 to be highly regioselective in mediating the reductive

a) Reductive amination with (S)-selective transaminase



Scheme 2. Preparative-scale (25 mm **1a**) reductive amination of diketone **1a** mediated by (S)-selective ATA113 or (*R*)-selective ATA117, followed by spontaneous cyclization.

amination of **1a** exclusively on the methyl ketone at a substrate concentration of 25 mM with L-alanine as the amine donor. The resulting 1,4-amino ketone (S)-**2a** subsequently underwent spontaneous cyclization to provide pyrroline (S)-**3a** in high yield (91%) and excellent *ee* (>99%). The lactate dehydrogenase (LDH)/glucose dehydrogenase (GDH) system was used to drive the equilibrium towards the product and recycle the NAD⁺ cofactor (see the Supporting Information). The (R)-selective transaminase ATA117 also catalyzed the regio- and stereoselective monoamination of **1a** to afford pyrroline (R)-**3a** in 65% yield and >99% *ee*.

Having established an effective means of accessing optically pure pyrrolines on a preparative scale, we subsequently explored a route for the diastereoselective synthesis of 2,5-disubstituted pyrrolidine **4a** starting from **3a** (Scheme 3). 2,6-Disubstituted piperidines have been prepared through a chemoenzymatic route employing an ω -transaminase followed by diastereoselective hydrogenation using Pd/C.^[9] However, the same strategy is not applicable to the diastereoselective synthesis of 2,5-disubstituted pyrrol-



Scheme 3. a, b) Analytical-scale synthesis of (2S,5R)- and (2R,5R)-4a mediated by MAO-N D5. c) One-pot TA/MAO-N cascade for the preparative-scale asymmetric synthesis of (2S,5R)-4a. Reduction of the starting diketone by NH₃·BH₃ prevented the addition of all of the reagents concurrently.

idines owing to poor diastereoselectivity in the reduction step. We envisaged using MAO-N variants in combination with NH₃·BH₃ for the asymmetric synthesis of 4a. Two MAO-N variants (D5 and D9) were selected based on their known activity and excellent selectivity towards structurally related amine frameworks, including pyrrolidines and piperidines.^[3h,k] Our strategy relies upon MAO-N variants displaying complete regio- and stereoselectivity to avoid stereorandomization of the C2-stereocenter generated by the (S)-selective ω -TA. Imine **3a** is in equilibrium with the open-chain amino ketone (S)-2a and hence optimization of the MAO-N/ NH3·BH3 oxidation/reduction cycle was necessary in order to prevent the formation of undesired amino alcohol as a side product. Ketone reduction was minimized by lowering the concentration of the MAO-N biocatalyst during the reaction while maintaining a high concentration of NH₃·BH₃. Under these conditions, rapid reduction of pyrroline 3a occurred, thus ensuring that a minimal concentration of the amino ketone was present during the biotransformation.

Treatment of (S)-**3a** with NH₃·BH₃ afforded **4a** initially as a mixture of diastereoisomers with a slight excess of the (2S,5S)-isomer ($de \approx 10\%$; Scheme 3a). Both MAO-N variants mediated the oxidation of the (2S,5S)-**4a** diastereoisomer exclusively and displayed complete regioselectivity for the more bulky phenyl side of the pyrrolidine. Following successive rounds of selective oxidation with the MAO-N D5 variant and nonselective reduction with NH₃·BH₃, (2S,5R)-**4a** was isolated in greater than 99% *de*. Despite a bias towards the formation of the *cis* diastereoisomer upon reduction with NH₃·BH₃, the combination with MAO-N yielded solely the *trans* reduction product (2S,5R)-**4a**. The complementary regioselectivity displayed by the ω -TA and MAO-N variants circumvents epimerization of the (S)-C2-center and provides a method for accessing optically pure 2,5-pyrrolidines.

Having developed efficient individual biocatalytic routes for the synthesis of optically pure pyrroline **3a** and the target chiral 2,5-pyrrolidine **4a**, we next sought to combine the ω -TA and MAO-N biocatalysts in a one-pot cascade (Scheme 3c). Diketone **1a** was exposed to ATA 113 followed by MAO-N and NH₃·BH₃, and the target (2*S*,5*R*)-**4a** was obtained in 82 % yield and >99% *de*.

To allow access to the (2R,5R)-4a diastereoisomer, the (R)-3a enantiomer, derived from the use of ATA117, was treated with the NH₃·BH₃/MAO-N combination (Scheme 3b). Following nonselective reduction to give a mixture of (2R,5S)-4a and (2R,5R)-4a, both MAO-N variants mediated the selective oxidation of the (2R,5S)-isomer to provide (2R,5R)-4a exclusively after successive rounds of oxidation/reduction. The stereochemistry at C2 has a minimal effect on the activity of the MAO-N enzyme and no effect on the stereoselectivity; the target (2R,5R)-3a was isolated in >99% *de*.

The generality of the TA/MAO-N cascade process was investigated by examining a series of diketones (1b-g) using ATA113, as well as a novel transaminase (*pf*-ATA) from *Pseudogulbenkiania ferrooxidans*^[10] (Table 1). *Pf*-ATA shares 95% sequence identity with the transaminase from *Chromobacterium violaceum* (*cv*-ATA; ATCC 12472).^[11] ATA113 mediated the reductive amination of diketones 1b-g, the

www.angewandte.org

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Table 1: TA-mediated reductive amination of 1 a-g.



products of which spontaneously cyclized to yield pyrrolines **3b-g** as the sole regioisomers with excellent conversion and high ee values. Unsurprisingly, replacement of the small methyl group by a larger ethyl substituent resulted in a slightly reduced ee (entries 11 and 13). The biotransformations performed with Pf-ATA proceeded with reduced selectivity, with ee values lower than those achieved with ATA113. Interestingly, replacing the methyl substituent by an ethyl group resulted in a switch in stereoselectivity to give (R)-3 f,g as the predominant enantiomers (entries 12 and 14, see the Supporting Information for absolute configuration and ee determination). We also compared the selectivity observed with pf-ATA to that of the related cv-ATA against diketones 1a and 1d, e and noted comparable conversion and selectivity (see the Supporting Information). The (R)-selective ATA117 also mediated the reductive amination of diketones 1a and 1d, e in >99% conversion and ee (see the Supporting Information).

The efficiency of the MAO-N/NH₃·BH₃ step with the isolated pyrrolines **3b–g** was next examined (Table 2). In general, the D9 variant showed higher selectivity and employing either the D5 or D9 MAO-N variants allowed access to all of the Me/Ar and Et/Ar substituted pyrrolidines in excellent *de*. We have also extended the one-pot TA/MAO-N cascade for the synthesis of (2S,5R)-**4b**, (2S,5R)-**4d**, and (2S,5R)-**4e** in > 99% conversion and > 99% *de* when starting from the corresponding diketones (Table 3), thus demonstrating the generality of this one-pot approach.

In summary, the combination of two complementary biocatalysts has been demonstrated in a novel one-pot chemoenzymatic cascade for the regio- and stereoselective synthesis of a panel of 2,5-disubstituted pyrrolidines from the corresponding 1,4-diketones. The transaminase ω -TA is highly selective for the sterically less demanding methyl ketone while the monoamine oxidase MAO-N shows an overwhelming preference for the more bulky portion of the

Table 2: MAO/NH₃·BH₃-mediated asymmetric synthesis of (2*S*,5*R*)-**4**a-g.



Entry	Substrate	MAO-N variant	de [%] (2S,5R)
1 2 3	3 a 3 b	D5 D9 D5	> 99 96 88
5 6	3c	D9 D5 D9	>99 >99 >99
7	3 d	D5	>99
8		D9	>99
9	3 e	D5	68
10		D9	>99
11	3 f	D5	64
12		D9	>99
13	3 g	D5	56
14		D9	96

Table 3: ATA113/MAO-N one-pot cascade for the synthesis of (25,5*R*)-**4a**, (25,5*R*)-**4b**, (25,5*R*)-**4d**, and (25,5*R*)-**4e**.

Ketone	ω-TA	MAO-N	Conv. [%]	de [%]
1 a ^[a]	ATA113	D5	> 99	>99 (2 <i>S</i> ,5 <i>R</i>)- 4 a
1 b ^[b]	ATA113	D9	>99	>99 (2 <i>S</i> ,5 <i>R</i>)- 4 b
1 d ^[b]	ATA113	D9	>99	>99 (2 <i>S</i> ,5 <i>R</i>)- 4 d
1 e ^[b]	ATA113	D9	>99	>99 (2 <i>S</i> ,5 <i>R</i>)- 4 e

[a] 25 mм substrate; [b] 5 mм substrate.

corresponding pyrrolidine. The compatibility of the two biocatalysts means that the reaction can be performed in one pot without the need for costly intermediate purification steps. The chemoenzymatic approach exploits four distinct biocatalytic operations and takes advantage of the complementary regioselectivity displayed by the ω -TA and MAO variants to establish two stereogenic centers. All of the biocatalysts described herein are commercially available^[12] and hence readily accessible for practical application.

Received: October 22, 2013 Revised: December 11, 2013 Published online:

Keywords: biocatalysis · cascade reactions · monoamine oxidase · pyrrolidines · transaminase

- [1] a) M. T. Reetz, J. Am. Chem. Soc. 2013, 135, 12480-12496;
 b) N. J. Turner, E. O' Reilly, Nat. Chem. Biol. 2013, 9, 285-288.
- [2] a) J. H. Schrittwieser, J. Sattler, V. Resch, F. G. Mutti, W. Kroutil, *Curr. Opin. Chem. Biol.* 2011, *15*, 249–256; b) I. Oroz-Guinea, E. García -Junceda, *Curr. Opin. Chem. Biol.* 2013, 17, 236–249; c) E. Ricca, B. Brucher, J. H. Schrittwieser, *Adv. Synth. Catal.* 2011, *353*, 2239–2262; d) J. H. Sattler, M. Fuchs, K. Tauber, F. G.

einheim www.angewandte.org These are not the final page numbers!



Mutti, K. Faber, J. Pfeffer, T. Haas, W. Kroutil, *Angew. Chem.* **2012**, *124*, 9290–9293; *Angew. Chem. Int. Ed.* **2012**, *51*, 9156–9159.

- [3] a) M. Höhne, U. Bornscheuer, ChemCatChem 2009, 1, 42-51; b) S. Mathew, H. Yun, ACS Catal. 2012, 2, 993-1001; c) C. K. Savile, J. M. Janey, E. C. Mundorff, J. C. Moore, S. Tam, W. R. Jarvis, J. C. Colbeck, A. Krebber, F. J. Fleitz, J. Brands, P. N. Devine, G. W. Huisman, G. J. Hughes, Science 2010, 329, 305-309; d) D. Koszelewski, K. Tauber, K. Faber, W. Kroutil, Trends Biotechnol. 2010, 28, 324-332; e) L. Frodsham, M. Golden, S. Hard, M. N. Kenworthy, D. J. Klauber, K. Leslie, C. Macleod, R. E. Meadows, K. R. Mulholland, J. Reilly, C. Squire, S. Tomasi, D. Watt, A. S. Wells, Org. Process Res. Dev. 2013, 17, 1123-1130; f) M. Alexeeva, A. Enright, M. J. Dawson, M. Mahmoudian, N. J. Turner, Angew. Chem. 2002, 114, 3309-3312; Angew. Chem. Int. Ed. 2002, 41, 3177-3180; g) R. Carr, M. Alexeeva, A. Enright, T. S. C. Eve, M. J. Dawson, N. J. Turner, Angew. Chem. 2003, 115, 4955-4958; Angew. Chem. Int. Ed. 2003, 42, 4807-4810; h) C. J. Dunsmore, R. Carr, T. Fleming, N. J. Turner, J. Am. Chem. Soc. 2006, 128, 2224-2225; i) I. Rowles, K. J. Malone, L. L. Etchells, S. C. Willies, N. J. Turner, ChemCatChem 2012, 4, 1259-1261; j) A. Znabet, M. M. Polak, E. Janssen, F. J. J. de Kanter, N. J. Turner, R. V. A. Orru, E. Ruijter, Chem. Commun. 2010, 46, 7918-7920; k) D. Ghislieri, A. P. Green, M. Pontini, S. C. Willies, I. Rowles, A. Frank, G. Grogan, N. J. Turner, J. Am. Chem. Soc. 2013, 135, 10863-10869; 1) D. Ghislieri, D. Houghton, A. P. Green, S. C. Willies, N. J. Turner, ACS Catal. 2013, 3, 2869-2872.
- [4] a) T. Li, J. Liang, A. Ambrogelly, T. Brennan, G. Gloor, G. Huisman, J. Lalonde, A. Lekhal, B. Mijts, S. Muley, L. Newman, M. Tobin, G. Wong, A. Zaks, X. Zhang, J. Am. Chem. Soc. 2012, 134, 6467–6472; b) B. de Lange, D. J. Hyett, P. J. D. Maas, D. Mink, F. B. J. van Assema, N. Sereinig, A. H. M. de Vries, J. G. de Vries, ChemCatChem 2011, 3, 289–292.

- [5] a) U. T. Bornscheuer, G. W. Huisman, R. J. Kazlauskas, S. Lutz, J. C. Moore, K. Robins, *Nature* **2012**, 485, 185–194; b) N. J. Turner, *Nat. Chem. Biol.* **2009**, 5, 567–573.
- [6] a) J. B. Brenneman, S. F. Martin, Org. Lett. 2004, 6, 1329–1331;
 b) F. Xu, Org. Lett. 2013, 15, 1324–1345;
 c) S. Hanessian, M. Bayrakdarian, X. Luo, J. Am. Chem. Soc. 2002, 124, 4716–4721.
- [7] a) V. K. Aggarwal, C. J. Astle, M. Rogers-Evans, Org. Lett. 2004, 6, 1469-1471; b) S. Zhang, L. Xu, L. Miao, H. Shu, M. L. Trudell, J. Org. Chem. 2007, 72, 3133-3136; c) A. Goti, S. Cicchi, V. Mannucci, F. Cardona, F. Guarna, P. Marino, T. Tejero, Org. Lett. 2003, 5, 4235-4238; d) B. M. Trost, D. B. Horne, M. J. Woltering, Angew. Chem. 2003, 115, 6169-6172; Angew. Chem. Int. Ed. 2003, 42, 5987-5990; e) E. A. Severino, C. R. D. Correia, Org. Lett. 2000, 2, 3039-3042.
- [8] a) F. A. Davis, M. Song, A. Augustine, J. Org. Chem. 2006, 71, 2779–2786; b) G. S. Lemen, J. P. Wolfe, Org. Lett. 2010, 12, 2322–2325; c) M. G. Moloney, T. Panchal, R. Pike, Org. Biomol. Chem. 2006, 4, 3894–3897; d) C. Enkisch, C. Schneider, Eur. J. Org. Chem. 2009, 5549–5564; e) K. R. Campos, A. Kalpars, J. H. Waldman, P. G. Dormer, C.-y. Chen, J. Am. Chem. Soc. 2006, 128, 3538–3539; f) F. A. Davis, J. Zhang, H. Qui, Y. Wu, Org. Lett. 2008, 10, 1433–1436.
- [9] R. C. Simon, B. Grischek, F. Zepeck, A. Steinreiber, F. Belaj, W. Kroutil, *Angew. Chem.* 2012, *124*, 6817–6820; *Angew. Chem. Int. Ed.* 2012, *51*, 6713–6716.
- [10] K. G. Byrne-Bailey, K. A. Weber, J. D. Coates, J. Bacteriol. 2012, 194, 2400-2401.
- [11] U. Kaulmann, K. Smithies, M. E. B. Smith, H. C. Hailes, J. M. Ward, *Enzyme Microb. Technol.* 2007, 41, 628–637.
- [12] a) MAO-N Screening Kits: http://www.discovery-bc.co.uk/ monoamineoxidase.php (2013); b) Transaminases: http://www. codexis.com.

www.angewandte.org

© 2014 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

Communications



E. O'Reilly, C. Iglesias, D. Ghislieri, J. Hopwood, J. L. Galman, R. C. Lloyd, N. J. Turner* ______ III-- III

A Regio- and Stereoselective ω-Transaminase/Monoamine Oxidase Cascade for the Synthesis of Chiral 2,5-Disubstituted Pyrrolidines



Mirror mirror on the wall: A ω -transaminase (ω -TA)/monoamine oxidase (MAO-N) cascade process for the asymmetric synthesis of chiral 2,5-disubstituted pyrrolidines is reported. The methodology exploits the complementary regio- and stereoselectivity displayed by both enzymes, which ensures that the stereogenic center established by the TA reaction is not affected by the MAO-N catalyzed step.