

Octahedral Metal Carbonyls.

61*. The Mechanism of *cis-trans* Isomerization During Ligand-substitution Reactions Affording *cis*-Bis(triphenylphosphine)tetracarbonyltungsten(0)


GERARD R. DOBSON**, HANI H. AWAD and STEPHEN S. BASSON†

Center for Organometallic Research and Department of Chemistry, North Texas State University, Denton, Tex. 76203-5068, U.S.A.

(Received March 4, 1986)

It has been reported that during thermal ligand substitution in $(\text{tmpa})\text{W}(\text{CO})_4$ ($\text{tmpa} = N,N,N',N'$ -tetramethyl-1,3-diaminopropane) proceeding by stoichiometry (1) and mechanism (2) ($L = \text{phosphine}$, $(\text{tmpa})\text{W}(\text{CO})_4 + 2L \longrightarrow L_2\text{W}(\text{CO})_4 + \text{tmpa}$ (1)

phosphite) [1], stereochemical rearrangement takes place in the five-coordinated intermediate (2c) [2].



This rearrangement thus would account for the creation of the two observed isomeric reaction products, *cis*- and *trans*- $L_2\text{W}(\text{CO})_4$, and it could be inferred that the five-coordinate species undergo stereochemical rearrangement on the time-scale of ligand-substitution. We report several lines of evidence which demonstrate


*For Part 60, see ref. 8.

**Author to whom correspondence should be addressed.

†Robert A. Welch Foundation Post-Doctoral Fellow, 1985. Permanent address: Department of Chemistry, University of the Orange Free State, Bloemfontein, South Africa.

that this proposal is not correct but, rather, that the isomerization takes place through a well-documented [3] non-dissociative scrambling process. Evidence further indicates that where a five-coordinate intermediate is produced, no stereochemical rearrangement takes place.

(a) Pulsed laser flash photolysis at 43.0 °C in xylene or chloroform of *cis*-(pip)(PPh_3) $\text{W}(\text{CO})_4$ (pip = piperidine) was carried out as previously described [4, 5]. The complex, synthesized via the method of Darensbourg and Kump [6], reacts upon photolysis via W-N bond fission [5]. At 430 nm two independent exponential decays are observed in the presence of PPh_3 . The faster decay (k_4 , eqn. (3)) $\approx 10^7 \text{ M}^{-1} \text{ s}^{-1}$ in xylene and $8.9(5) \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$ in chloroform) has been shown [5] to involve



bimolecular reaction of the specifically solvated *cis*- $[\text{Ph}_3\text{PW}(\text{CO})_4]$ transient with PPh_3 . The slower decay ($k_4' = 6.8 \times 10^5 \text{ M}^{-1} \text{ s}^{-1}$ in xylene and $8.4(1) \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$ in chloroform) has been attributed to the reaction of the corresponding *trans* isomer [4]. Only one decay would be observed if these species were undergoing rapid intramolecular inter-conversion. Thus the overall photolysis mechanism is that in eqn. (3).

(b) ^{31}P NMR studies of chelate ring-displacement (eqn. (2)) by PPh_3 (0.305 M, Jeol FX90Q spectrometer) in $\text{CHCl}_3/\text{CDCl}_3$ at 52.3 °C clearly demonstrate that the peak at 27.529 ppm (downfield from a PPh_3 internal standard), attributable to the *cis*-(Ph_3P) $_2\text{W}(\text{CO})_4$ product, grows in intensity at the reaction's earlier stages but that the corresponding *trans* peak (at 32.678 ppm) appears only after an

induction period. The chemical shifts of the two isomers were identified through comparison with those of authentic samples*. These observations indicate that the *cis*-(Ph₃P)₂W(CO)₄ reaction product is produced first, with subsequent rearrangement to the *trans* isomer. Mosbo and coworkers have made similar observations for analogous systems for L = PPhMe₂ and PPh₂Me [3g].

(c) The rates of the thermal *cis*-to-*trans* isomerization of *cis*-(Ph₃P)₂W(CO)₄ were studied spectrophotometrically at 400 nm in toluene at 52.5 °C in the absence and in the presence of P(OEt)₃. The results demonstrate that the process is unimolecular and that the derived rate constant ($k_5 + k_{-5}$, eqn. (3)) is independent of the concentration of P(OEt)₃ (k in the absence of P(OEt)₃ = $6.06(3) \times 10^{-4} \text{ s}^{-1}$; in its presence (three concentrations over the range 0.1–0.5 M) $k_5 + k_{-5} = 5.79(9) \times 10^{-4} \text{ s}^{-1}$). Further, in the presence of added L (=P(OEt)₃, pip), no (L)(PPh₃)W(CO)₄ products are observed to form during the isomerization process. Thus, a five-coordinate species formed via W–PPh₃ bond-fission is not produced, since pulsed laser flash photolysis studies indicate that L readily trap [Ph₃PW(CO)₄] [5].

Thus it may be concluded that *cis*-(Ph₃P)₂W(CO)₄ undergoes isomerization exclusively via a slow, intramolecular scrambling process probably involving a trigonal twist [3].

The results may be contrasted to those for the analogous Mo complex, for which thermal isomerization *does* take place via a mechanism involving Mo–P bond-fission; it has been suggested that isomerization of the *cis*-[PPh₃Mo(CO)₄] intermediate to afford *trans*-(PPh₃)₂Mo(CO)₄ may be a concerted process [7].

*The *cis* complex was prepared photolytically employing a 2:1 ratio of W(CO)₆ to PPh₃ dissolved in hexane (400 w medium pressure Hg lamp for 20 min). The precipitated product was washed with hexane until the peak characteristic of (Ph₃P)W(CO)₅ at ca. 2075 cm⁻¹ has disappeared. The *trans* isomer was synthesized via displacement of tmpa from (tmpa)W(CO)₄ in chlorobenzene at 52 °C over three hours.

Acknowledgements

The support of this research by the National Science Foundation (Grant CHE 84 15153), the Robert A. Welch Foundation (Grant B-434) and the North Texas State University Faculty Research Fund is gratefully acknowledged. The pulsed laser flash photolysis studies and analyses of the data produced were performed at the Center for Fast Kinetics Research (CFKR) at the University of Texas at Austin. The CFKR is supported jointly by the Biotechnology Branch of the Division of Research Resources of the National Institutes of Health (Grant RR00886) and the University of Texas at Austin. The experimental help and technical expertise of the staff at CFKR, in particular that of Stephen J. Atherton, are greatly appreciated. We are also grateful for the experimental assistance of Mr. Corby Young, Mr. Charles B. Dobson and Dr. Paul H. Wermer.

References

- 1 G. R. Dobson and A. Moradi-Araghi, *Inorg. Chim. Acta*, **31**, 263 (1978).
- 2 D. J. Darensbourg, G. R. Dobson and A. Moradi-Araghi, *J. Organomet. Chem.*, **116**, C17 (1976).
- 3 (a) W. Majunke, D. Leibfritz, T. Mack and H. tom Dieck, *Chem. Ber.*, **108**, 3025 (1975); (b) D. J. Darensbourg, *Inorg. Chem.*, **18**, 14 (1979); (c) D. J. Darensbourg and B. J. Baldwin, *J. Am. Chem. Soc.*, **101**, 6447 (1979); (d) D. J. Darensbourg, R. Kudarowski and W. Schenk, *Inorg. Chem.*, **21**, 2488 (1982); (e) D. J. Darensbourg and R. L. Gray, *Inorg. Chem.*, **23**, 2993 (1984); (f) D. T. Dixon, J. C. Kola and J. A. S. Howell, *J. Chem. Soc., Dalton Trans.*, 1307 (1984); (g) M. L. Boyles, D. V. Brown, D. A. Drake, C. K. Hostetler, C. K. Maves and J. A. Mosbo, *Inorg. Chem.*, **24**, 3126 (1985).
- 4 G. R. Dobson, I. Bernal, M. G. Reisner, C. B. Dobson and S. E. Mansour, *J. Am. Chem. Soc.*, **107**, 525 (1985).
- 5 G. R. Dobson, K. J. Asali, S. S. Basson and C. B. Dobson, *Inorg. Chem.*, submitted for publication.
- 6 D. J. Darensbourg and R. J. Kump, *Inorg. Chem.*, **17**, 2680 (1978).
- 7 D. J. Darensbourg and A. H. Graves, *Inorg. Chem.*, **18**, 1257 (1979).
- 8 P. H. Wermer, C. B. Dobson and G. R. Dobson, *J. Organomet. Chem.*, submitted for publication.