

Synthesis of 2-pyranylidene or (2-furyl)carbene–chromium complexes from conjugated enyne carbonyl compounds with $\text{Cr}(\text{CO})_5(\text{THF})$

Koji Miki, Tomomi Yokoi, Fumiaki Nishino, Kouichi Ohe *, Sakae Uemura *

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan

Received 21 August 2001; received in revised form 18 September 2001; accepted 23 September 2001

Abstract


The reaction of β -ethynyl α,β -unsaturated esters or amides with $\text{Cr}(\text{CO})_5(\text{L})$ ($\text{L} = \text{Et}_2\text{O}$, THF), derived from chromium hexacarbonyl, gives 2-pyranylidene–chromium complexes in good yields. The conjugated ene-carbonyl-vinylidene complex is a key intermediate to these Fischer-type oxacarbene complexes. In contrast, β -ethynyl α,β -unsaturated ketones react with $\text{Cr}(\text{CO})_5(\text{THF})$ to give (2-furyl)carbene complexes via nucleophilic attack of the carbonyl oxygen atom to the intermediate π -alkyne–chromium complexes. Some chemical transformation, such as [4 + 2] type cyclization of 2-pyranylidene complexes with alkynes and aldehyde formation from (2-furyl)carbene complexes, has also been described. © 2002 Elsevier Science B.V. All rights reserved.

Keywords: Chromium; Pyranylidene; Furylcarbene; Vinylidene

1. Introduction

Transition metal carbene complexes, especially Fischer-type oxacarbene complexes, have been widely studied and applied to synthetic organic chemistry as versatile organometallic reagents [1,2]. Typical oxacarbene complexes like $(\text{CO})_5\text{M}=\text{C}(\text{OR})\text{R}'$ are readily prepared by the reaction of metal hexacarbonyl with a range of organolithium and organomagnesium reagents. Nucleophilic addition of an alcohol to the α - and β -carbons of a vinylidene–metal complex ($\text{M} = \text{C}_\alpha = \text{C}_\beta$, $\text{M} = \text{Cr}$, Mo , W) also leads to the formation

of oxacarbene complexes in catalytic and stoichiometric reactions [3–5]. We have recently reported that 2-pyranylidene–molybdenum and –tungsten complexes, as cyclic Fischer-type oxacarbene complexes, were prepared from $\text{M}(\text{CO})_5(\text{L})$ ($\text{M} = \text{Mo}$, W ; $\text{L} = \text{THF}$, NEt_3) and β -ethynyl α,β -unsaturated esters or amides via nucleophilic attack of the carbonyl oxygen to the α -carbon of a vinylidene–metal intermediate (Scheme 1) [6,7]. However, the preparation of 2-pyranylidene–chromium complexes from $\text{Cr}(\text{CO})_5(\text{L})$ ($\text{L} = \text{Et}_2\text{O}$, THF), which was prepared by photo-irradiation, was unsuccessful. The complex $\text{Cr}(\text{CO})_5(\text{NMe}_3)$ [8] prepared from $\text{Cr}(\text{CO})_6$ and trimethylamine-*N*-oxide in situ was only slightly effective for the preparation of 2-pyranylidene–chromium complex. After we investigated the optimal conditions to obtain 2-pyranylidene–chromium complexes, we found that the reaction of β -ethynyl α,β -unsaturated esters or amides with $\text{Cr}(\text{CO})_5(\text{sol})$, which was prepared from $\text{Cr}(\text{CO})_6$ by irradiation in a suitable solvent, in the presence of Et_3N as an additive gave effectively the corresponding pyranylidene–chromium complexes. In sharp contrast with this result, the reaction of β -ethynyl α,β -unsaturated ketones with $\text{Cr}(\text{CO})_5(\text{THF})$ afforded new (2-furyl)carbene–



Scheme 1.


* Corresponding authors. Tel.: +81-75-753-5687; fax: +81-75-753-3573.

E-mail address: uemura@scl.kyoto-u.ac.jp (S. Uemura).

chromium complexes. The 5-*exo*-dig nucleophilic cyclization in a π -alkyne complex of the ketone with $\text{Cr}(\text{CO})_5$ might be responsible for the formation of nonstabilized carbene complexes. 2-Pyranylidene–chromium complexes undergo [4 + 2] type cycloaddition with dimethyl acetylenedicarboxylate followed by demetallation to give tetralin derivatives, while (2-furyl)carbene–chromium complexes react with molecular oxygen to give the corresponding furfurals.


2. Results and discussion

We have reported that 2-pyranylidene–tungsten or –molybdenum complexes were prepared from β -ethynyl α,β -unsaturated esters or amides with $\text{W}(\text{CO})_5(\text{THF})$ [9] or $\text{Mo}(\text{CO})_5(\text{NEt}_3)$ [8]. However, the corresponding pyranlydene–chromium complexes could not be prepared under similar conditions using $\text{Cr}(\text{CO})_5(\text{Et}_2\text{O})$ or $\text{Cr}(\text{CO})_5(\text{THF})$. When we examined the reaction with these chromium complexes in more detail, we found that the reaction of β -ethynyl α,β -unsaturated ester **1a** with $\text{Cr}(\text{CO})_6$ and $\text{Me}_3\text{N}^+\text{O}^-$ gave 2-pyranylidene–chromium complex **2a** in 9% yield (Eq. (1)). We consid-




ered that trimethylamine, which is generated from the reaction between $\text{Cr}(\text{CO})_6$ and $\text{Me}_3\text{N}^+\text{O}^-$ might play some role in the formation of pyranlydene complex. Then we tried the reaction using an amine as an additive. The reaction of **1a** with $\text{Cr}(\text{CO})_5(\text{Et}_2\text{O})$ in the absence of triethylamine gave none of **2a** (Table 1). However, the reaction proceeded in the presence of triethylamine to give complex **2a** in 27% isolated yield together with the recovered **1a** (30–50%). The use of THF instead of Et_2O as a solvent gave a better yield of **2a**. The reactions of other substrates with $\text{Cr}(\text{CO})_5(\text{THF})$ in the presence of triethylamine were next examined, typical results of which being shown in Table 2.

Reactions of **1a** with a lesser amount of $\text{Cr}(\text{CO})_5(\text{THF})$ gave lower yields of **2a** in 15% (1.2 equivalents of Cr) and 33% (2.0 equivalents of Cr), respectively (entries 1 and 2). The reaction of an amide **1b** with $\text{Cr}(\text{CO})_5(\text{THF})$ at room temperature for 0.5 h also gave the corresponding 2-pyranylidene–chromium



Scheme 2.

Table 1
Preparation of **2a**



Solvent	Additive	Isolated yield (%)
Et_2O (30 ml)		
Et_2O (30 ml)	Et_3N (0.1 ml)	27
THF (20 ml)	Et_3N (0.1 ml)	62

^a The solution of $\text{Cr}(\text{CO})_5(\text{L})$ was prepared by irradiating $\text{Cr}(\text{CO})_6$ (in Et_2O or THF) for 4 h with a high-pressure Hg lamp.

Table 2


Preparation of pyranlydene–chromium complexes from conjugated enyne carbonyls and $\text{Cr}(\text{CO})_5(\text{THF})$ ^{a,b}

entry	1	product	yield (%) ^c
1 ^d			15
2 ^e	1a	2a	33
3	1a	2a	62
4 ^f			68
5 ^f			69
6 ^f			50

^a Reactions were carried out with **1** (0.2 mmol) at room temperature for 2 h under Ar. ^b The solution of $\text{Cr}(\text{CO})_5(\text{L})$ was prepared by irradiating a solution of $\text{Cr}(\text{CO})_6$ (0.6 mmol) and triethylamine (0.1 ml) in THF (20 ml) for 4 h with a high-pressure Hg lamp. ^c Isolated yield. ^d **1a** (0.5 mmol). ^e **1a** (0.3 mmol). ^f Reaction time was 0.5 h.

complex **2b** in 68% yield (entry 4). The ester **1c** and the amide **1d** gave similar complexes **2c** and **2d** in 69 and 50% yields, respectively (entries 5 and 6). At present, we assume that triethylamine facilitates the formation of a vinylidene–chromium intermediate from a π -alkyne–chromium complex (Scheme 2) [7b].


The structure of pyranilidene–chromium complexes represents α -metallopyrone. Therefore, we envisaged cycloaddition reaction of these 2-pyranilidene–chromium complexes with dienophiles. The reaction of **2a** with an excess amount of dimethyl acetylenedicarboxylate (DMAD, 15 equivalents) gave the tetralin derivative **3a** in 26% isolated yield (Table 3). The use of solvent, i.e., toluene, did not positively affect the reaction. The reaction in DMSO resulted in demetallation leading to the formation of compound **1a** in 59% yield. The complex **2b** also reacted with DMAD to give the corresponding tetralin derivative **3b** in 25% yield. However, the reaction of **2a** with an electron-rich dienophile, i.e., ethyl ethynyl ether, gave only a trace amount of tetralin derivative as one isomer (Eq. (2)). The transformation was explained by assuming [4 + 2] cycloaddition reaction between 2-pyranilidene–chromium complex and the acetylene followed by a pericyclic demetallation of chromium hexacarbonyl (Scheme 3).



Next, we examined the reactions of β -ethynyl α,β -unsaturated ketone instead of β -ethynyl α,β -unsaturated ester or amide with chromium and tungsten carbonyls. When 2-ethynyl-1-cyclohexenyl phenyl ketone (**4a**) was treated with three equivalents of $\text{Cr}(\text{CO})_5(\text{THF})$ in the presence of triethylamine, the corresponding 2-pyranilidene–chromium complex **2e** was obtained only in 8% yield together with many unidentified products (Eq. (3)). On the other hand, in the absence of triethylamine the reaction of **4a** with three equivalents of $\text{Cr}(\text{CO})_5(\text{THF})$ did not afford **2e**, and instead (2-furyl)carbene–chromium **5a** was isolated in 52% yield as a blue solid (Eq. (4)) [10]. The complex **5a** is marginally stable during purification with silica gel column

Table 3

Reaction of pyranilidene–chromium complex with dimethyl acetylene dicarboxylate





Entry	Substrate	Solvent	Product	Yield (%) ^a
1	2a	–	3a	26
2	2b	–	3b	25
3	2a	Toluene	3a	9
4	2a	Dioxane	3a	0
5	2a	DMSO	3a	0 ^b

^a Isolated yield.


^b An enyne carbonyl compound **1a** was obtained in 59% yield.

chromatography. 2-Thienyl ketone **4b** also gave the corresponding (2-furyl)carbene–chromium complex **5b** in 59% yield. When **4a** was treated with three equivalents of $\text{W}(\text{CO})_5(\text{THF})$, (2-furyl)carbene–tungsten complex **5c** was isolated in 56% yield [11] (Eq. (4)).




However, in the reaction of an alkyl vinyl ketone, such as ethyl 2-ethynyl-1-cyclohexenyl ketone (R = Et in **4**), we could not succeed in the isolation of the corresponding (2-furyl)carbene complex. These complexes **5a–5c** were stored for several days under N_2 atmosphere, but they gradually decomposed in CDCl_3 or under oxygen atmosphere. When the isolated carbene complex **5a** was stirred in THF under oxygen atmosphere, furfural derivative **6a** was obtained in 76% yield. The displacement of metal carbonyl moiety with the oxygen atom in carbene complex by molecular oxygen is a well-known process [12]. Most plausible pathway from **4** to **5** is shown in Scheme 4.

In the pre-equilibrium between a π -alkyne complex **7** and a vinylidene complex **8**, 5-*exo*-dig cyclization takes place as a favorable process to give the furylcarbene



Scheme 3.



Scheme 4.

complex. The presence of triethylamine, which may promote the isomerization of a π -alkyne complex to a vinylidene complex, in fact, precluded the formation of the (2-furyl)carbene complex. The mode of the complex formation is also attributed to the difference in the reactivity of carbonyl functionality of β -ethynyl α,β -unsaturated carbonyl compounds.

3. Conclusions

We have demonstrated that 2-pyranylidene–chromium complexes were prepared from β -ethynyl α,β -unsaturated esters and amides in the presence of triethylamine. In sharp contrast, the reaction of β -ethynyl α,β -unsaturated ketones with $M(\text{CO})_5(\text{THF})$ ($M = \text{Cr}, \text{W}$) gave (2-furyl)carbene complexes selectively via nucleophilic attack of a carbonyl oxygen to a π -alkyne complex. Some chemical transformation of newly prepared 2-pyranylidene–chromium complexes and (2-furyl)carbene–chromium and tungsten complexes has been described: [4 + 2] type cyclization with alkynes in the former cases and aldehyde formation in the latter cases, respectively.

4. Experimental

4.1. General

Tetrahydrofuran (THF) and Et_2O were distilled from sodium benzophenone ketyl under Ar. Analytical thin-layer chromatographies (TLC) were performed with silica gel 60 Merck F-254 plates. Column chromatographies were performed with Merck silica gel 60. The NMR spectra were measured for solutions in CDCl_3 or $\text{THF}-d_8$ with Me_4Si as an internal standard (^1H and ^{13}C) and the following abbreviations are used: s, singlet; d, doublet; t, triplet; q, quartet; quint, quintet; m, multiplet. IR spectra were recorded with a FT-IR spectrometer. Melting points (m.p.) are uncorrected. Elemental analyses were performed at Microanalytical Center of Kyoto University. All new compounds were prepared and fully characterized.

4.2. Synthesis of 2-pyranylidene–chromium complex **2** from **1** or **4a**

4.2.1. 2-Pyranylidene–chromium complex **2a** from **1a**

A solution of $\text{Cr}(\text{CO})_6$ (0.13 g, 0.60 mmol) in THF (20 ml) and Et_3N (0.1 ml) under Ar was irradiated by a Hg lamp (350 nm) at room temperature (r.t.) for 4 h. To this orange solution under Ar was added a solution of **1a** (33 mg, 0.20 mmol) in THF (1 ml) by a syringe. The mixture was stirred at r.t. for 0.5–2 h. The solvent was removed under reduced pressure, and the residue

was subjected to column chromatography on SiO_2 with hexane– EtOAc (v/v = 10:1) as an eluent to afford **2a** (44 mg, 62% yield) as an orange solid, m.p. 91.5–94.3 °C. IR (KBr, cm^{-1}): 1897, 1935, 1976, 2053. ^1H -NMR (CDCl_3 , 300 MHz, 25 °C): δ = 1.72–1.84 (m, 4H), 2.38–2.49 (m, 2H), 2.55–2.64 (m, 2H), 4.32 (s, 3H), 7.33 (s, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz, 25 °C): δ = 20.4, 21.2, 21.4, 29.0, 56.9, 108.2, 134.7, 154.9, 171.9, 218.6 (Cr–CO), 223.4 (Cr–CO), 251.0 (Cr=C). HRMS (FAB); m/z : 355.9995 ($[\text{M}^+]$, calc. for $\text{C}_{15}\text{H}_{12}\text{CrO}_7$; 355.9988).

4.2.2. 2-Pyranylidene–chromium complex **2b** from **1b**

An orange solid (68% yield), m.p. 118.4–121.5 °C. IR (KBr, cm^{-1}): 1888, 1909, 1965, 2045. ^1H -NMR (CDCl_3 , 300 MHz, 25 °C): δ = 1.36 (t, J = 6.9 Hz, 6H), 1.69–1.79 (m, 4H), 2.44–2.50 (m, 2H), 2.53–2.60 (m, 2H), 3.63 (q, J = 6.9 Hz, 4H), 6.96 (s, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz, 25 °C): δ = 13.7, 21.1, 22.6, 25.4, 29.5, 44.7, 108.6, 132.0, 151.2, 170.1, 219.3 (Cr–CO), 223.9 (Cr–CO), 240.7 (Cr=C). Anal. Calc. for $\text{C}_{18}\text{H}_{19}\text{CrNO}_6$: C, 54.41; H, 4.82; N, 3.53. Found: C, 54.27; H, 5.03; N, 3.40%.

4.2.3. 2-Pyranylidene–chromium complex **2c** from **1c**

An orange solid (69% yield), m.p. 71.0–73.5 °C. IR (KBr, cm^{-1}): 1898, 1917, 1976, 2053. ^1H -NMR (CDCl_3 , 300 MHz, 25 °C): δ = 4.32 (s, 3H), 6.15 (d, J = 8.1 Hz, 1H), 7.15 (dd, J = 8.1, 8.1 Hz, 1H), 7.68 (d, J = 8.1 Hz, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz, 25 °C): δ = 57.0, 96.9, 134.1, 139.6, 175.0, 217.9 (Cr–CO), 223.4 (Cr–CO), 269.6 (Cr=C). HRMS (FAB); m/z : 301.9518 ($[\text{M}^+]$, calc. for $\text{C}_{11}\text{H}_6\text{CrO}_7$; 301.9519).

4.2.4. 2-Pyranylidene–chromium complex **2d** from **1d**

An orange solid (50% yield), m.p. 128.3–129.5 °C. IR (KBr, cm^{-1}): 1906, 1966, 2052. ^1H -NMR (CDCl_3 , 300 MHz, 25 °C): δ = 3.75–3.87 (m, 8H), 6.00 (m, 1H), 6.97 (m, 1H), 7.26 (m, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz, 25 °C): δ = 44.7, 65.9, 95.4, 129.4, 137.6, 169.2, 218.7 (Cr–CO), 223.8 (Cr–CO), 258.1 (Cr=C). HRMS (FAB); m/z : 356.9934 ($[\text{M}^+]$, calc. for $\text{C}_{14}\text{H}_{11}\text{CrNO}_7$; 356.9941).

4.2.5. 2-Pyranylidene–chromium complex **2e** from **4a**

The ketone **4a** (0.13 g, 0.60 mmol) and Et_3N (1.0 ml) were used. A red solid (8% yield), m.p. 132.0–134.6 °C. IR (KBr, cm^{-1}): 1918, 1974, 2051. ^1H -NMR ($\text{THF}-d_8$, 300 MHz, 25 °C): δ = 1.78–1.91 (m, 4H), 2.76–2.84 (m, 4H), 7.56–7.62 (m, 3H), 7.79–7.86 (m, 2H), 7.95 (s, 1H); ^{13}C -NMR ($\text{THF}-d_8$, 75 MHz, 25 °C): δ = 22.0, 23.0, 26.2, 29.6, 124.8, 129.6, 129.7, 132.0, 133.1, 141.0, 150.6, 175.5, 219.2 (Cr–CO), 224.3 (Cr–CO), 268.8 (Cr=C). HRMS (FAB); m/z : 402.0185 ($[\text{M}^+]$, calc. for $\text{C}_{20}\text{H}_{14}\text{CrO}_6$; 402.0196).

4.3. Typical procedure for [4 + 2] type cycloaddition of **2**

4.3.1. Dimethyl 5,6,7,8-tetrahydro-1-methoxy-2,3-naphthalenedicarboxylate (**3a**)

A mixture of complex **2a** (0.034 g, 0.095 mmol) and dimethyl acetylenedicarboxylate (0.2 ml, 1.6 mmol) in a sealed tube was stirred at 90 °C for 12 h under N₂. The mixture was subjected to column chromatography on SiO₂ with hexane–EtOAc (v/v = 10:1) as an eluent to afford **3a** (7 mg, 0.025 mmol, 26% yield) as a colorless liquid. ¹H-NMR (CDCl₃, 300 MHz, 25 °C): δ = 1.77–1.84 (m, 4H), 2.74–2.82 (m, 4H), 3.79 (s, 3H), 3.86 (s, 3H), 3.94 (s, 3H), 7.52 (s, 1H). ¹³C-NMR (CDCl₃, 75 MHz, 25 °C): δ = 22.2, 22.4, 23.7, 29.3, 52.3, 52.6, 61.7, 125.3, 126.6, 127.2, 136.9, 140.4, 155.0, 165.8 (CO), 168.4 (CO).

4.3.2. Dimethyl 1-diethylamino-5,6,7,8-tetrahydro-2,3-naphthalenedicarboxylate (**3b**)

A yellow liquid (25% yield). ¹H-NMR (CDCl₃, 300 MHz, 25 °C): δ = 1.00 (t, J = 7.2 Hz, 6H), 1.72–1.82 (m, 4H), 2.65–2.75 (m, 2H), 2.75–2.86 (m, 2H), 2.90–3.12 (m, 4H), 3.85 (s, 3H), 3.89 (s, 3H), 7.57 (s, 1H). ¹³C-NMR (CDCl₃, 75 MHz, 25 °C): δ = 14.4, 22.4, 22.6, 26.6, 29.6, 47.4, 52.1, 52.2, 124.7, 128.1, 134.9, 139.4, 144.2, 146.8, 166.2 (CO), 169.8 (CO).

4.4. Synthesis of furylcarbene complexes **5**

4.4.1. Chromium complex **5a**

A solution of Cr(CO)₆ (0.13 g, 0.60 mmol) in THF (20 ml) under Ar was irradiated by a Hg lamp (350 nm) at r.t. for 4 h. To the yellow solution under Ar was added a solution of **4a** (42 mg, 0.20 mmol) in THF (1 ml) by a syringe. The mixture was stirred at r.t. for 0.5 h. The solvent was removed under reduced pressure, and the residue was subjected to column chromatography on SiO₂ with hexane–EtOAc (v/v = 10:1) as an eluent to afford **5a** (42 mg, 0.10 mmol, 52% yield) as a blue solid. IR (KBr, cm^{−1}): 651, 826, 1934, 1994, 2045. ¹H-NMR (THF-*d*₈, 300 MHz, 25 °C): δ = 1.84–2.00 (m, 4H), 2.75–2.83 (m, 2H), 2.95–3.04 (m, 2H), 7.60–7.69 (m, 3H), 8.16–8.25 (m, 2H), 13.5 (s, 1H). ¹³C-NMR (THF-*d*₈, 75 MHz, 25 °C): δ = 23.0, 24.0, 24.1, 24.4, 127.7, 128.8, 129.9, 130.8, 132.9, 138.4, 168.2, 171.7, 219.6 (Cr–CO), 233.9 (Cr–CO), 284.4 (Cr=C). HRMS (FAB); m/z : 402.0195 ([M⁺], calc. for C₂₀H₁₄CrO₆: 402.0219).

4.4.2. Chromium complex **5b**

A blue solid (59% yield). IR (KBr, cm^{−1}): 1933, 1970, 2041. ¹H-NMR (THF-*d*₈, 300 MHz, 25 °C): δ = 1.73–1.90 (m, 4H), 2.66 (t, J = 3.9 Hz, 2H), 2.77 (t, J = 3.9 Hz, 2H), 7.27–7.31 (m, 1H), 7.88–7.94 (m, 2H), 13.0 (s, 1H). ¹³C-NMR (THF-*d*₈, 75 MHz, 25 °C):

δ = 21.5, 21.6, 22.2, 22.7, 125.3, 129.2, 130.3, 131.5, 132.8, 137.2, 163.3, 169.9, 218.2 (Cr–CO), 232.3 (Cr–CO), 276.4 (Cr=C). HRMS (FAB); m/z : 407.9756 ([M⁺], calc. for C₁₉H₁₂CrO₆S: 407.9760).

4.4.3. Tungsten complex **5c**

A solution of W(CO)₆ (0.21 g, 0.60 mmol) in THF (20 ml) under Ar was irradiated by a Hg lamp (350 nm) at r.t. for 4 h. To the yellow solution under Ar was added a solution of **4a** (42 mg, 0.2 mmol) in THF (1 ml) by a syringe. The mixture was stirred at r.t. for 0.5 h. The solvent was removed under reduced pressure, and the residue was subjected to column chromatography on SiO₂ with hexane–EtOAc (v/v = 10:1) as an eluent to afford **5c** (60 mg, 0.11 mmol, 56% yield) as a blue solid. IR (KBr, cm^{−1}): 580, 677, 825, 1920, 1934, 2052. ¹H-NMR (THF-*d*₈, 300 MHz, 25 °C): δ = 1.94–2.10 (m, 4H), 2.54–2.60 (m, 2H), 3.07–3.13 (m, 2H), 7.64–7.78 (m, 3H), 8.26–8.32 (m, 2H), 13.3 (s, 1H). ¹³C-NMR (THF-*d*₈, 75 MHz, 25 °C): δ = 21.3, 22.5, 22.5, 23.5, 127.4, 127.9, 129.5, 129.8, 131.5, 141.7, 166.5, 171.3, 198.1 (W–CO), 210.2 (W–CO), 247.8 (W=C). HRMS (FAB); m/z : 534.0300 ([M⁺], calc. for C₂₀H₁₄WO₆: 534.0286).


4.5. Synthesis of furfural derivatives **6**

4.5.1. Furfural **6a**

A solution of **5a** (22 mg, 0.054 mmol) in THF (5 ml) was stirred at r.t. under O₂ atmosphere for 12 h. The solvent was removed under reduced pressure and the residue was subjected to column chromatography on SiO₂ with hexane–EtOAc (v/v = 8:1) as an eluent to afford **6a** (9.2 mg, 0.041 mmol, 76% yield) as a white solid, m.p. 91.3–94.8 °C. IR (KBr, cm^{−1}): 696, 773, 828, 1650, 1669. ¹H-NMR (THF-*d*₈, 300 MHz, 25 °C): δ = 1.73–1.86 (m, 4H), 2.81–2.87 (m, 2H), 2.87–2.93 (m, 2H), 7.31–7.38 (m, 1H), 7.41–7.49 (m, 2H), 7.76–7.81 (m, 2H), 9.73 (s, 1H). ¹³C-NMR (THF-*d*₈, 75 MHz, 25 °C): δ = 22.8, 23.4, 23.9, 24.4, 122.6, 124.7, 127.1, 129.9, 130.2, 132.1, 147.8, 152.9, 178.2 (CHO). HRMS (FAB); m/z : 227.1067 ([M + H⁺], calc. for C₁₅H₁₄O₂: 227.1072).

4.5.2. Furfural **6b**

A white solid (76% yield), m.p. 100.5–101.6 °C. IR (KBr, cm^{−1}): 698, 819, 853, 1445, 1635, 1663. ¹H-NMR (THF-*d*₈, 300 MHz, 25 °C): δ = 1.75–1.90 (m, 4H), 2.68–2.76 (m, 2H), 2.87–2.94 (m, 2H), 7.14 (dd, J = 3.6, 4.8 Hz, 1H), 7.42 (d, J = 4.8 Hz, 1H), 7.47 (d, J = 3.6 Hz, 1H), 9.69 (s, 1H). ¹³C-NMR (THF-*d*₈, 75 MHz, 25 °C): δ = 22.7, 23.1, 23.5, 24.1, 121.6, 126.6, 128.2, 129.3, 134.0, 136.7, 147.4, 149.6, 177.8 (CHO). Anal. Calc. for C₁₃H₁₂O₂S: C, 67.21; H, 5.21. Found: C, 66.97; H, 5.17%.



Scheme 5.

4.6. Synthesis of substrates

Methyl 2-(trimethylsilyl)-1-cyclohexenecarboxylate (**10**) was prepared from cyclohexanone in several steps: see the supporting information of Ref. [6a] (Scheme 5).

4.6.1. *N*-Methyl *N*-methoxy

2-(trimethylsilyl)-1-cyclohexenecarboxamide (**11**) [13]

A solution of methyl 2-(trimethylsilyl)-1-cyclohexenecarboxylate (**10**) (3.6 g, 15 mmol) in benzene (5 ml) was added to benzene (40 ml) and hexane (45 ml) solution of $\text{Me}_2\text{AlNMe(OMe)}$ prepared from a hexane solution of Me_3Al (45 ml, 45 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (4.4 g, 45 mmol). The resulting solution was heated under reflux for 3 h. The reaction mixture was cooled down to r.t., and hydrolyzed by slow and cautious addition of 1.0 N HCl (30 ml). The upper organic layer was separated, and the aqueous layer was extracted with EtOAc (30 ml $\times$ 3). The organic extracts were combined, washed with brine, and dried over MgSO_4 . The solvent was removed under reduced pressure to give the residual liquid, which was subjected to column chromatography on SiO_2 with hexane–EtOAc ($v/v = 4:1$) as an eluent to afford *N*-methyl *N*-methoxy 2-(trimethylsilyl)-1-cyclohexenecarboxamide (**11**) (3.6 g, 14 mmol, 92% yield) as a pale yellow liquid. IR (KBr, cm^{-1}): 845, 1659 (C=O), 2144 ($\text{C}\equiv\text{C}$). $^1\text{H-NMR}$ (CDCl_3 , 300 MHz, 25 $^\circ\text{C}$): $\delta = 0.12$ (s, 9H), 1.57–1.65 (m, 4H), 2.14–2.27 (m, 4H), 3.23 (s, 3H), 3.71 (s, 3H). $^{13}\text{C-NMR}$ (CD_2Cl_2 , 100 MHz, -25°C): $\delta = -0.1$ (SiCH_3), -0.1 (SiCH_3), 21.5 (CH_2), 21.5 (CH_2), 21.8 (CH_2), 21.9 (CH_2), 26.5 (CH_2), 26.7 (CH_2), 28.8 (CH_2), 29.5 (CH_2), 31.8 (NCH_3), 35.7 (NCH_3), 60.3 (OCH_3), 62.0 (OCH_3), 96.4 ($\text{C}\equiv$), 97.2 ($\text{C}\equiv$), 103.0 ($\text{C}\equiv$), 104.5 ($\text{C}\equiv$), 116.7 ($\text{C}=\text{}$), 119.4 ($\text{C}=\text{}$), 140.0 ($\text{C}=\text{}$), 143.1 ($\text{C}=\text{}$), 166.4 (C=O), 170.8 (C=O) as a mixture of rotamers. Anal. Calc. for $\text{C}_{14}\text{H}_{23}\text{NO}_2\text{Si}$: C, 63.35; H, 8.73; N, 5.28. Found: C, 63.07; H, 8.47; N, 5.21%.

4.7. Preparation of β -ethynyl- α,β -unsaturated carbonyl compounds **4**

4.7.1. 2-Ethynyl-1-cyclohexenyl phenyl ketone (**4a**)

To a solution of **11** (0.47 g, 2.0 mmol) in THF (10

ml) was added preformed phenylmagnesium bromide (4.2 mmol) in THF (15 ml) at 0 $^\circ\text{C}$, and the mixture was stirred at r.t. for 4 h. The reaction mixture was poured into saturated NH_4Cl solution (30 ml), and the aqueous layer was extracted with EtOAc (20 ml $\times$ 3). The combined organic layer was dried over MgSO_4 . The solvent was removed under reduced pressure and then 1 N KOH solution (2.0 ml) was added to the residue in MeOH (20 ml) at r.t. After stirring for 0.5 h, this solution was poured into saturated NH_4Cl solution (50 ml). The aqueous layer was extracted with EtOAc (20 ml $\times$ 3) and the combined organic layer was dried over MgSO_4 . The solvent was removed under reduced pressure and the residue was subjected to column chromatography on SiO_2 with hexane–EtOAc ($v/v = 20:1$) as an eluent to afford **4a** (0.22 g, 1.0 mmol, 52% yield) as a colorless liquid. IR (neat, cm^{-1}): 648, 690, 709, 732, 916, 1662 (C=O), 2093 ($\text{C}\equiv\text{C}$), 3292 (=C-H). $^1\text{H-NMR}$ (CDCl_3 , 300 MHz, 25 $^\circ\text{C}$): $\delta = 1.71$ – 1.77 (m, 4H), 2.28–2.38 (m, 4H), 2.81 (s, 1H), 7.40–7.48 (m, 2H) 7.50–7.57 (m, 1H), 7.85–7.92 (m, 2H). $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz, 25 $^\circ\text{C}$): $\delta = 21.4$, 21.8, 27.3, 29.8, 82.2, 82.3, 119.8, 128.5, 129.5, 133.2, 136.0, 145.8, 199.0 (CO). HRMS (FAB); m/z : 211.1119 ($[\text{M} + \text{H}^+]$, calc. for $\text{C}_{15}\text{H}_{14}\text{O}$: 211.1123).

4.7.2. 2-Ethynyl-1-cyclohexenyl 2-thienyl ketone (**4b**)

To a solution of thiophene (0.46 ml, 6.0 mmol) in THF (5 ml) was added *n*-BuLi (4 ml, 1.6 M in hexane, 6.4 mmol) at 0 $^\circ\text{C}$ under N_2 , and the mixture was stirred at r.t. for 2 h. This mixture was added to a solution of **10** (0.80 g, 3.0 mmol) in THF (5 ml) at 0 $^\circ\text{C}$, and the mixture was stirred at r.t. for 10 min. The reaction mixture was poured into saturated NH_4Cl solution (20 ml), and the aqueous layer was extracted with EtOAc (10 ml $\times$ 3). The combined organic layer was dried over MgSO_4 . The solvent was removed under reduced pressure and then K_2CO_3 (1.0 g, 7.5 mmol) was added to the residue in MeOH (30 ml) at r.t. After stirring for 12 h, this solution was poured into saturated NH_4Cl solution (100 ml). The aqueous layer was extracted with EtOAc (30 ml $\times$ 3) and the combined organic layer was dried over MgSO_4 . The solvent was removed under reduced pressure and the residue was subjected to column chromatography on SiO_2 with hexane–EtOAc ($v/v = 20:1$) as an eluent to afford **4b** (0.28 g, 1.3 mmol, 43% yield) as a colorless liquid. IR (neat, cm^{-1}): 643, 734, 774, 856, 1262, 1285, 1411, 1644 (C=O), 2094 ($\text{C}\equiv\text{C}$), 2935, 3288 (=C-H). $^1\text{H-NMR}$ (CDCl_3 , 300 MHz, 25 $^\circ\text{C}$): $\delta = 1.71$ – 1.77 (m, 4H), 2.28–2.40 (m, 4H), 2.92 (s, 1H), 7.13 (dd, $J = 4.2$, 4.2 Hz, 1H), 7.68 (d, $J = 4.2$ Hz, 2H). $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz, 25 $^\circ\text{C}$): $\delta = 21.4$, 21.7, 27.3, 29.7, 82.0, 82.2, 119.4, 128.1, 134.6, 134.7, 143.0, 145.8, 191.2 (CO). HRMS (FAB); m/z : 217.0685 ($[\text{M} + \text{H}^+]$, calc. for $\text{C}_{13}\text{H}_{12}\text{OS}$: 217.0687).

Acknowledgements

We thank Mr Yasuaki Hisamoto in our department for measuring HRMS spectra. This research was supported on Priority Areas (A) “Exploitation of Multi-Element Cyclic Molecules” from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- [1] L.S. Hegedus, *Transition Metals in the Synthesis of Complex Organic Molecules*, University Science Books, Mill Valley, 1994, p. 151.
- [2] M.J. Winter, in: E.W. Abel, F.G.A. Stone, G. Wilkinson (Eds.), *Comprehensive Organometallic Chemistry II*, vol. 5, Pergamon/Elsevier, Oxford, 1995, p. 155.
- [3] For a review, see: F.E. McDonald, *Chem. Eur. J.* 5 (1999) 3103.
- [4] For leading references, see: (a) E.O. Fischer, D. Plabst, *Chem. Ber.* 107 (1974) 3326;
(b) C.P. Casey, R.L. Anderson, *J. Chem. Soc. Chem. Commun.* (1975) 895;
(c) A. Parlier, H. Rudler, *J. Chem. Soc. Chem. Commun.* (1986) 514;
(d) K.H. Dötz, W. Sturm, H.G. Alt, *Organometallics* 6 (1987) 1424;
(e) F.E. McDonald, C.B. Connolly, M.M. Gleason, T.B. Towne, K.D. Treiber, *J. Org. Chem.* 58 (1993) 6952;
(f) P. Quayle, S. Rahman, E.L.M. Ward, J. Herbert, *Tetrahedron Lett.* 35 (1994) 3801;
(g) F.E. McDonald, M.M. Gleason, *J. Am. Chem. Soc.* 118 (1996) 6648;
(h) B. Schmidt, P. Kocienski, G. Reid, *Tetrahedron* 52 (1996) 1617;
(i) C.F. Bernasconi, *Chem. Soc. Rev.* 26 (1997) 299;
(j) F.E. McDonald, H.Y.H. Zhu, *J. Am. Chem. Soc.* 120 (1998) 4246;
(k) F.E. McDonald, K.S. Reddy, Y. Díaz, *J. Am. Chem. Soc.* 122 (2000) 4304;
- [5] For reviews on vinylidene–metal complexes, see: (a) M.I. Bruce, A.G. Swincer, *Adv. Organomet. Chem.* 22 (1983) 59;
(b) M.I. Bruce, *Chem. Rev.* 91 (1991) 197;
(c) C. Bruneau, P.H. Dixneuf, *Acc. Chem. Res.* 32 (1999) 311.
- [6] Preparation of pyranlydene–metal complexes via vinylidene–metal intermediates, see: (a) K. Ohe, K. Miki, T. Yokoi, F. Nishino, S. Uemura, *Organometallics* 19 (2000) 5525;
(b) N. Iwasawa, M. Shido, K. Maeyama, H. Kusama, *J. Am. Chem. Soc.* 122 (2000) 10226.
- [7] Transformation of vinylidene–metal intermediates in conjugated systems, see: (a) Y. Wang, M.G. Finn, *J. Am. Chem. Soc.* 117 (1995) 8045;
(b) K. Ohe, M. Kojima, K. Yonehara, S. Uemura, *Angew. Chem. Int. Ed. Engl.* 35 (1996) 1823;
(c) C.A. Merlic, M.E. Pauly, *J. Am. Chem. Soc.* 118 (1996) 11319;
(d) T. Manabe, S. Yanagi, K. Ohe, S. Uemura, *Organometallics* 17 (1998) 2942;
(e) K. Maeyama, N. Iwasawa, *J. Am. Chem. Soc.* 120 (1998) 1928;
(f) K. Maeyama, N. Iwasawa, *J. Org. Chem.* 64 (1999) 1344;
- [8] (a) M. Wrighton, *Chem. Rev.* 74 (1974) 401;
(b) F.E. McDonald, C.C. Schultz, *J. Am. Chem. Soc.* 116 (1994) 9363.
- [9] W. Strohmeier, K. Gerlach, *Chem. Ber.* 94 (1961) 398.
- [10] A part of this work has been presented at the 79th Annual Meeting of the Chemical Society of Japan in Kobe, March 31, 2001.
- [11] Most recently, Iwasawa and co-workers have reported the intervention of a similar carbene complex in the tungsten-catalyzed cyclization of 2-ethynylphenyl alkyl ketones. N. Iwasawa, M. Shido, H. Kusama, *J. Am. Chem. Soc.* 123 (2001) 5814.
- [12] Chromium moieties are readily displaced with the oxygen atom by molecular oxygen, see: (a) R.B. Silveanu, R.A. Olofson, *Chem. Commun.* (1968) 1313;
(b) E.O. Fischer, S. Riedmüller, *Chem. Ber.* 107 (1974) 915.
- [13] For preparation of ketones, see: (a) M.F. Lipton, A. Baska, S.M. Weinreb, *Org. Synth.* 59 (1979) 49;
(b) J.I. Levin, E. Turos, S.M. Weinreb, *Synth. Commun.* 12 (1982) 989;
(c) S. Nahm, S.M. Weinreb, *Tetrahedron Lett.* 22 (1981) 3815.