Synthesis, Molecular Structure, and Reactivity of a Half-Sandwich Vanadium(III) Imido Complex: The First Vanadium(V) Alkylidene

Jan-Karel F. Buijink and Jan H. Teuben*

Groningen Center for Catalysis and Synthesis, Department of Chemistry, University of Groningen, Nijenborgh 4, NL-9747 AG Groningen, The Netherlands

Huub Kooijman and Anthony L. Spek[†]

Bijvoet Center for Biomolecular Research, Crystal and Structural Chemistry, University of Utrecht, Padualaan 8, NL-3584 CH Utrecht, The Netherlands

Received May 11, 1994[®]

Summary: One of the trimethylphosphine ligands in $CpV(NAr)(PMe_3)_2$ (3; $Ar = 2,6-C_6H_3$ -i- Pr_2), obtained by magnesium reduction of $CpV(NAr)Cl_2$ (2), can be substituted by CO, ethylene, or diphenylacetylene to give the corresponding π -acceptor (L) complex CpV(NAr)(L)- (PMe_3) (L = CO, 4; $L = C_2H_4, 5$; L = PhC = Ph, 6). 3 reacts with $Ph_3P = CHPh$ to form the first vanadium(V) alkylidene, $CpV(NAr)(=CHPh)(PMe_3)$ (7).

Imido ligands have been widely used as stabilizing ligands in high-oxidation-state transition-metal complexes¹ and have played an important role in the development of well-defined olefin metathesis² and ringopening metathesis polymerization³ (ROMP) catalysts, i.e. alkylidenes. The use of imido ligands in organovanadium chemistry has allowed the isolation of thermally stable alkylvanadium(V) complexes⁴ but did not lead to the isolation of vanadium(V) alkylidenes.⁵ Half-sandwich vanadium imido complexes have received little attention,^{4a-c,6} in spite of the potentially useful isolobal relationship between these complexes and titanocene derivatives.⁷

(3) (a) Schrock, R. R.; Feldman, J.; Cannizzo, L. F.; Grubbs, R. H. Macromolecules 1987, 20, 1169-1172. (b) Feldman, J.; Schrock, R. R. In Progress in Inorganic Chemistry; Lippard, S. J., Ed.; Wiley-Interscience: New York, 1991; Vol. 39, 1-74.

(7) Pez, G. P.; Armor, J. N. In Advances in Organometallic Chemistry; Stone, F. G. A., West, R., Eds.; Academic Press: New York, 1981; Vol. 19, pp 1-50. Here we wish to report the synthesis and characterization of a simple half-sandwich imidovanadium(III) complex, as well as some aspects of its reactivity, including the synthesis and molecular structure of the first vanadium(V) alkylidene.

In analogy with the established route to 4-substituted (phenylimido)vanadium complexes,^{4a} reaction of VOCl₃ with ArNCO (Ar = 2,6-C₆H₃-*i*-Pr₂) in refluxing *n*-octane gave V(NAr)Cl₃ (1)⁸ in a nearly quantitative yield (Scheme 1). 1 reacts cleanly with CpSiMe₃ to produce the moderately air sensitive half-sandwich imidovanadium dichloride CpV(NAr)Cl₂ (2)⁹ in good yield (Scheme 1). Reduction of 2 with magnesium in the presence of trimethylphosphine (Scheme 1) yields the diamagnetic⁹ vanadium(III) bis(trimethylphosphine) complex CpV-(NAr)(PMe₃)₂ (3).¹⁰

An X-ray structure determination of 3^{11} (Figure 1) shows a nearly tetrahedral geometry about vanadium with a Cp_{centroid}-V(1)-N(1) bond angle of 129.99(11)°, P-V(1)-N(1) bond angles of 99° (average) and P-V-(1)-Cp_{centroid} bond angles of 114° (average). All such interligand angles are similar to those found in structurally characterized d² group 4 metallocene complexes

 $^{^{\}dagger}$ To whom correspondence concerning crystallographic data should be addressed.

[®] Abstract published in Advance ACS Abstracts, June 15, 1994.
(1) Nugent, W. A.; Mayer, J. M. Metal-Ligand Multiple Bonds;
Wiley-Interscience: New York, 1988.
(2) (a) Schrock, R. R.; DePue, R. T.; Feldman, J.; Schaverien, J. C.;

^{(2) (}a) Schrock, R. R.; DePue, R. T.; Feldman, J.; Schaverien, J. C.;
Dewan, J. C.; Liu, A. H. J. Am. Chem. Soc. 1988, 110, 1423–1435. (b)
Schaverien, C. J.; Dewan, J. C.; Schrock, R. R. J. Am. Chem. Soc. 1986, 108, 2771–2773. (c) Murdzek, J. S.; Schrock, R. R. Organometallics 1987, 6, 1373–1374.

^{(4) (}a) Devore, D. D.; Lichtenhan, J. D.; Takusagawa, F.; Maatta,
E. A. J. Am. Chem. Soc. 1987, 109, 7408-7416. (b) Preuss, F.; Becker,
H.; Häusler, H.-J. Z. Naturforsch. 1987, 42B, 881-888. (c) Preuss, F.;
Becker, H.; Wieland, T. Z. Naturforsch. 1986, 41B, 185-190. (e) de With, J.;
Horton, A. D.; Orpen, A. G. Organometallics 1990, 9, 2207-2209. (f) de With, J.; Horton, A. D.; Orpen, A. G. Organometallics 1993, 12, 1493-1496. (g) de With, J.; Horton, A. D. Angew. Chem., Int. Ed. Engl. 1993, 32, 903-905.

⁽⁵⁾ One "Schrock-type" vanadium(III) alkylidene ((5a) Hessen, B.; Buijink, J.-K. F.; Meetsma, A.; Teuben, J. H.; Helgesson, G.; Håkansson, M.; Jagner, S.; Spek, A. L. Organometallics **1993**, *12*, 2268-2276) and several "Fischer-type" vanadium carbenes ((5b) Berlekamp, M.; Erker, G.; Petersen, J. L. J. Organomet. Chem. **1993**, 458, 97-103 and references therein) are known.

^{(6) (}a) Preuss, F.; Becker, H.; Kaub, J.; Sheldrick, W. S. Z. Naturforsch. 1988, 43B, 1195-1200. (b) Preuss, F.; Wieland, T.; Perner, J.; Heckmann, G. Z. Naturforsch. 1992, 47B, 1355-1362. (c) Preuss, F.; Wieland, T.; Günther, B. Z. Anorg. Allg. Chem. 1992, 609, 45-50.
(7) Pez, G. P.; Armor, J. N. In Advances in Organometallic Chem-

⁽⁸⁾ All complexes gave satisfactory elemental analyses and have also been characterized by ¹H, ¹³C, and ⁵¹V NMR and IR. Full details may be found in the supplementary material.

⁽⁹⁾ To a solution of VCl₃[N-2,6-C₆H₃-(*i*-Pr)₂] (7.50 g, 22.5 mmol) in 60 mL of CH₂Cl₂ was added CpSiMe₃ (3.12 g, 22.5 mmol). After the mixture was stirred for 48 h at 40 °C, the color had changed from green to golden brown and the mixture was filtered. After removal of the solvent in vacuo 15 mL of pentane was added to the residue and removed in vacuo, yielding 6.59 g (18.2 mmol; 81%) of product as a brown powder: ¹H NMR δ 6.82 (d, 2H, ³J_{HH} = 7 Hz, NAr H_m), 6.72 (t, 1H, ³J_{HH} = 7, NAr H_p), 5.89 (s, 5H, Cp), 3.91 (sept, 2H, ³J_{HH} = 6.8, CHMe₂), 1.19 (d, 12H, ³J_{HH} = 6.8, CHMe₂); ¹³C NMR δ 159 (br, NAr C_{*ipso*}), 150.5 (NAr C₀), 129.3 (NAr C_p), 122.8 (NAr C_m), 116.6 (Cp), 28.2 (CHMe₂), 24.4 (CHMe₂); ⁵¹V NMR δ -183 ($\Delta v_{1/2} = 875$ Hz). (10) A mixture of CpVCl₂[N-2,6-C₆H₃-(*i*-Pr)₂] (2; 1.09 g, 3.01 mmol), PMe₃ (0.46 g, 6.02 mmol), and magnesium (1.0 g, 41.1 mmol) in 20 mL of THF was stirred for 16 h under argon at room temperature.

⁽¹⁰⁾ A mixture of CpVCl₂[N-2,6-C₆H₃-(*i*-Pr)₂] (2; 1.09 g, 3.01 mmol), PMe₃ (0.46 g, 6.02 mmol), and magnesium (1.0 g, 41.1 mmol) in 20 mL of THF was stirred for 16 h under argon at room temperature. After removal of the solvent in vacuo, the dark residue was extracted with 40 mL of pentane. Concentrating and cooling the green extract to -25 °C gave 0.90 g (2.03 mmol, 67%) of product as dark green crystals in two crops: ¹H NMR δ 7.13 (d, 2H, $^{3}J_{HH} = 7$ Hz, NAr H_m), 7.00 (t, 1H, $^{3}J_{HH} = 7$, NAr H_p), 4.81 (t, 5H, $J_{PH} = 2.1$, Cp), 4.55 (sept. 2H, $^{3}J_{HH} = 6.8$, CHMe₂), 1.22 (d, 12H $^{3}J_{HH} = 6.8$, CHMe₂), 0.99 (virtual t, 18H, PMe₃); ¹³C NMR δ 142.9 (NAr C_{ipso}), 129.3 (NAr C_o), 122.7 (NAr C_p), 121.1 (NAr C_m), 94.1 (Cp), 26.8 (CHMe₂), 24.4 (CHMe₂), 22.6 (dd, ¹J_{PC} = 9.2 Hz, $^{3}J_{PC} = 6.1$); ⁶V NMR δ 265 (t, $J_{PV} = 380$ Hz). (11) 3 crystallizes in the monoclinic space group P2/e (No. 14) with

^{(11) 3} crystallizes in the monoclinic space group $P2_0/c$ (No. 14), with a = 9.0608(3) Å, b = 19.0538(6) Å, c = 16.0943(5) Å, $\beta = 110.466(3)^\circ$, V = 2603.18(15) Å³, Z = 4, 6788 reflections measured, 5360 unique reflections, 298 K, Cu K α radiation, 2.32° < θ < 75.0°. The structure was solved by automated Patterson methods (DIRDIF-92). Refinement on F^2 (SHELXL-93), with H atoms at calculated positions, gave wR2 = 0.115, R1 = 0.047, and S = 1.01.



Figure 1. ORTEP diagram of 3. The thermal ellipsoids are drawn at the 30% probability level. Selected bond distances (Å): V(1)-N(1) = 1.698(2), V(1)-P(1) = 2.3994-(9), V(1)-P(2) = 2.3859(10), V(1)-C(19) = 2.357(4), V(1)-C(20) = 2.340(5), V(1)-C(21) = 2.228(3), V(1)-C(22) = 2.187(4), V(1)-C(23) = 2.268(4). Selected bond angles (deg): V(1)-N(1)-C(1) = 170.7(2), P(1)-V(1)-P(2) = 92.60(4), P(1)-V(1)-N(1) = 101.66(8), P(2)-V(1)-N(1) = 96.69(9), Cp_{centroid}-V(1)-N(1) = 129.99(11), P(1)-V-(1)-Cp_{centroid} = 113.24(6), P(2)-V(1)-Cp_{centroid} = 115.49-(7).

Scheme 1. Synthesis and Reactivity of 3



such as Cp₂Ti(PMe₃)₂.¹² The V–N distance of 1.698(2) Å is characteristic of V–N multiple bonding but is relatively long compared to most known V=N bonds in V(IV) or V(V) systems (usually 1.60–1.68 Å)¹ and similar to the one found in the comparatively electron rich V(III) complex CpV[=NC(CMe₃)=CHCMe₃](dmpe) (1.707(2) Å).^{5a} The V–N–Ar bond angle is almost linear

 $(170.7(2)^{\circ})$. Three short and two long V-C_{Cp} bond distances indicate a trend toward η^3 coordination of the cyclopentadienyl moiety. Similar behavior has been observed for several cyclopentadienyl imido complexes of group 5 metals.^{4b,13}

One of the trimethylphosphine ligands in 3 can be substituted by CO, ethylene, or diphenylacetylene to give the corresponding π -acceptor (L) complex CpV-(NAr)(L)(PMe₃) (L = CO, 4; L = C₂H₄, 5; L = PhC=CPh, 6; Scheme 1). The reactions proceed under conditions (25-50 °C, 1 atm) milder than those reported for the analogous niobium complex (C₅Me₅)Nb(NAr)(PMe₃)₂,¹⁴ and no disubstitution, as has been reported for titanocene complex Cp₂Ti(PMe₃)₂,^{12,15} is observed. The products can be obtained in good yields as crystalline solids.

The CO stretch for carbonyl complex 4,¹⁶ at 1885 cm⁻¹, and the ¹³C NMR shifts of the vanadium-bonded ethylene carbons of 5^{17} (δ 40.7 and 38.8 ppm) are consistent with considerable back-donation from the vanadium(III) centers of 4 and 5. The ¹³C NMR spectrum of [¹³C]-4,¹⁸ prepared from 3 and ¹³CO, shows a broad plateau-form resonance for the carbonyl carbon at 306 ppm, the broadening being caused by a combination of unresolved coupling to the $I = \frac{7}{2}$ vanadium nucleus and quadrupolar relaxation effects.4e For the same reason no olefinic C-H coupling constants can be determined in the ¹³C NMR spectrum of 5. The ²H NMR spectrum of $[{}^{2}H_{4}]$ -5, ¹⁹ prepared from 3 and $C_{2}D_{4}$, shows resonances at 1.92, 1.76, and 0.75 ppm, which correspond to partially hidden multiplets in the ¹H NMR of 5. The relatively high IR frequency of the $\nu_{C=C}$ vibration (1778 cm^{-1}) in alkyne complex 6 indicates that the alkyne acts as a two-electron donor, in contrast with the alkyne acting as a four-electron donor in structurally characterized CpV(η^2 -PhC=CPh)(PMe_3)₂²⁰ ($\nu_{C=C} = 1600$ cm^{-1}).

Alkylidene transfer from phosphoranes to reduced transition-metal complexes has been reported for Cp₂-Zr²¹ and Cp₂TaMe systems²² and recently for W(IV) imido complexes.²³ Reaction of **3** with Ph₃P=CHPh (Scheme 1) proceeds with loss of PMe₃ and PPh₃ to produce the first vanadium(V) alkylidene, CpV(NAr)-

(17) Selected spectroscopic data for 5: ¹H NMR δ 6.93 (d, 2H, ³J_{HH} = 7 Hz, NAr H_m), 6.85 (t, 1H, ³J_{HH} = 7, NAr H_p), 5.14 (d, 5H, J_{PH} = 2.5, Cp), 3.60 (sept, 2H, ³J_{HH} = 6.8, CHMe₂), 2.00 (m, 2H, C₂H₄), 1.80 (m, 1H, C₂H₄), 1.22 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 1.17 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 0.90 (d, 9H, ²J_{PH} = 7.3, PMe₃), 0.81 (m, 1H, C₃H₄); ¹³C NMR NAr C_{ipso} not observed, δ 145.4 (NAr C_o), 122.7 (NAr C_p), 122.6 (NAr C_m), 99.8 (Cp), 40.8, 38.8 (br s, C₂H₄), 27.0 (CHMe₂), 24.6, 24.4 (CHMe₂), 17.3 (d, ¹J_{PC} = 19.8); ⁵¹V NMR δ -702 (d, J_{FY} = 272).

(18) Selected spectroscopic data for [¹³C]-4: ¹³C NMR (C₆D₆) δ 306 (CO, $\Delta \nu_{1/2} = 1230$ Hz).

(19) Selected spectroscopic data for [²H₄]-5: ²H NMR (C₆D₆) δ 1.92 (br s, 2D, $\Delta \nu_{1/2} = 9$ Hz), 1.76 (br s, 1D, $\Delta \nu_{1/2} = 11$), 0.75 (br s, 1D, $\Delta \nu_{1/2} = 7$).

(20) Hessen, B.; Meetsma, A.; van Bolhuis, F.; Teuben, J. H.; Helgesson, G.; Jagner, S. Organometallics **1990**, 9, 1925-1936.

⁽¹²⁾ Kool, L. B.; Rausch, M. D.; Alt, H. G.; Herberhold, M.; Thewalt, U.; Wolf, B. Angew. Chem., Int. Ed. Engl. 1985, 24, 394-401.

⁽¹³⁾ Williams, D. N.; Mitchell, J. P.; Poole, A. D.; Siemeling, U.; Clegg, W.; Hockless, D. C. R.; O'Neil, P. A.; Gibson, V. C. J. Chem. Soc., Dalton Trans. 1992, 739-751.

⁽¹⁴⁾ Siemeling, U.; Gibson, V. C. J. Organomet. Chem. 1992, 426, C25-C27.

⁽¹⁵⁾ Kool, L. B.; Rausch, M. D.; Alt, H. G.; Herberhold, M.; Honold, B.; Thewalt, U. J. Organomet. Chem. 1987, 320, 37-45.

⁽¹⁶⁾ Selected spectroscopic data for 4: ¹H NMR (toluene- d_8) δ 7.06 (d, 2H, ³J_{HH} = 7 Hz, NAr H_m), 6.96 (t, 1H, ³J_{HH} = 7, NAr H_p), 5.01 (d, 5H, J_{PH} = 2.1, Cp), 4.22 (sept, 2H, ³J_{HH} = 6.8, CHMe₂), 1.32 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 1.26 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 0.82 (d, 9H, ²J_{PH} = 7.1, PMe₃); ¹³C NMR (toluene- d_8) NAr C_{ipso} not observed, δ 144.0 (NAr C_o), 123.1 (NAr C_p), 122.4 (NAr C_m), 95.3 (Cp), 27.7 (CHMe₂), 24.5, 23.3 (CHMe₂), 19.6 (d, ¹J_{PC} = 20.6); ⁵¹V NMR (toluene- d_8) δ -548 (d, J_{PV} = 390 Hz).

 $(= CHPh)(PMe_3) (7).^{24} Transfer of the alkylidene moiety from phosphorus to vanadium is clearly indicated by the downfield shift of the alkylidene H_{\alpha} (13.45 ppm) and C_{\alpha} (304 ppm) resonances. Nuclear Overhauser effect experiments²⁵ indicate a syn arrangement of the alkylidene ligand relative to the imido group in the single alkylidene rotamer that is formed.$

The structure of 7 was confirmed by an X-ray diffraction study²⁶ (Figure 2), showing a slightly distorted tetrahedral geometry about the metal, similar to the geometry in 3. All interligand angles are similar to those found in the structurally characterized niobium analogue (C₅Me₅)Nb(NAr)(=CHPh)PMe₃,²⁷ whereas the differences in metal-ligand distances correlate well with the difference in ionic radii of V(V) and Nb(V)²⁸ (e.g. V=C = 1.922(6) Å versus Nb=C = 2.026(4) Å). The V(1)-C(24)-C(25) angle of 136.9(4)° is indicative of an undistorted²⁹ alkylidene ligand, which is in marked contrast with the very obtuse V=C-C angle of 173.3-(3)° that was observed in the only other structurally characterized 3d-metal alkylidene, CpV(CHCMe₃)-(dmpe).^{5a}

(22) (a) Sharp, P. R.; Schrock, R. R. J. Organomet. Chem. 1979, 171, 43-51. (b) van Asselt, A.; Burger, B. J.; Gibson, V. C.; Bercaw, J. E. J. Am. Chem. Soc. 1986, 108, 5347-5349.

(23) Johnson, L. K.; Frey, M.; Ulibarry, T. A.; Virgil, S. C.; Grubbs,
 R. H.; Ziller, J. W. J. Am. Chem. Soc. 1993, 115, 8167–8177.

(24) A mixture of CpV[N-2,6-C₆H₃-(*i*-Pr)₂](PMe₃)₂ (0.81 g, 1.82 mmol) and PhCH=PPh₃ (0.65 g, 1.82 mmol) in 20 mL of toluene was stirred at 70 °C for 16 h. Occasionally a vacuum was applied to remove PMe₃ liberated during reaction. After the mixture was cooled to room temperature, solid CuCl (0.18 g, 1.82 mmol) was added to coordinate free PPh₃ and the resulting mixture was stirred at room temperature for 20 h. Removal of the solvent in vacuo gave a sticky, brown residue, which was extracted two times with 50 mL of pentane. Concentrating and cooling the combined extracts to -25 °C yielded 0.41 g (1.11 mmol, 61%) of red crystals in two crops: ¹H NMR δ 13.45 (d, 1H, ³J_{PH} = 7.5 Hz, H_a), 7.65 (d, 2H, ³J_{HH} = 7.9, Ph H_o), 7.11 (t, 2H, ³J_{HH} = 7.9, Ph H_m), 7.00 (m, 3H, NAr H_m, Ph H_p), 6.83 (t, 1H, ³J_{HH} = 6.8, CHMe₂), 1.18 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 1.15 (d, 6H, ³J_{HH} = 6.8, CHMe₂), 1.18 (d, 6H, ³J_{HH} = 6.9, PMe₈); ¹³C NMR (toluene-d₈) δ 304 ($\Delta\nu_{12}$ = 600 Hz, C_a), 148.5 (NAr C_{ipso}), 144.8 (NAr C_o), 137.8 (Ph C_{ipso}), 127.8 (Ph C_o), 126.4 (Ph C_p), 125.1 (Ph C_m), 123.1 (MAr C_p), 122.5 (NAr C_m), 101.4 (Cp), 26.8 (CHMe₂), 24.4, 24.1 (CHMe₂), 18.0 (d, ¹J_{PC} = 22.5); ⁶¹V NMR δ -630 (s, $\Delta\nu_{12}$ = 1000). (25) In a 2D-NOESY of 7 an interaction between H_a and the Cp

(25) In a 2D-NOESY of 7 an interaction between H_{α} and the Cp protons is observed, whereas NOE's between this proton and the *i*-Pr protons are absent. Calculations on the molecular structure of 7 support the observed NOE: H(24)-H(20) = 2.430 Å.

(26) 7 crystallizes in the monoclinic space group $P2_1/c$ (No. 14), with a = 8.8870(12) Å, b = 15.816(2) Å, c = 18.719(2) Å, $\beta = 104.930(10)^\circ$, V = 2542.3(6) Å³, Z = 4, 8455 reflections measured, 3950 unique reflections, 150 K, Mo Ka radiation, $1.13^\circ < \theta < 24.2^\circ$. The structure was solved by automated Patterson methods (DIRDIF-92). Refinement on F^2 (SHELXL-93), with H atoms at calculated positions, except H24 (which was located on a difference Fourier map), gave wR2 = 0.131, R1 = 0.063, and S = 0.94.

(27) Cockroft, J. K.; Gibson, V. C.; Howard, J. A. K.; Poole, A. D.; Siemeling, U.; Wilson, C. J. Chem. Soc., Chem. Commun. **1992**, 1668-1670.

(28) Crystal ionic radii are V(V) = 0.59 Å and Nb(V) = 0.69 Å: Handbook of Chemistry and Physics, 60th ed.; Weast, R. C., Ed.; CRC Press: Boca Raton, FL, 1980; pp F214-215.

(29) See ref 1, p 155.



Figure 2. ORTEP diagram of 7. The thermal ellipsoids are drawn at the 30% probability level. Selected bond distances (Å): V(1)-N(1) = 1.679(5), V(1)-P(1) = 2.4010-(19), V(1)-C(24) = 1.922(6), $V(1)-Cp_{centroid} = 1.990(3)$. Selected bond angles (deg): V(1)-N(1)-C(1) = 169.5(4), P(1)-V(1)-C(24) = 87.62(17), P(1)-V(1)-N(1) = 99.24(15), N(1)-V(1)-C(24) = 105.1(2), $Cp_{centroid}-V(1)-N(1) = 126.96-(16)$, $P(1)-V(1)-Cp_{centroid} = 113.12(9)$, V(1)-C(24)-C(25) = 136.9(4), V(1)-C(24)-H(24) = 113(3).

In an initial reactivity study, benzylidene complex 7 shows no activity in ROMP of norbornene³⁰ and no reaction with acetone,³¹ presumably due to its electronic and coordinative saturation and lack of dissociation of PMe₃ from the Lewis acidic metal center.

Currently we are investigating both the reactivity of the compounds mentioned here and the scope of alkylidene transfer from phosphoranes to low-valent vanadium complexes.

Acknowledgment. This investigation was supported by the Netherlands Foundation for Chemical Research (SON) with financial aid from the Netherlands Organization for Scientific Research (NWO).

Supplementary Material Available: Text giving experimental details and analytical and spectral data for all compounds and tables giving details of the structure determination, crystal data, positional and thermal parameters, bond distances and angles, and torsion angles for 3 and 7 (38 pages). Ordering information is given on any current masthead page.

OM940361L

(31) 7 can be dissolved in acetone- d_6 without decomposition (NMR).

⁽²¹⁾ Schwartz, J.; Gell, K. I. J. Organomet. Chem. 1980, 184, C1-C2.

⁽³⁰⁾ A 0.035 M solution of 7 in C_6D_6 was allowed to react with 34 equiv of norbornene at 75 °C. No polymerization of norbornene or decomposition of 7 was observed after 72 h (NMR).