


Light-induced Reaction of Boracyclanes with Bromine in the Presence of Water. Ring Contraction of Boracyclanes to Produce Carbocyclic Structures

By YOSHINORI YAMAMOTO† and HERBERT C. BROWN*


(R. B. Wetherill Laboratory, Purdue University, Lafayette, Indiana 47905)

Summary Six-membered boracyclanes undergo ring contraction to produce the corresponding five-membered carbocyclic compounds by treatment with bromine in the presence of water.

THE light-induced α -bromination of trialkylboranes, dialkylborinic acids, and thexyldialkylboranes in the presence of water provides a convenient method for the synthesis of new carbon-carbon bonds.¹ Various types of straight-chain or branched carbon structures are now available *via* this procedure.




If bromine attacks hydrogen α to the boron atom of boracyclanes to produce the α -bromo-derivatives, and if the B-C¹ bond rearranges from boron to Cⁿ⁺¹, ring contraction of boracyclanes to achieve the formation of carbocyclic structures would be possible [equation (1)].




Accordingly, the representative boracyclanes (I), (II), and (III) were prepared *via* hydroboration of the corresponding dienes.² These boranes were treated with bromine in the presence of water under normal laboratory light, and then oxidized with alkaline hydrogen peroxide.

The six-membered boracyclanes (I) and (II) undergo ring contraction to the five-membered carbocyclic compounds. The bromination-oxidation of (I) affords cyclopentanone in 48% yield [equation (2)]. This conversion requires 2 mol. equiv. of bromine, presumably because of a competing α -bromination of the cyclopentylboronic acid.



The bromination-oxidation of (II) yields *cis*-8-hydrindanol in 60% yield [equation (3)]. The predominant formation of *cis*-isomer might be due to the preferred



formation of *cis*-9-bromo-1-thexyl-1-boradecalin in the α -bromination stage, followed by migration of the B-C² bond.

† Postdoctorate research associate supported by the National Science Foundation.

The versatility of this new development is indicated by the fact that the same borane (II) is converted by carbonylation-oxidation into pure *trans*-1-decalone.³

Five-membered boracyclanes (IIIa,b) fail to undergo this ring contraction, but in the case of (IIIa) undergo instead preferential migration of the *tert*-butyl group to produce the diol (IV) in 55% yield [equation (4)]. This result is presumably a reflection of the large ring strain involved in a contraction to the cyclobutane structure, forcing the migration to take another path. Bromination of 2-methyl-1-boracyclopentane-1-ol (IIIb) was also tried, but here also

the ring contracted product, 1-methylcyclobutanol, was not detected.

The present results provide the first example of a new type of ring contraction of cyclic boranes. A number of cyclic boranes is available from dienes and trienes *via* hydroboration.⁴ Consequently, the present results may also provide a convenient and general procedure for the synthesis of carbocyclic structures from the corresponding polyenes.

(Received, 13th August 1973; Com. 1165.)

¹ H. C. Brown and Y. Yamamoto, *Synthesis*, 1972, 699, and references cited therein.

² For preparation of (I), see H. C. Brown, E. Negishi, and S. K. Gupta, *J. Amer. Chem. Soc.*, 1970, **92**, 6648; (II), H. C. Brown and C. D. Pfaffenberger, *ibid.*, 1967, **89**, 5475; (III), H. C. Brown, E. Negishi, and P. L. Burke, *ibid.*, 1972, **94**, 3561.

³ H. C. Brown and E. Negishi, *Chem. Comm.*, 1968, 594.

⁴ H. C. Brown and E. Negishi, *J. Amer. Chem. Soc.*, 1972, **94**, 3567.