HETEROCYCLES, Vol. 80, No. 1, 2010, pp. 199 - 206. © The Japan Institute of Heterocyclic Chemistry Received, 15th July, 2009, Accepted, 18th August, 2009, Published online, 19th August, 2009 DOI: 10.3987/COM-09-S(S)55

SUBSTITUENT EFFECTS ON THE ELECTRON TRANSFER-INITIATED PHOTOCHEMICAL TRANSFORMATION OF 1,2,4-TRIAZOLE-SUBSTITUTED α -DEHYDROARYLALANINAMIDES INTO 2(1*H*)-QUINOLINONE DERIVATIVES

Yuichi Yazawa, Minoru Suzuki, Tetsutaro Igarashi, and Tadamitsu Sakurai*

Department of Material and Life Chemistry, Faculty of Engineering, Kanagawa University, Kanagawa-ku, Yokohama 221-8686, Japan

Abstract – An investigation was undertaken to elucidate substituent effects on the photoreactivity of 1,2,4-triazole-substituted α -dehydroarylalaninamides [(Z)-1] as well as on the selectivity of 2(1*H*)-quinolinone derivatives (2) from a synthetic point of view. It was found that photoinduced electron transfer-initiated cyclization of (Z)-1 bearing a *meta*-substituted phenyl or a 4-substituted naphthalen-1-yl group in methanol proceeds with a moderate to good efficiency affording the corresponding product 2 in a selectivity ranging from 33 to 100%.

Excited-state chemistry for organic molecules has continued to contribute to the development of convenient methods for synthesizing pharmaceutically useful heterocyclic compounds.¹ Particularly, photoinduced electron transfer (PET) reactions have attracted much recent attention owing to the potential that these reactions are able to construct many types of heteroatom-containing ring systems with high efficiencies.^{1a,c,d} In the course of a systematic study on the PET reaction of *N*-acyl- α -dehydroarylalanines, we found that irradiation of (*Z*)-2-(3,5-dimethyl-1,2,4-triazol-4-yl)-3-(1-naphthyl)-2-propenamide and related derivatives [hereafter, referred to as 1,2,4-triazol-substituted α -dehydro(1-naphthyl)alaninamides] in methanol-triethylamine (TEA) afforded benzo[*f*]quinolinones in high selectivities, along with minor amounts of dihydrobenzo[*f*]quinolinones.² In addition, anion radicals derived from triazole-substituted α -dehydro(1-naphthyl)alaninamides were shown to have a strong tendency to dissociate into the corresponding triazole anion and α -dehydronaphthylalanyl radicals, which readily cyclize to 2(1*H*)-benzo[*f*]quinolinones in the presence of the TEA cation radical. Although there are several synthetic studies aiming at the construction of a pharmaceutically useful quinolinone ring,³ convenient photochemical routes to quinolinone and its derivatives are scarcely known.⁴ It is, thus, of

significance to develop a novel method for synthesizing 2(1H)-quinolinone derivatives through the PETinitiated cyclization of triazole-substituted α -dehydroarylalaninamides. In this communication we synthesized (*Z*)-2-(3,5-dimethyl-1,2,4-triazol-4-yl)-3-aryl-2-propenamides [(*Z*)-1**a**–**i**] and (*Z*)-2-(3methyl-5-phenyl-1,2,4-triazol-4-yl)-3-aryl-2-propenamides [(*Z*)-1**j**–**n**] to examine substituent effects on the photoreactivity of (*Z*)-1 as well as on the selectivity of substituted 2(1H)-quinolinone as one of the (*Z*)-1-derived photoproducts (Chart 1).



Chart 1

The starting (*Z*)-isomers $[(Z)-1\mathbf{a}-\mathbf{n}]$ were prepared in good yields by the ring-opening reactions of (*Z*)-4arylmethylene-2-methyl-5(4*H*)-oxazolones with equimolar amounts of the corresponding *N*-substituted hydrazides in acetonitrile, followed by the condensation of the resulting 1,2,4-triazole-substituted 2propenoic acids with methyl amine in 1,4-dioxane or dimethyl formamide.⁵ After a nitrogen-saturated methanol solution of (*Z*)-1**a** (4.0×10^{-3} mol dm⁻³, 500 mL) containing TEA (0.10 mol dm⁻³) was irradiated at wavelengths longer than 280 nm from a 400 W high-pressure Hg lamp for 60 min at room temperature (Pyrex glass filter, conversion 95%), the reaction mixture was subjected to preparative thin layer chromatography over silica gel (eluent: EtOAc). Usual workup allowed us to isolate 1-methyl-6methoxy-2(1*H*)-quinolinone (**2a**; isolated yield, 20%), the regional isomer of which [1-methyl-8methoxy-2(1*H*)-quinolinone] was not detected.⁶ On the other hand, a comparison of the ¹H NMR spectrum of this reaction mixture with those of 3,4-dihydro-3-(3,5-dimethyl-1,2,4-triazol-4-yl)-1-methyl-2(1*H*)-benzo[*f*]quinolinone (**3a**) and triazole derivatives were produced along with (*E*)-**1a** and **4a** (Scheme 1).² The arylalaninamide derivative **4a** was isolated by repeated preparative thin layer



Table 1. Substituent effects on the conversion and composition of each compound obtained by the 30 min irradiation of (*Z*)-1a-h in MeOH–TEA at room temperature^a

(<i>Z</i>)-1	Conversion (%)	Composition (%)					Selectivity of
		<i>(Z)</i> -1	(E)- 1	2	3	4	2 (%) ^b
1a	24	68	8	8	14	2	33
1b	6	77	17	6	0	0	100
1c	0	36	64	0	0	0	0
1d	32	34	34	20	12	0	63
1e	0	77	23	0	0	0	0
1f	17	71	11	2	15	0	12
1g	48	27	25	8	40	0	17
1h	0	59	41	0	0	0	0

 $a[(Z)-1a-h] = 4.0 \times 10^{-3} \text{ mol } dm^{-3}; [TEA] = 0.10 \text{ mol } dm^{-3}.$

^bSelectivity of **2** was estimated by dividing the composition for **2** by the sum of composition for **2**, **3**, and **4**.

chromatography and the ¹H NMR spectral data substantiated the structure of this reduction product.⁷ The result that the photocyclization reaction of (*Z*)-1a proceeds without forming any byproducts enabled us to monitor this reaction by means of ¹H NMR spectroscopy (Scheme 1). However, prolonged irradiation (>60 min) resulted in a side reaction to a definitely detectable extent depending on the structure of (*Z*)-1a–h and, hence, the ¹H NMR composition of each compound was determined and compared after 30 min irradiation to analyze substituent effects on the photoreactivity of 1 and the selectivity of 2 (Table 1).

The data collected in Table 1 demonstrate that substituents introduced into the benzene ring of (*Z*)-1e exert great electronic and steric effects (which are beyond our expectations) on both the photoreactivity and the product composition. Replacement of hydrogens at the *para-* and *meta-*positions on the ring by methoxy or chloro groups increases the conversion of 1 but replacement at the latter position lowers the composition ratio of 2 to 3. In addition, the presence of a bulky methoxy group at the *ortho-*position (1c) or an electron-withdrawing cyano group at the *meta-*position (1h) resulted in a considerable decrease in photoreactivity. The finding that irradiation of (*Z*)-1i under the same conditions enables the complete conversion into the corresponding photoproducts 2i (isolated yield, 72%; selectivity, 76%) and 3i (R^4 = H, R^5 = Me in Scheme 2) confirms that the phenyl moiety in 1e has a much lower ability to accept an electron from TEA in its excited state, as compared to the naphthyl in 1i. As mentioned above, either a methoxy or a chloro substituent introduced at the *meta-*position on the benzene ring of (*Z*)-1e has a tendency to enhance the electron-accepting ability of this ring to exhibit the so-called "*meta* effect" although the selectivity of the quinolinone derivative 2 is decreased by this effect.⁸



In a previous study we found (through analysis of the effects of substituents attached to the triazole ring on the product composition) that the phenyl substituent increases the composition ratio of 2 to 3 by a factor of about 2 with keeping the conversion of 1 nearly constant.² Additionally, taking into account

the fact that methoxy and benzo groups increase both the photoreactivity of 1 and the selectivity of 2, we synthesized (Z)-1j-n to explore the *meta* effect as well as the effect of substituent (attached to the naphthalene ring) on the reactivity and selectivity. In Table 2 are summarized the conversion of 1 and the product composition, obtained under the same irradiation conditions as those for (Z)-1a-i. Replacement of one of the two methyl groups attached to the triazole ring by the phenyl increased the selectivity of 2 from 74% (2i) to 84% (2m), being consistent with our previous results $(76\% \rightarrow 86\%)$.² Interestingly, substitution of two methoxy groups for the *meta*-hydrogens (1) enhanced not only the photoreactivity (conversion) of **1a** ($24\% \rightarrow 79\%$) but also the selectivity of **2a** ($33\% \rightarrow 71\%$). However, the presence of the fluoro-substituted naphthalene ring in 1n (R⁴= F, R⁵= Ph in Scheme 2) fairly reduced both the selectivity of the benzoquinolinone derivative 2m (84% for 2m and 40% for 2n) and the photoreactivity of 1m (R⁴= H, R⁵= Ph). Furthermore, the introduction of an electron-donating methyl group (11) exerted only a very minor effect on these two parameters but replacement of this methyl group by a methoxy (1k) substantially reduced the conversion of 1m (100% for 1m and 13% for 1k) with the selective formation of 2k. As seen from these considerations, the selectivity of the benzoquinolinone derivative 2 has a clear tendency to increase with an increase in the electron-donating ability of the substituent R^4 in (Z)-1k-n, whereas this substituent has a complicated influence on the excited-state reactivity of the triazole-substituted 1-naphthylalaninamide derivative 1.

(<i>Z</i>)-1	Conversion (%)	Composition (%)					Selectivity of
		<i>(Z)</i> -1	(E)- 1	2	3	4	2 (%) ^b
1j	79	16	5	56	20	3	71
1k	13	33	54	13	0	0	100
11	100	0	0	90	10	0	90
1m	100	0	0	84	16	0	84
1n	48	21	31	19	29	0	40

Table 2. Substituent effects on the conversion and composition of each compound obtained by the 30 min irradiation of (Z)-1j–n in MeOH–TEA at room temperature^a

 $a[(Z)-1j-n] = 4.0 \times 10^{-3} \text{ mol } \text{dm}^{-3}; [TEA] = 0.10 \text{ mol } \text{dm}^{-3}.$

^bSelectivity of **2** was estimated by dividing the composition for **2** by the sum of composition for **2**, **3**, and **4**.

We previously proposed (through a detailed analysis of solvent and substituent effects on the composition ratio of **2i** to **3i**) that the relative stability of the (*E*)-**1i**-derived anion radical intermediates (*E*)-**IA** and (*E*)-**IB** (which are in equilibrium with each other) determines the magnitude of this ratio (Scheme 3).² Since electron-donating substituents such as methoxy and methyl is considered to shift the equilibrium to the



Scheme 3

(*E*)-**IB** side by destabilizing the former intermediate, the finding that these substituents tend to enhance the selectivity of **2** provides additional evidence in support of our proposal. Taking into account that the photocyclization eventually yielding **2** and **3** is initiated by ET from TEA to the excited-state (*E*)-**1**, we propose that the photoreactivity of this (*E*)-isomer may be determined by the relative rate for ET to the (*E*)-isomer as well as by the relative rate for back ET from (*E*)-**IA** and (*E*)-**IB** to the TEA cation radical. It is thus likely that substituents introduced into the benzene or naphthalene ring of **1** exert their electronic and steric effects on these ET and back ET processes in a complicated manner. Negligible formation of the cyano-substituted products **2h** and **3h** may be due to either the exclusive deactivation of the excitedstate (*E*)-**1h** or the exclusive back ET in (*E*)-**IA**. On the other hand, it seems strange that **1a** and **1j** underwent PET-initiated reduction to give **4a** and **4j** as minor products (Schemes 1 and 3). The *meta* effect of electron-donating methoxy groups in these two α -dehydroarylalanines might be responsible for the increased reactivity of the anion radical intermediate (*E*)-**IA** toward hydrogen abstraction from methanol.

As described above, the PET-initiated cyclization reactions of (*Z*)-1 bearing a methoxy group attached at the *meta*- or the *para*-position on the benzene ring (as well as an electron-donating or an electron-attracting group introduced into the naphthalene ring) were found to proceed with moderate to good efficiencies affording the corresponding quinolinone derivatives **2** in 33–100% selectivities. Taking the relatively high photostability of these derivatives into account, we may conclude that the photocyclization reactions provide a novel synthetic method for the construction of several substituted 2(1H)-quinolinone rings although the introduction of an electron-attracting group into the benzene or the naphthalene ring renders this method less attractive.

205

ACKNOWLEDGMENTS

This research was partially supported by a "Scientific Frontier Research Project" from the Ministry of Education, Sports, Culture, Science and Technology, Japan.

REFERENCES AND NOTES

- a) 'Synthetic Organic Photochemistry,' ed. by W. M. Horspool, Plenum, New York, 1984; b) 'Handbook of Photochemistry and Photobiology,' Vol. 2, ed. by H. S. Nalwa, American Scientific Publishers, Stevenson Ranch, 2003; c) 'Synthetic Organic Photochemistry,' ed. by A. G. Griesbeck and J. Mattay, Marcel Dekker, New York, 2005; d) A. G. Griesbeck, N. Hoffmann, and K. Warzecha, *Acc. Chem. Res.*, 2007, 40, 128.
- 2. K. Maekawa, A. Tomoda, T. Igarashi, and T. Sakurai, *Heterocycles*, 2009, 77, 739.
- O. Meth-Cohn, *Heterocycles*, 1993, **35**, 539; C. W. Holzapfel and C. Dwyer, *Heterocycles*, 1998, **48**, 215; K. Y. Lee and J. N. Kim, *Bull. Korean Chem. Soc.*, 2002, **23**, 939; B. Joseph, F. Darro, A. Behard, B. Lesur, F. Collignon, C. Decaestecker, A. Frydman, G. Guillaumet, and R. Kiss, *J. Med. Chem.*, 2002, **45**, 2543; D. V. Kadnikov and R. C. Larock, *J. Organomet. Chem.*, 2003, **687**, 425; M. Hadjeri, C.Beney, and A. Boumendjel, *Curr. Org. Chem.*, 2003, **7**, 679; S. K. Chattopadhyay, R. Dey, and S. Biswas, *Synthesis*, 2005, 403; Q. Li, K. W. Woods, W. Wang, N.-H. Lin, A. Claiborne, W.-Z. Gu, J. Cohen, V. S. Stoll, C. Hutchins, D. Frost, S. H. Rosenberg, and H. L. Sham, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2033; M. Asahara, M. Ohtsutsumi, M. Tamura, N. Nishiwaki, and M. Ariga, *Bull. Chem. Soc. Jpn.*, 2005, **78**, 2235; R. Bernini, S. Cacchi, G. Fabrizi, *Synlett*, 2006, 2947.
- 4. T. Horaguchi, N. Hosokawa, K. Tanemura, and T. Suzuki, J. Heterocycl. Chem., 2002, 39, 61.
- 5. Selected data for (Z)-1a: IR (KBr) ν/cm⁻¹= 3232, 1666; ¹H NMR (600 MHz, DMSO-d₆) δ= 2.03 (6H, s), 2.70 (3H, d, J= 4.6 Hz), 3.60 (3H, s), 6.18 (1H, s), 6.44 (1H, d, J= 8.0 Hz), 6.95 (1H, d, J= 7.4 Hz), 7.25 (1H, dd, J= 7.4, 8.0 Hz), 7.78 (1H, s), 8.24 (1H, q, J= 4.6 Hz); ¹³C NMR (150 MHz, DMSO-d₆) δ= 10.8 (2C), 27.1, 55.5, 114.0, 117.2, 122.4, 125.9, 130.9, 133.5, 135.6, 150.6, 160.0, 163.3.
- 6. Selected data for 2a: IR (KBr) ν/cm⁻¹= 1678; ¹H NMR (600 MHz, DMSO-d₆) δ= 3.56 (3H, s), 3.78 (3H, s), 6.58 (1H, d, J= 9.7 Hz), 7.20 (1H, dd, J= 2.8, 9.7 Hz), 7.23 (1H, d, J= 2.8 Hz), 7.40 (1H, d, J= 9.2 Hz), 7.79 (1H, d, J= 9.2 Hz); ¹³C NMR (150 MHz, DMSO-d₆) δ= 29.0, 55.5, 110.5, 115.9, 119.0, 120.8, 121.6, 134.2, 138.6, 154.1, 154.8, 160.6.
- Selected data for 4a: ¹H NMR (600 MHz, DMSO-*d*₆) *δ*= 2.05 (6H, s), 2.63 (3H, d, *J*= 4.6 Hz), 2.99 (1H, dd, *J*= 11.6, 14.3 Hz), 3.38 (1H, dd, *J*= 4.6, 14.3 Hz), 3.63 (3H, s), 4.95 (1H, dd, *J*= 4.6, 11.6 Hz), 6.56 (1H, s), 6.57 (1H, d, *J*= 7.4 Hz), 6.73 (1H, d, *J*= 8.0 Hz), 7.10 (1H, dd, *J*= 7.4, 8.0 Hz), 8.06 (1H,

q, *J*= 4.6 Hz).

 H. E. Zimmerman and V. R. Sandel, J. Am. Chem. Soc., 1963, 85, 915; H. E. Zimmerman, J. Am. Chem. Soc., 1995, 117, 8988; H. E. Zimmerman, J. Phys. Chem. A, 1998, 102, 5616.