This article was downloaded by: [RMIT University] On: 23 July 2013, At: 06:42 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



# Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/uopp20

# IMPROVED SYNTHESES OF 20- AND 26-MEMBERED bis(5-CARBOMETHOXY-1,3-PHENYLENE) CROWN ETHERS

Devdatt S. Nagvekar<sup>a</sup> & Harry W. Gibson<sup>a</sup>

<sup>a</sup> Department of Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, VA, 24061 Published online: 09 Feb 2009.

To cite this article: Devdatt S. Nagvekar & Harry W. Gibson (1997) IMPROVED SYNTHESES OF 20-AND 26-MEMBERED bis(5-CARBOMETHOXY-1,3-PHENYLENE) CROWN ETHERS, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 29:2, 237-240, DOI: <u>10.1080/00304949709355194</u>

To link to this article: http://dx.doi.org/10.1080/00304949709355194

# PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <a href="http://www.tandfonline.com/page/terms-and-conditions">http://www.tandfonline.com/page/terms-and-conditions</a>

### **IMPROVED SYNTHESES OF 20- AND 26-MEMBERED**

## bis(5-CARBOMETHOXY-1,3-PHENYLENE) CROWN ETHERS

Submitted by (6/03/96)

Devdatt S. Nagvekar and Harry W. Gibson<sup>\*</sup>

Department of Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061

Recently a number of useful functionalized *bis(m*-phenylene) crown ethers were synthesized in a one step reaction of 5-substituted resorcinols with linear dihalides in DMF,<sup>1-6</sup> but this method gave relatively poor yields. Controlling the rates of *intra*- versus *inter*-molecular reactions during the condensation is a major problem in any single step cyclization,<sup>7,8</sup> although a number of other factors are known to affect yields.<sup>8</sup> Taking advantage of the preorganization effect, we achieved improved syntheses of these macrocycles *via* a two-step methodology.

The disodium salt of methyl 3,5-dihydroxybenzoate (1) in DMF was added slowly to a large excess of oligo(ethylene glycol) dichloride (2 or 3), which also acted as solvent. This reaction gave intermediate dichlorides 4 and 5 in 81 and 71% yields, respectively, after purification *via* column chromatography. Cyclization was accomplished with alkali metal salts in the presence of the phase transfer agent n-Bu<sub>4</sub>NI. It is well known that crown ether yields are greatly dependent on the nature of the metal ion<sup>9</sup> and the ring size of the macrocycle, in addition to concentration.<sup>10</sup> The yields of macrocycles 6 and 7 were higher *via* the two-step method (Table). In the two-step method K<sub>2</sub>CO<sub>3</sub> gave the highest yield for both macrocycles, much higher than NaH or CsF. Macrocycles 6 and 7 were purified by column chromatography.



The hydrolysis of bis(5-carbomethoxy-1,3-phenylene)-26-crown-8 (7) was achieved with LiOH in THF, affording the diacid 8. The syntheses of polyesters and polyamides from the macrocyclic diesters and diacids are currently under investigation and will be reported elsewhere.

Method	Base	Temp. (°C)	Yield of <b>6</b> (%)	Yield of <b>7</b> (%)
One-step	NaH	85	7.0ª	6.0 <sup>c</sup>
One-step	KH	85	18.0 <sup>a</sup>	
One-step	Cs <sub>2</sub> CO <sub>3</sub>	85	4.5 <sup>a</sup>	_
Two-step	NaH	110	16.0 <sup>b</sup>	
Two-step	K <sub>2</sub> CO <sub>3</sub>	110	58.0 <sup>b</sup>	46.0 <sup>d</sup>
Two-step	ĊsF	110	26.0 <sup>b</sup>	26.0 <sup>d</sup>

TABLE. Yields of Macrocycles 6 and 7 by One-step and Two-step Methods with Various Bases in DMF

a) One-step reaction<sup>6</sup> of 1 and 2, 0.12 M each. b) Reaction of 1 and 4 under pseudo-high dilution with n-Bu<sub>4</sub>NI as catalyst. c) One-step reaction<sup>6</sup> of 1 and 3, 0.12 M each. d) Reaction of 1 and 5 under pseudo-high dilution with n-Bu<sub>4</sub>NI as catalyst.

## **EXPERIMENTAL SECTION**

Methyl 3,5-dihydroxybenzoate was synthesized according to the literature procedure.<sup>1</sup> Melting points were taken in capillary tubes with a Haake-Buchler apparatus and have been corrected. NMR spectra were obtained at 20° on a Varian Unity 400 MHz spectrometer using CDCl<sub>3</sub> unless otherwise noted and TMS as internal standard. Infrared spectra were recorded on a Nicolet MX-1 FTIR spectrometer. A Harvard syringe infusion pump. Model 22 was used to control the addition rates in the cyclization reactions. Mass spectroscopy was carried out at the Mass Spectrometry Center at the University of Nebraska-Lincoln. Elemental analysis was performed by Atlantic Microlabs of Norcross, GA.

Methyl 3,5-*bis*(5-Chloro-3-oxapentyloxy)benzoate (4).- Sodium hydride (2.92 g, 73 mmol, 60% in mineral oil) was added to a solution of methyl 3,5-dihydroxybenzoate (1, 6.06 g, 36 mmol) in DMF (30 mL). The mixture was stirred for 2 hrs at 110° and cooled to 25°; the resulting suspension of dianion was added to a solution of di(ethylene glycol) dichloride (2, 51.81 g, 362.3 mmol, 10 equiv.) in DMF (30 mL) over 6 hrs and then the mixture was stirred for five days at 50° and filtered to remove NaCl. DMF and excess dichloride 2 were removed *in vacuo*. Purification of the residue by flash column chromatography with 20:1 CHCl<sub>3</sub>:EtOAc gave 11.1 g (81%) of 4, an oil; IR (neat): 3100 (=CH), 2951 (-CH), 1716 (C=O), 1601 (C=C), 1126 (C-O-C); <sup>1</sup>H NMR:  $\delta$  7.21 (d, J = 2.2Hz, 2H), 6.53 (t, J = 2.2Hz, 1H), 4.16 (t, J = 4.6Hz, 4H), 3.88 (m, 7H), 3.83 (t, J = 5.8Hz, 4H), 3.66 (t, J = 5.8Hz, 4H); <sup>13</sup>C NMR:  $\delta$  42.64, 52.18, 67.67, 69.53, 71.45, 106.86, 108.02, 131.88, 159.56, 166.59; MS (EI) m/z: 380 (M<sup>+</sup>, 3.8%), 349 [(M-OCH<sub>3</sub>)<sup>+</sup>, 3.5%], 274 [(M-C<sub>4</sub>H<sub>7</sub>ClO)<sup>+</sup>, 5.3%], 106 [(M-C<sub>12</sub>H<sub>15</sub>ClO<sub>5</sub>)<sup>+</sup>, 53%], 63 (C<sub>2</sub>H<sub>4</sub>Cl<sup>+</sup>, 100%).

Methyl 3,5-*bis*(8-Chloro-3,6-dioxaoctyloxy)benzoate (5).- The above procedure with 1 and 3 on a similar scale produced (71%) 5, an oil: IR (neat): 3090 (=CH), 2884 (-CH), 1722 (C=O), 1596 (C=C), 1118 (C-O-C); <sup>1</sup>H NMR: δ 7.20 (d, J = 2.2Hz, 2H), 6.70 (t, J = 2.2Hz, 1H), 4.15 (t, J = 4.6Hz, 4H), 3.90 (s, 3H), 3.87 (t, J = 4.6Hz, 4H), 3.77 (t, J = 5.8Hz, 4H), 3.73 (m, 8H), 3.64 (t, J = 5.8Hz, 4H); <sup>13</sup>C NMR: δ 42.68, 52.19, 67.68, 69.60, 70.64, 70.77, 71.34, 106.87, 107.99, 131.84, 159.67, 166.70; MS (FAB) m/z: 469 [(M+H)<sup>+</sup>, 57%] and 437 [(M-OCH<sub>3</sub>)<sup>+</sup>, 100%].

*bis*(5-Carbomethoxy-1,3-phenylene)-20-crown-6 (6).- A solution of methyl 3,5-*bis*(5-chloro-3-oxapentyloxy)benzoate (4, 6.04 g, 15.8 mmol) and methyl 3,5-dihydroxybenzoate (1, 2.66 g, 15.8 mmol) in DMF (total volume 22 mL) was added *via* a syringe pump at 0.75 mL/hr to a suspension containing  $K_2CO_3$  (22.04 g, 159.5 mmol) and n-Bu<sub>4</sub>NI (20 mg) in DMF (750 mL) at 110°. After complete addition, the mixture was stirred vigorously at 110° for five days, cooled and evaporated. The residue in CH<sub>2</sub>Cl<sub>2</sub> was filtered to remove salts and subjected to flash column chromatography using Et<sub>2</sub>O to give 4.36 g (58%) of pure **6**, mp. 185.9-187.5°, lit.<sup>6</sup> 179-181°; <sup>1</sup>H NMR:  $\delta$  7.09 (d, J = 2.4Hz, 4H), 6.67 (br s, 2H), 4.12 (br s, 8H), 3.83 (br s, 14H); <sup>13</sup>C NMR:  $\delta$  52.16, 68.03, 69.72, 108.27, 131.73, 159.68, 166.68.

*bis*(5-Carbomethoxy-1,3-phenylene)-26-crown-8 (7).- The above procedure with 1 and 5 on a similar scale produced (46%) pure 7: mp. 131.4-132.9°, lit.<sup>6</sup> 131.2-133.2°; <sup>1</sup>H NMR:  $\delta$  7.16 (d, J = 2.2Hz, 4H), 6.68 (t, J = 2.2Hz, 2H), 4.11 (t, J = 4.6Hz, 8H), 3.87 (s, 6H), 3.85 (t, J = 4.6Hz, 8H), 3.74 (s, 8H); <sup>13</sup>C NMR:  $\delta$  52.16, 67.65, 69.50, 70.92, 107.06, 107.86, 131.74, 159.71, 166.73.

*bis*(5-Carboxy-1,3-phenylene)-26-crown-8 (8).- A solution of LiOH+ $H_2O$  (0.764g, 19.1 mmol), *bis*(5-carbomethoxy-1,3-phenylene)-26-crown-8 (7, 1.35 g, 2.39 mmol), water (10 mL) and THF (60 mL) was refluxed for 4 days. The solvent was removed and the residue was treated with  $H_2O$  (30 mL), and then filtered. The filtrate was acidified with 2N HCl to give a white precipitate. Filtration followed by drying gave 1.20 g (94%) of 8, mp. 259.1-260.8°; <sup>1</sup>H NMR (DMSO- $d_6$ ):  $\delta$  7.05 (d, J = 1.6Hz, 4H), 6.71 (s, 2H), 4.10 (brs, 8H), 3.73 (br s, 8H), 3.60 (s, 8H); <sup>13</sup>C NMR (DMSO- $d_6$ ):  $\delta$  67.45, 68.72, 70.03, 106.29, 107.12, 132.85, 159.49, 166.93.

Anal. Calcd. for C<sub>26</sub>H<sub>32</sub>O<sub>12</sub>: C, 58.21; H, 6.01. Found: 58.24; H, 6.04

Acknowledgement.- We acknowledge support from the National Science Foundation (DMR-93-20196) for this work and the Center for Mass Spectrometry at the University of Nebraska-Lincoln for mass spectrometry.

#### REFERENCES

- 1. Y. Delaviz and H. W. Gibson, Macromolecules, 25, 18 (1992).
- 2. H. W. Gibson and Y. Delaviz, Polymer, 35, 1109 (1994).
- 3. Y. Delaviz and H. W. Gibson, Org. Prep. Proced. Int., 23, 382 (1991).
- 4. Y. Delaviz and H. W. Gibson, Polym. Commun., 32, 103 (1991).
- 5. Y. Delaviz, Y. X. Shen and H. W. Gibson, *ibid.*, **32**, 21 (1991).
- 6. Y. Delaviz, J. S. Merola, M. A. G. Berg and H. W. Gibson, J. Org. Chem., 60, 516, (1995).
- a) A. J. Kirby, Adv. Phys. Org. Chem., 17, 1831 (1980); b) D. A. Laidler and J. F. Stoddart in "Crown Ethers and Analogs"; S. Patai and Z. Rappoport, eds., John Wiley & Sons, New York, 1989, pp. 1-58.

- a) B. Dietrich, P. Viout and J.-M. Lehn, "Macrocyclic Chemistry", VCH Publishers, New York, 1993, pp. 1-165; b) "Frontiers in Supramolecular Organic Chemistry and Photochemistry", H.-J. Schneider and H. Durr, eds., VCH Publishers, New York, 1991; c) J.-M. Lehn, "Supramolecular Chemistry", VCH Publishers, Inc., New York, 1995.
- 9. G. Dijkstra, W. H. Kruizinga and R. M. Kellogg, J. Org. Chem., 52, 4230 (1987).
- a) P. R. Story and P. Bush, in "Advances in Organic Chemistry", C. E. Taylor, ed., Wiley Interscience, New York, 1972, vol. 8, pp. 67-95; b) P. Knops, N. Sendhoff, H.-B. Mekelburger and F. Vögtle, Topics Curr. Chem., 161, 3 (1992); c) A. Ostrowicki, E. Koepp and F. Vögtle, *ibid.*, 161, 39 (1992).

### 5-BENZYLOXYRESORCINOL, A MONOPROTECTED PHLOROGLUCINOL

\*\*\*\*\*\*\*

Submitted by (6/07/96)

Devdatt S. Nagvekar and Harry W. Gibson\*

Department of Chemistry, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061

5-Benzyloxyresorcinol (6) is a key intermediate in our synthetic schemes directed toward diand mono-functionalized bis(m-phenylene) crown ethers<sup>1-7</sup> and related cryptands. The previous onestep synthesis of 5-benzyloxyresorcinol (6) from phloroglucinol (1) gave only a 9% yield.<sup>8</sup> Although our attempts to improve this method failed, herein we report an alternative synthesis that affords an improved overall yield of the target compound.

Treatment of phloroglucinol (1) with two moles of benzoyl chloride produced a mixture of benzoates 2, 3 and 4. The monoester 4 was insoluble in  $CHCl_3$  and hence could be easily removed from the mixture, since both 2 and 3 were soluble. Analysis of the soluble mixture *via* <sup>1</sup>H NMR spectroscopy revealed a 36:64 ratio of 2: 3. Pure 3 was prepared as a reference material from pure 2<sup>9</sup> by hydrolysis with  $Cs_2CO_3$ . It was also converted to the desired 6 (34% overall), but this route was less efficient than the one described here. The mixture of 2 and 3, without further purification, was treated with NaH and one equivalent of benzyl bromide to give a mixture of 2 and 5, which was hydrolyzed with aqueous methanolic KOH. Neutralization with acid, followed by removal of benzoic acid with saturated aqueous NaHCO<sub>3</sub> and removal of phloroglucinol (1) derived from 2 by washing with water afforded the desired 5-benzyloxyresorcinol (6) in 41% overall yield.

The application of 5-benzyloxyresorcinol (6) to the synthesis of monomeric macrocycles, cryptands and polymacrocycles is the subject of current research and will be reported elsewhere. We note that Büchi reported the dibenzyl ether and the monobenzenesulfonate of phloroglucinol.<sup>10</sup>