

Li[Cp₂Zr(C≡CPh)(η²:1,2-PhC₂C≡CPh)]: an anionic zirconium(II) intermediate for carbon–carbon coupling

Robert Choukroun,* Jianshe Zhao, Christian Lorber, Patrick Cassoux and Bruno Donnadieu


Equipe Précurseurs Moléculaires et Matériaux, Laboratoire de Chimie de Coordination du CNRS, 205 Route de Narbonne, 31077, Toulouse Cedex, France. E-mail: choukrou@lcc-toulouse.fr

Received (in Basel, Switzerland) 5th June 2000, Accepted 5th July 2000

Published on the Web 24th July 2000

The unexpected anionic Zr^{II} complex [Cp₂Zr(C≡CPh)(η²:1,2-PhC₂C≡CPh)][−] was isolated from LiC≡CPh and Cp₂ZrCl₂ in THF, giving clear evidence for a CC coupling between two alkynyl moieties, one of them being η²-bonded to the zirconium atom.

New prospects towards non linear optical (NLO) materials have induced, in recent years, a resurgence of interest in the chemistry of acetylenic transition metals complexes.¹ Numerous studies have focused on the CC coupling of the acetylenic moieties, which underlies the synthesis of the building block of the poly-ynes.² We have been interested in the reaction of vanadocene with poly-ynes.³ An unexpected heterodimetallic complex Cp₂V(μ:η²:η⁴-PhC≡CC≡CPh)ZrCp₂, containing a butadiene framework with two internal planar tetracoordinate carbons was isolated from Cp₂V and Cp₂Zr(C≡CPh)₂ (Cp' = C₅H₅, C₅H₄Me, C₅H₄Bu^t, C₅H₄SiMe₃).^{4,5} The chemistry of these bis(alkynyl)metallocene precursors has previously been detailed^{6–8} and their synthesis, generally in diethyl ether solvent, well described.⁶ On the other hand, a CC bond forming reaction was reported by Negishi *et al.*⁹ as resulting from the reaction of Cp₂ZrCl₂ with 3 equiv. of LiC≡CPh in THF followed by hydrolysis affording the isomerically pure (Z)-1,4-diphenylbut-1-ene-3-yne. In this process, Li[Cp₂Zr(C≡CPh)₃] was postulated as an intermediate species.⁹ The authors underline the necessity of the third equiv. of LiC≡CPh to Cp₂Zr(C≡CPh)₂ to produce the CC coupling. More recently, the [Zr(C≡CR)₃][−] anion moiety was suggested as being an intermediate species in the trimerization of *tert*-butyl acetylene to 1,3,6-tri(*tert*-butyl)fulvene.¹⁰ We report here the X-ray structure of the intermediate anion species Li[Cp₂Zr(C≡CPh)(η²:1,2-PhC₂C≡CPh)] and some aspects of its formation mechanism.



When the reaction of Cp₂ZrCl₂ with 2 equiv. of LiC≡CPh is carried out in THF, followed by slow diffusion of pentane, the unexpected Zr^{II} lithium salt species Li[Cp₂Zr(C≡CPh)(η²:1,2-PhC₂C≡CPh)][−] 1,† was obtained as a red crystalline complex and fully characterized by an X-ray structure determination (Fig. 1).‡ The main feature of this structure gives clear evidence for a CC coupling between two alkynyl moieties, one of them being η²-bonded to the zirconium atom. A titanium complex related to 1, namely (C₅Me₅)₂Ti(η²:1,2-RC₂C≡CR)] was also recently isolated by Rosenthal *et al.* by Mg reduction of (C₅Me₅)₂TiCl₂ in the presence of Me₃SiC₄SiMe₃.^{11a} Depending on the ratio of the reactants, this reaction also affords either the tweezer Ti^{III} complex [(C₅Me₅)₂Ti(η¹-C≡CSiMe₃)₂][Mg(THF)Cl)] (A) or

[(C₅Me₅)₂Ti(η³-Me₃SiC₃=C(C≡CSiMe₃)SiMe₃)]^{11b} (B). The tweezer Ti^{III} complex (A) could be described as an intermediate for the formation of complex (B).

Different crossing reactions were monitored by ¹H NMR to gain an understanding of the formation of 1.§ When the reaction of Cp₂ZrCl₂ and 2 equiv. of LiC≡CPh is carried out in THF and in the absence of sunlight, Cp₂Zr(C≡CPh)₂ 2 was obtained as the sole product and can be kept unchanged at least one week in the dark, whereas in presence of daylight only 1 is formed.¹² Starting from 2 and the solid lithium salt LiC≡CPh in THF-d₈, complete consumption of the Li salt gives 1 nearly immediately and quantitatively in absence or in presence of daylight. We checked that in the absence of daylight no reaction occurs in C₆D₆ between 2 and LiC≡CPh (1:1) whereas still in the absence of daylight the addition of THF-d₈ in the NMR tube, which dissolves the lithium salt, immediately generates 1.

Complementary hydrolysis experiments on complex 1 containing the preformed CC coupling were performed to ensure that the 1,4-diphenylbut-1-en-3-yne is also formed in this case. In presence of HCl and at room temperature, hydrolysis of 1 leads to the formation of *E* and *Z* isomers of the enyne PhCH=CHC≡CPh with a high selectivity when the reaction is carried out in toluene (toluene: *Z*:*E* = 98:2; THF: *Z*:*E* = 30:70).¶ Adding LiC≡CBu^t to 2 followed by HCl hydrolysis gives *Z* enynes, namely PhCH=CHC≡CBu^t, PhC≡CCH=CHBu^t and PhCHCHC≡CPh (roughly 30, 15, 55% respectively, characterized by GC/MS and ¹H NMR).|| This result suggests that the first step of the reaction is the formation of the




Fig. 1 Molecular structure of anionic [1][−] with selected bond distances (Å) and angles (°), hydrogen atoms omitted. Zr–C(11) 2.206(2), Zr–C(12) 2.342(1), Zr–C(15) 2.314(2), C(11)–C(12) 1.335(2), C(12)–C(13) 1.412(2), C(13)–C(14) 1.208(2), C(15)–C(16) 1.223(2), Zr–Cp 2.252(av.); Zr–C(11)–C(111) 147.8(1), Zr–C(15)–C(16) 171.9(1), Zr–C(12)–C(13) 127.9(1), C(12)–C(13)–C(14) 177.5(2), C(11)–Zr–C(12) 33.97(6), C(12)–Zr–C(15) 88.60(5), C(11)–Zr–C(15) 122.53(5), Zr–C(15)–C(16) 171.9(1), C(15)–C(16)–C(166) 176.3(2), Cp–Zr–Cp 129.57(av.) [Cp are the centroids of the C₅H₅ rings C(1)–C(5), C(6)–C(10)].



At this stage we are not in a position to prove the involvement of $\text{LiC}\equiv\text{CPh}$ in a photoassisted reaction with **2** leading to **1**. However, it is noteworthy that when the reaction is carried out in the dark, it yields only **2**. Our results clearly suggest that the alkynyl coupling reaction is induced by a third alkynyl ligand *via* the formation of the unstable ‘ate’ intermediate Zr^{IV} species $\text{Li}[\text{Cp}_2\text{Zr}(\text{C}\equiv\text{CPh})_3]$, or an assumed tweezer Zr species $[\text{Cp}_2\text{Zr}(\text{C}\equiv\text{CPh})_2][\text{Li}(\text{C}\equiv\text{CPh})]$, which may subsequently rearrange to **1**.¹⁶

Notes and references

§ A suspension of Cp_2ZrCl_2 (0.900 g, 3.08 mmol) was treated with 2 equiv. solid $\text{LiC}\equiv\text{CPh}$ (0.665 g, 6.16 mmol) in benzene for 4 h and species such as Cp_2ZrCl_2 , $\text{Cp}_2\text{ZrCl}(\text{C}\equiv\text{CPh})$ and **2** were identified by ^1H NMR (in nearly 1:4:1 ratio respectively). After 24 h stirring and work-up, **2** was obtained as a crystalline solid (0.840 g, 64% yield). With 3 equiv. $\text{LiC}\equiv\text{CPh}$ for 24 h,

†† ^1H and $^{13}\text{C}\{^1\text{H}\}$ NMR of the reaction show complex spectra in which three main cyclopentadienyl signals can be observed at 5.76, 5.71, 5.67/105.0, 104.9, 104.7 ppm; low field quaternary carbons at 228, 225.9, 208.2, 205.9, 203, 202 ppm were also observed.

- 1 H. S. Nalwa and S. Miyata, *Nonlinear Optics of Organic Molecules and Polymers*, CRC Press, Boca Raton, FL, 1997.
- 2 (a) A. de Meijeire, *Top. Curr. Chem.*, 1999, **201**, 000; (b) F. Diederich, in *Modern Acetylene Chemistry*, ed. P. J. Stang and F. Diederich, VCH, Weinheim, 1995, p. 443; (c) F. Diederich and Y. Rubin, *Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 1101.
- 3 (a) R. Choukroun, C. Lorber, B. Donnadiou, B. Henner, R. Frantz and C. Guerin, *Chem. Commun.*, 1999, 1099 and references therein; (b) R. Choukroun, B. Donnadiou, I. Malfant, S. Haubrich, R. Frantz, C. Guerin and B. Henner, *Chem. Commun.*, 1997, 2315.
- 4 C. Danjoy, J. Zhao, B. Donnadiou, J. P. Legros, L. Valade, R. Choukroun, A. Zwick and P. Cassoux, *Chem. Eur. J.*, 1998, **4**, 1100.
- 5 R. Choukroun and P. Cassoux, *Acc. Chem. Res.*, 1999, **32**, 494.
- 6 W. Ahlers, B. Temme, G. Erker, R. Fröhlich and T. Fox, *J. Organomet. Chem.*, 1997, **527**, 191.
- 7 (a) G. Erker, W. Frömberg, R. Mynott, B. Gabor and C. Krüger, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 463; (b) G. Erker, W. Frömberg, R. Benn, R. Mynott, K. Angermund and C. Krüger, *Organometallics*, 1989, **8**, 911; (c) G. L. Wood, C. B. Knobler and M. F. Hawthorne, *Inorg. Chem.*, 1989, **28**, 382; (d) U. Rosenthal and H. Görls, *J. Organomet. Chem.*, 1992, **439**, 36.
- 8 (a) G. Erker, M. Albrecht, C. Krüger, M. Nolte and S. Werner, *Organometallics*, 1993, **12**, 4979; (b) A. D. Jenkins, M. F. Lappert and R. C. Scriverstava, *J. Organomet. Chem.*, 1970, **23**, 165.
- 9 K. Takagi, C. J. Rousset and E. Negishi, *J. Am. Chem. Soc.*, 1991, **113**, 1440.
- 10 E. S. Johnson, G. J. Balaich, P. E. Fanwick and I. P. Rothwell, *J. Am. Chem. Soc.*, 1997, **119**, 11 086.
- 11 (a) P. M. Pellny, F. G. Kirchbauer, V. V. Burlakov, W. Baumann, A. Spannenberg and U. Rosenthal, *J. Am. Chem. Soc.*, 1999, **121**, 8313; (b) P. M. Pellny, F. G. Kirchbauer, V. V. Burlakov, A. Spannenberg, K. Mach and U. Rosenthal, *Chem. Commun.*, 1999, 2505 and references therein.
- 12 R. Choukroun, B. Donnadiou, J. Zho, P. Cassoux, C. Lepetit and B. Silvi, *Organometallics*, 2000, **10**, 1901.
- 13 (a) V. V. Burlakov, N. Peulecke, W. Baumann, A. Spannenberg, R. Kempe and U. Rosenthal, *J. Organomet. Chem.*, 1997, **536**, 293; (b) S. Pulst, P. Arndt, B. Heller, W. Baumann, R. Kempe and U. Rosenthal, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1112.
- 14 U. Rosenthal, A. Ohff, W. Baumann, R. Kempe, A. Tillack and V. V. Burlakov, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1605.
- 15 B. Temme, G. Erker, R. Fröhlich and M. Grehl, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1480.
- 16 A stable tweezer bimetallic alkynylcopper(i) titanium acetylide has been characterized. M. D. Janssen, M. Herres, L. Zsolnai, D. Grove, A. L. Spek, H. Lang and G. van Koten, *Organometallics*, 1995, **14**, 1098.