

Pergamon

PII: S0040-4020(96)01076-9

Charge-transfer Complex Formation and Photo-induced Electron-transfer Reaction of Dibenzo-7-silabicyclo[2.2.1]hepta-2,5-dienes

Masahiro Kako, Masahiro Mori, Kaname Hatakenaka, Seiji Kakuma, Yasuhiro Nakadaira,* Masanori Yasui,[†] and Fujiko Iwasaki[†]

Department of Chemistry and Department of Applied Physics and Chemistry, [†] The University of Electro-Communications, Chofu, Tokyo 182, Japan

Abstract: Dibenzo-7-silabicyclo[2.2.1]hepta-2,5-dienes (1a, 1b) are excellent electron donors because of effective σ - π conjugation between the orbitals of C-C π bonds and Si-C σ bonds. Some of their donor properties are demonstrated by the reactions with some electron acceptors. When 1a and 1b are mixed with tetracyanoethylene, facile formation of charge-transfer complexes was observed. In the 2,4,6-triphenylpyrylium tetrafluoroborate-sensitized photoreaction of 1b, the corresponding difluorosilane and anthracene were obtained in good yields. The structural and electronic features of radical cation 1a^{+*} were provided by semiemperical molecular orbital calculation. In addition, the structure of 1a in crystals was determined by X-ray crystallography and compared with that obtained by the calculation. © 1997, Elsevier Science Ltd. All rights reserved.

INTRODUCTION

Electron-transfer chemistry of group 14 organometallic compounds has been producing fascinating results.¹⁻⁶⁾ β -silylated olefinic and aromatic compounds are the most commonly used electron-donors since the HOMOs of these substrates are significantly destabilized due to σ - π conjugation between the orbitals of C-C π bonds and Si-C σ bonds.⁷⁾ Recently, we indicated that dibenzo-7-silabicyclo[2.2.1]hepta-2,5-dienes (1) and 7,8-disilabicyclo[2.2.2]octa-2,5-dienes (2) act as electron donors to undergo skeletal rearrangement and C-Si bond cleavage, respectively.⁸⁾ Matsumoto and coworkers have also reported electron donating properties of some polycyclic carbosilanes by investigating charge-transfer (CT) complex formation.⁹⁾ As a part of our continuing studies of electron-transfer reaction of organosilicon compounds, we now report donor-acceptor interaction of 1 with some electron acceptors. Also described are structural and electronic properties of both the ground state neutral 1 and the corresponding radical cation 1^{+•} on the basis of semiemperical molecular orbital calculation.





1a: R = 2,6-dimethylphenyl, 1b: R = mesityl

2a: R = Me, 2b: R = ⁱPr, 2c: R = CH₂^tBu

RESULTS AND DISCUSSION

Since 1a and 1b¹⁰ have fairly low Formation of CT Complexes between 1 and Electron Acceptors. oxidation potentials (1a: $E_{ox} = +1.45$ V vs SCE, 1b: $E_{ox} = +1.40$ V vs SCE), we investigated CT complex Upon mixing 1a and tetracyanoethylene (TCNE) in CH₂Cl₂, an intense formation with electron acceptors. coloration with an absorption maximum at 467 nm developed immediately. According to Mulliken's chargetransfer theory,¹¹ the absorption bands of the CT complex shift to a longer wavelength region as the magnitude of electron donating abilities of donors increases. Thus, when 1b, which has a lower oxidation potential than 1a, was used as the electron donor, a substantial red shift was observed affording an absorption maximum Previously, we^{8a)} and Matsumoto's group^{9c)} have reported that 2a-c and TCNE form CT at 507 nm. complexes with absorption maxima at 630~650 nm, and the donor properties of 2a-c are attributable to the effective overlapping of π -orbitals of the aromatic ring with the benzylic Si-C σ bonds. Meanwhile, in the case of 1, such σ - π conjugation should operate less effectively than does in 2 as expected from that dihedral angle between the π -orbitals of the anthracene moiety and the benzylic Si-C σ bonds is 43° (calculated by the MOPAC PM3 method, see below), much larger than that of 2c (179).^{9c)} Similarly, 1b and benzoquinone derivatives, chloranil (CL), tetracyanoquinodimethane (TCNQ), and dichlorodicyanobenzoquinone (DDQ) afforded CT complexes with absorption maxima as summarized in Table 1, whereas those of 1a could not be observed even with DDQ probably due to spectral overlapping by intense absorption of the acceptors. There seems to be a linear correlation between the charge transition energies (hv_{cT}) and the reduction potentials (E_{red}) of acceptors for the 1 b-quinone system as expected from Mulliken's charge-transfer theory.^{If,g,11)}



Table 1. UV Absorption Maxima of CT Complexes of 1b with Acceptors.

Acceptor	E _{red} [V] vs SCE ¹²⁾	λ_{CT} [nm]	hv _{CT} [eV]
TCNE	+0.24	507	2.45
DDQ	+ 0.52	500	2.48
TCNQ	+ 0.18	445	2.79
CL	+ 0.04	420	2.95

Photosensitized Reaction of 1 with a Pyrylium Salt. Recently we reported that 1b underwent skeletal rearrangement under photolytic conditions using 9, 10-dicyanoanthracene (DCA) as a sensitizer (Scheme 1, eq.

1).^{8b)} It is of particular interest to examine whether such a rearrangement takes place under other electron-Since pyrylium salts have been used in various photo-induced electron-transfer transfer conditions. reactions,¹³⁾ we studied the photoreaction of 1b in the presence of 2,4,6-triphenylpyrylium tetrafluoroborate (TPP^+BF_4) . When a CH₂Cl₂ solution of 1b (2.8 x 10^{-2} M) in the presence of TPP⁺BF₄⁻ (1.5 x 10^{-2} M) was irradiated with 500 W tungsten-halogen lamps (passing through an aqueous NaNO₂ solution filter, cutoff < 400 nm), 1b was rapidly consumed and anthracene (5) and difluorodimesitylsilane (6b)¹⁴⁾ were obtained in 89 and 59 %, respectively, as shown in Table 2 (Scheme 1, eq. 2). The following observation is supportive of an (i) The fluorescence of TPP⁺BF₄ was efficiently quenched with 1b ($k_a \tau = 70$ electron-transfer mechanism. (ii) The free energy change $(\Delta G)^{15}$ of electron-transfer process between 1b and TPP⁺BF₄ is calculated M¹). to be -29.9 kcal/mol, which is indicative of exothermic electron-transfer from 1b to the excited singlet state of The reaction did not occur without the sensitizer or under a dark condition and was suppressed by TPP⁺BF₄. addition of electron donors such as 1,4-diazabicyclo[2.2.2]octane ($E_{ox} = + 0.70$ V vs SCE). On the other hand, neither exciplex emission nor charge-transfer absorption was observed between 1b and the acceptors Whereas the results presented above cannot exclude possibilities of initial formation of $(TPP^+BF_4$ and DCA). an exciplex or a charge-transfer complex, it would be reasonable to discuss these reactions as a process induced by initial electron-transfer from 1b to the acceptors.



A proposed reaction mechanism for 1b is shown in Scheme 2. Initially, the bridge Si-C bond would be oxidized and cleaved by a nucleophilic attack of BF_4^- to form an open intermediate A, which may subsequently release neutral anthracene and fluorosilyl radical B. Since it has been reported that silyl radicals have quite low ionization potentials, ¹⁶ B might undergo further one-electron oxidation to give a silyl cation C, which should react with BF_4^- as a F⁻ source to afford difluorosilane 6b. Recent reports have demonstrated that nucleophilies such as alcohols have an important role in the cleavage reactions of Si-Si and Si-C bonds of organosilicon radical cations.¹⁰ Furthermore, the electrochemical oxidation of cyclic polysilanes $(R_2Si)_n$ using n-Bu₄N⁺BF₄⁻ as supporting electrolyte has been reported to give α, ω -difluoropolysilanes F-(R₂Si)_n-F, and explained in terms of fluorinative Si-Si bonds cleavage of polysilane radical cations.¹⁷⁾ These reports would offer a basis for our mechanistic account.

In the TPP⁺BF₄⁻-sensitized reaction of 1b, neither 3b nor 4b was detected in contrast to the case of the DCA-sensitization,^{8b)} which suggests that the fluorinative cleavage reaction should occur at an earlier stage than

the rearrangement in the total reaction pathway. The Si-F bonds have so large bond formation energies¹⁸) that fluoride ions might be transferred quite rapidly from BF_4^- to radical cation $1b^{+\circ}$ and eventually furnish 6b. In fact, 6b was readily produced when DCA-sensitized photoreaction of 1b was carried out in the presence of n-Bu₄N⁺BF₄⁻ whereas production of the rearrangement products, 3b and 4b, was suppressed. Under the same conditions, 3b was confirmed to be stable and 4b was found to be photoreactive, but afforded neither 5 nor 6b at all. Since TCNE is also known as a strong acceptor to oxidize various donors¹⁹ we performed photolysis of a CH₂Cl₂ solution of 1b and TCNE in the presence of n-Bu₄N⁺BF₄⁻ with visible light. As expected, 1b was facilely consumed to give 6b along with the Diels-Alder adduct of anthracene and TCNE (7)²⁰ as shown in Table 2.

Table 2.	Photosensitized	Fluorination	

Substrate	Reaction Condition	Time/h	Coversion/%	Pro	ducts	and Y	ields/	%	
				_3b	4b	5	6b	7	
1b	hv / $TPP^+BF_4^- / CH_2Cl_2$	2	100	0	0	89	59		
1b	hv / DCA / $n-Bu_4N^+BF_4^-$ /CH ₂ Cl ₂	3	78	18	18	36	14		
1b	hv / DCA /CH ₂ Cl ₂	3	89	31	36	18	0		
1b	$hv / TCNE / n-Bu_4N^+BF_4^-/CH_2Cl_2$	2 3	85	18	18	0	14	36	



Scheme 2.





Structure Determination and Semiemperical MO Calculation. The optimized structure of 1a and the corresponding radical cation $1a^{+*}$ are calculated by the MOPAC PM3²¹⁾ method, which are shown in Figure 1 with selected geometrical parameters in Table 3. To verify the validity of the calculation, we carried out the

X-ray analysis of 1a in crystals. Two sets of four molecules of 1a (1a-A and 1a-B) are contained in the unit cell and the molecular structure of 1a-A is shown with important bond lengths and angles in Figure 2 and Table The bond lengths and angles are fairly close to with those reported for related silanorbornadiene 3. derivatives.²²⁾ The bridge Si-C bonds are slightly longer than normal Si-C bonds.²³⁾ Similar bond elongation in silanorbornadiene skeleton was reported and explained in terms of delocalization of electron density of the bridge Si-C σ bond to the π bonds of the anthracene moiety.^{22c)} Validity of the optimized structure of **1a** was thus supported by a comparison with that obtained by the X-ray analysis. Adiabatic ionization potentials (IP) of 1a and 1b were also estimated to be 7.42 and 7.36 eV, respectively, by calculating the gap of heats of formation between the neutral and the radical cation. As shown in Table 4, these values are reasonable since the calculated IP values for bis- and tetrakis-Me₃SiCH₂ benzenes as related compounds show good consistency with those measured by photoelectron spectroscopy.²⁴⁾ The calculated HOMO drawing of 1a is illustrated in Figure 3. As reported by Bock and Alt,^{7a)} the HOMO is surely delocalized on both the aromatic π systems and the benzylic C-Si bond.

We investigated the characteristics of the radical cation $1a^{+\bullet}$ to gain insight into the reaction mechanism. The most important structural features of the $1a^{+\bullet}$ are as follows, (i) the one of the bridge Si-C bond (Si(1)-C(1), 2.60Å) being much longer compared to the other (Si(1)-C(4), 1.98Å) and to those of neutral 1a (Si(1)-C(1), 2.02Å; Si(1)-C(4), 2.03Å), (ii) the marked flattening of both the Si(1) atom and the C(1) atom (sum of

	X-ray	Data	Calcula	ated Data
	1a-A	<u>1a-B</u>	<u>1a</u>	1a**
0 '(1) ((1)	1044(5)	1.020(5)	2.024	2 500
Si(1)-C(1)	1.944(5)	1.939(3)	2.024	2.599
$S_1(1) - C(4)$	1.930(5)	1.930(3)	2.020	1.983
Si(1)-C(7)	1.905(5)	1.904(5)	1.875	1.821
Si(1)-C(8)	1.886(5)	1.914(5)	1.8/8	1.835
C(1)-C(6)	1.509(7)	1.533(7)	1.491	1.434
C(1)-C(2)	1.527(7)	1.504(7)	1.491	1.433
C(4)-C(5)	1.527(7)	1.506(7)	1.488	1.480
C(4)-C(3)	1.554(8)	1.508(7)	1.493	1.479
C(6)-C(5)	1.413(7)	1.397(7)	1.417	1.426
C(2)-C(3)	1.375(8)	1.417(7)	1.417	1.427
C(1)-Si(1)-C(4)	80.4(2)	79.5(2)	78.379	70.752
C(1)-Si(1)-C(7)	112.3(2)	112.3(2)	111.543	104.552
C(1)-Si(1)-C(8)	117.9(2)	118.9(2)	119.713	114.344
C(4)-Si(1)-C(7)	119.2(2)	119.4(2)	117.881	123.273
C(4)-Si(1)-C(8)	110.5(2)	110.3(2)	114.132	114.917
C(7)-Si(1)-C(8)	113.0(2)	112.9(2)	111.715	117.580
Si(1)-C(1)-C(6)	97.8(3)	97.6(3)	96.418	82.076
Si(1)-C(1)-C(2)	97.0(3)	98.1(3)	95.637	83.154
C(6) - C(1) - C(2)	107.4(4)	107.1(4)	107.362	117.662
Si(1)-C(4)-C(5)	96.4(3)	96.2(3)	96.485	97.047
Si(1)-C(4)-C(3)	97.1(3)	98.8(3)	96.233	99.124
C(5) - C(4) - C(3)	107.6(4)	107.7(4)	107.663	111.146
C(1)-C(6)-C(5)	111.6([°] 4)	109.8(4)	112.721	117.444
C(4)-C(5)-C(6)	110.3(4)	112.3(4)	112.208	114.007
C(1)-C(2)-C(3)	111.2(5)	111.5(4)	112.547	117.383
C(4)-C(3)-C(2)	111.7(5)	110.2(4)	112.363	113.932

Table 3. Selected Bond Distances (Å) and Angles (°)

angles, C(7)-Si(1)-C(8), C(7)-Si(1)-C(4), and C(4)-Si(1)-C(8) being 355.8°, sum of angles, C(2)-C(1)-C(6), C(2)-C(1)-H(1), and C(6)-C(1)-H(1) being 356.4°), (iii) the slight but appreciable shortening of Si(1)-C(4), Si(1)-C(7), Si(1)-C(8), C(1)-C(2), and C(1)-C(6) bonds on the transition from 1 a to $1a^{+\circ}$. Some electronic properties of the optimized radical cation are summarized in Table 5. A significant population of the positive charge is observed at the Si(1) atom whereas the spin population is more predominant at the anthracene unit than at the silicon atom and the 2,6-dimethylphenyl group. Therefore, it is suggested that Si(1)-C(1) bond of $1a^{+\circ}$ has a tendency to dissociate to a cationic silyl site and a diarylmethyl radical site. Similar structural and electronic aspects were reported for radical cation [t-Bu₃SiH]^{+•} calculated by the MOPAC AM1 method.²⁵⁾

In addition to the HOMO of 1a, these findings described above indicate that one electron is removed from one of the bridge Si-C bonds, which would be weakened subsequently to afford the relaxed form of $1a^{+\bullet}$ as depicted in Figure 1. The presence of nucleophiles such as alcohol and fluoride ion should accelerate the cleavage of the Si-C bond.

Figure 1. Optimized Structures of the Neutral 1a (left) and the Radical Cation of 1a (right).



Figure 2. ORTEP Drawing of 1a.

Figure 3. Calculated HOMO Orbital for 1a.



*Experimental values are in parentheses.(see text)

	opin Demoir of Audre	
Atom	Charge (∆Charge*)	Spin Density
Si(1)	±0.905 (±0.443)	+0.200
$\mathbf{C}(1)$	-0.072 (-0.006)	-0.154
$\tilde{C}(2)$	-0.061 (-0.001)	+0.467
C(3)	-0.073 (-0.018)	-0.417
C(4)	-0.058 (+0.029)	+0.633
C(5)	-0.069 (-0.018)	-0.425
C(6)	-0.061 (+0.001)	+0.499
C(7)	-0.257 (-0.063)	-0.155
C(8)	-0.264 (-0.067)	-0.111
Fragment	Charge(∆Charge*)	Spin Density
Si atom	+0.904(+0.443)	+0.200
2.6-dimethylphenyl	-0.134 (+0.178)	+0.022
anthracene	+0.229 (+0.379)	+0.778

Table 5.	Charge and	l Snin	Density	of	Radical	Cation	19#
•		-		`	•		

	ŧŌ	hange of	the cl	narge from	n the ne	utral to	the cation	radica
--	----	----------	--------	------------	----------	----------	------------	--------

EXPERIMENTAL

NMR spectra were recorded with a Varian Unity-plus 500 spectrometer. Deuteriated chloroform and benzene were used as the solvent. Mass spectral data were obtained on a Shimadzu QP-1000 mass spectrometer. UV-visible spectra were obtained with a HITACHI U-3300 spectrometer. GLC-analyses were carried out on a Shimadzu GC-14A equipped with a 0.25 mm x 25 m CBP1 capillary column. Cyclic voltammograms of 1 were obtained on 0.1 M n-Bu₄NClO₄/CH₂Cl₂ solution (vs SCE; scan rate, 200 mV/s; Hokuto Denko Ltd., a potentiostat/galvanostat HA-501 and a function generator HB-104). Oxidation potentials (E_{ox}) of 1a and 1b are + 1.40 and +1.45 eV vs SCE (irreversible), respectively. The ΔG values were calculated according to the Rehm-Weller equation $(\Delta G(\text{kcal/mol})=23.06[E(D/D^+)-E(A/A)-e_o^2/\epsilon_a-\Delta E_{0,0}]^{9)}$ by using the excited singlet energies of $TPP^+BF_4^-$ (2.83 V), the reduction potentials of $TPP^+BF_4^-$ (-0.29 V vs SCE),¹³⁾ and -0.12 eV for e₀²/εa in CH₂Cl₂.²⁶⁾ Quenching of TPP⁺BF₄ fluorescence was measured by using a HITACHI F-4500 fluorescence spectrometer. Irradiation was carried out by two 500-W tungsten-halogen lamps using a Pyrex filter in a water bath while N₂ passed through the photolysate. DCA, CL, DDQ, TCNQ, n-Bu₄N⁺BF₄ (Tokyo Kasei) and TPP⁺BF₄ (Aldrich) were used as received. TCNE (Tokyo Kasei) was used after sublimation in vacuo. Compounds $1a^{10}$, $1b^{10}$, $6b^{14}$, and 7^{20} were prepared according to the literature.

 $TPP^*BF_4^-$ -sensitized Photoreactions of 1b. In a typical experiment, 1b (2.8 x 10⁻² M) was dissolved in a CH₂Cl₂ solution (4 ml) with TPP⁺BF₄⁻ (1.5 x 10⁻² M) as a sensitizer, and this solution was photolyzed. The reaction was monitored by GLC. Formation of 5 and 6b was confirmed by comparison of their retention time on GLC with those of the authentic sample and GC-MS analysis. The yields of 5 and 6b were determined by ¹H-NMR analysis.

DCA-sensitized Photoreactions of 1b. In a typical experiment, a CH₂Cl₂ solution (4 ml) of 1b (1.7 x 10^{-2} M) and DCA (1.3 x 10^{-3} M) in the presence of n-Bu₄N⁺BF₄ (1.7 x 10^{-1} M) was photolyzed and the reaction was monitored by GLC. Formation of 5 and 6b was confirmed by comparison of its retention time The yields of **3b**, **4b**, **5**, and **6b** were determined by ¹H-NMR analysis. on GLC. 2: colorless crystals; mp 177~179°C; ¹H-NMR(CDCl₁) & 7.63(d, 1H, J=7.3Hz), 7.48(d, J=7.3Hz, 1H), 7.32(t, J=7.3Hz, 1H), 7.13(t, 1H, J=7.3Hz), 6.93~6.90(m, 2H), 6.745(1H, t, J=7.3Hz), 6.744(s, 2H), 6.67(s, 2H), 5.89(d, 1H, J=7.3Hz), 5.21(d, 1H, J=4.6Hz), 4.32(d, 1H, J=4.6Hz), 2.25(s, 9H), 2.21(s, 3H), 1.94(brs, 6H); ¹³C- $NMR(CDCl_3) \delta 150.90(s), 149.97(s), 145.62(s), 143.07(s), 142.92(s), 139.84(s), 138.55(s), 138.33(s), 138.33$ 134.97(d), 133.62(s), 133.26(s), 129.67(d), 129.12(d), 128.32(d), 126.86(d), 126.69(d), 125.47(d), 125.39(d), 123.34(d), 120.37(d), 55.10(d), 39.30(d), 24.86(q), 24.49(q), 21.11(q), 20.96(q); ²⁹Si-NMR(CDCl₃) δ -7.32; MS m/z (%) 444(M, 19), 429(100), 368 (23); Anal. Calcd for C₃₂H₃₂Si: C, 86.43; H, 3: colorless crystals; mp 226~227°C; ¹H-NMR(CDCl₃) δ 7.70(d, 1H, 7.25. Found: C, 86.54; H, 7.27. J=7.3Hz), 7.52(d, 1H, J=7.3Hz), 7.36(d, 1H, J=7.3Hz), 7.13~7.05(m, 2H), 7.02~6.93(m, 5H), 6.70(s, 2H), 4.47 (d, 1H, J=1.4Hz), 3.64(d, 1H, J=1.4Hz), 2.51(s, 6H), 2.31(s, 3H), 2.13(s, 3H), 2.07(brs, 6H); 13 C-NMR(CDCl₂) δ 156.37(s), 153.23(s), 145.52(s), 143.03(s), 141.97(s), 140.77(s), 137.70(s), 136.07(s), 136.07 134.72(s), 132.14(d), 132.11(d), 129.69(d), 128.67(d), 128.04(d), 127.69(d), 125.71(d), 125.34(s), 125.03(d), 124.81(d), 122.34(d), 67.18(s), 59.05(s), 23.27(q), 22.77(q), 21.23(q), 20.57(q); ²⁹Si-NMR(CDCl₃) δ -9.70; MS m/z (%) 444(M⁺, 15), 429(100); Anal. Calcd for C₃₂H₃₂Si: C, 86.43; H, 7.25. Found: C, 86.22; H, 7.40.

Photolysis of the 1b-TCNE CT complex. A CH_2Cl_2 solution (4 ml) containing 1b (4.5 x 10^{-2} M), TCNE (1.7 x 10^{-1} M), and n-Bu₄N⁺BF₄⁻ (9.0 x 10^{-2} M) was photolyzed and the reaction was monitored by TLC. Formation of 6b and 7 was confirmed by comparison of ¹H-NMR spectra with those of the authentic samples. The yields of 6b and 7 were also determined by ¹H-NMR analysis.

X-ray Crystal Analysis. Intensity data were collected with Mo K α radiation (graphite monochrometer λ =0.71073Å) on a Rigaku AFC-4 diffractometer. Crystal data for 1a,: C₃₀H₂₈Si (fw 416.64), orthorhombic Pna2₁; a= 16.681(2) Å; b = 8.701(2) Å; c= 31.817(7) Å; V=4618(2) Å³; Z=8. A total of 5725 reflections within 20 = 55° were measured by the ω scan method with a scan rate of 49min. The final R factor was 0.056 (Rw= 0.072) for 3943 reflections of Fo> 3 σ (Fo).

Semiemperical Calculation. The calculations were carried out by the CAChe MOPAC program (Release 3.8, available from SONY Tektronix Co. Ltd.) on an Apple Macintosh computer. Optimized structures of the neutrals and the radical cations were obtained by using restricted and unrestricted Hartree-Fock wave function, respectively.

Acknowledgment: This work was supported in part by Grants-in-Aid for Scientific Research on Priority Areas No. 0623922 and No. 06227226 and for Encouragement of Young Scientists No. 06854031 from the Ministry of Education, Science, and Culture in Japan. Y. N. acknowledges SUNBOR Grant from SUNTORY Institute for Bioorganic Research.

REFERENCES

- (a) Sakurai, H.; Sakamoto, K.; Kira, M. Chem. Lett. 1984, 1213. (b) Nakadaira, Y.; Komatsu, N.; Sakurai, H. ibid. 1985, 1781. (c) Kyushin, S.; Ehara, Y.; Nakadaira, Y.; Ohashi, M. J. Chem. Soc., Chem. Commun. 1989, 279. (d) Kyushin, S.; Masuda, Y.; Matsushita, K.; Nakadaira, Y.; Ohashi, M. Tetrahedron Lett. 1990, 31, 6395. (e) Nakadaira, Y.; Ohkura, Y.; Kyushin, S.; Ohashi, M.; Ueno, K.; Kanouchi, S.; Sakurai, H. ibid., 1992, 33, 4013. (f) Kyushin, S.; Ohkura, Y.; Nakadaira, Y.; Ohashi, M.; Yasui, M.; Iwasaki, F. Chem. Lett. 1991, 883. (g) Kyushin, S.; Baba, Y.; Nakadaira, Y.; Ohashi, M. Main Group Metal Chemistry 1995, 299, and references cited therein.
- (a) Nakanishi, K.; Mizuno, K.; Otsuji, Y. Bull. Chem. Soc. Jpn. 1993, 66, 2371. (b) Tamai, T.; Mizuno, K.; Hashida, I.; Otsuji, Y. *ibid.* 1993, 66, 3747. (c) Mizuno, K.; Nakanishi, K.; Chosa, J.; Otsuji, Y. J. Organomet. Chem. 1994, 473, 35, and references cited therein.
- (a) Yoon, U. C.; Mariano, P. S. Acc. Chem. Res. 1992, 25, 233. (b) Fagnoni, M.; Mella, M.; Albini, A. Tetrahedron 1994, 21, 6401. (c) Mella, M.; Fusan, E.; Albini, A. J. Org. Chem. 1992, 57, 6210. (d) d'Alessandro, N.; Albini, A.; Falvey, D. E.; Mariano, P. S. J. Am. Chem. Soc. 1994, 116, 4211, and references cited therein.
- (a) Fukuzumi, S.; Kitano, T.; Mochida, K. J. Am. Chem. Soc. 1990, 112, 3246. (b) idem. J. Chem. Soc., Chem. Commun. 1989, 279. (c) idem. Chem. Lett. 1984, 1213, and references cited therein.
- (a) Akasaka, T.; Sato, K.; Kako, M.; Ando, W. Tetrahedron Lett. 1991, 32, 6605. (b) idem, Tetrahedron, 1992, 48, 3283; (c) Ando, W.; Kako, M.; Akasaka, T. Chem. Lett. 1993, 1679, and references cited therein.
- (a) Nakadaira, Y.; Sekiguchi, A.; Funada, Y.; Sakurai, H. Chem. Lett. 1991, 327. (b) Nakadaira, Y.; Otani, S.; Kyushin, S.; Ohashi, M.; Sakurai, H.; Funada, Y.; Sakamoto, K.; Sekiguchi, A. *ibid.* 1991, 601. (c) Dinnocenzo, J. P.; Farid, S.; Goodman, J. L.; Gould, I. R.; Mattes, S. L.; Todd, W. P. J. Am. Chem. Soc. 1989, 111, 8973. (d) Todd, W. P.; Dinnocenzo, J. P.; Farid, S.; Goodman, J. L.; Gould, I. R. *Tetrahedron Lett.* 1993, 34, 2863. (e) Dinnocenzo, J. P.; Farid, S.; Goodman, J. L.; Gould, I. R.; Todd, W. P. Mol. Cryst. Liq. Cryst. 1991, 194, 151. (f) Watanabe, H.; Kato, M.; Tabei, E.; Kuwabara, H.; Hirai, N.; T. Sato, T.; Nagai, Y. J. Chem. Soc., Chem. Commun. 1986, 1662, and references cited therein.
- (a) Bock, H.; Alt, H. J. Am. Chem. Soc. 1970, 92, 1569. (b) Bock, H. Angew. Chem. Int. Ed. Engl. 1989, 28, 1627.
- (a) Nakadaira, Y.; Gomi, Y.; Hosoe, H.; Kyushin, S; Kako, M.; Hatakenaka, K.; Ohashi, M. Bull. Chem. Soc. Jpn. 1993, 66, 344. (b) Kako, M.; Kakuma, S.; Hatakenaka, K.; Nakadaira, Y.; Yasui, M.; Iwasaki, F. Tetrahedron Lett. 1995, 35, 6293.
- (a) Kyushin, S.; Izumi, Y.; Tsunakawa, S.; Matsumoto, H. Chem. Lett. 1992, 1393. (b) Kyushin, S.;
 Ikarugi, M.; Tsunakawa, S.; Izumi, Y.; Miyake, M.; Matsumoto, H.; Goto, M. J. Organomet. Chem.
 1994, 473, 19. (c) Kyushin, S.; Ikarugi, M.; Takatsuna, K.; Goto, M.; Matsumoto, H. *ibid.* 1996, 510, 121.
- 10) Sakurai, H.; Oharu, K.; Nakadaira, Y. Chem. Lett. 1986, 1797.
- 11) Mulliken, R. S. J. Am. Chem. Soc. 1952, 74, 811.

- (a) Chanon, M.; Eberson, L. in "Photoinduced Electron Transfer" Fox, M. A.; Chanon, M., Eds.;
 Elsevier: Amsterdam, 1988, Part A. Chapter 1.10, p. 409. (b) Foster, R.; Foreman, M. I. in "The Chemistry of the Quinonoid Compounds" Patai, S. Ed.; Wiley, New York 1974, Chapter 6, p. 257.
- 13) For a review, see; Miranda, M. A.; García, H. Chem. Rev. 1994, 94, 1063.
- 14) Damrauer, R.; Simon, R. A.; Kanner, B. Organometallics 1988, 7, 1161.
- 15) Rehm, D.; Weller, A. Isr. J. Chem. 1970, 8, 259.
- 16) Chatgilialoglu, C. Chem. Rev. 1995, 95, 1229.
- 17) Becker, J. Y.; Shakkour, E.; West, R. Tetrahedron Lett. 1992, 33, 5633.
- Corey, J. Y. in "The Chemistry of Organosilicon Compounds" Patai, S.; Rappoport, Z., Eds.; Wiley, New York 1989, Chapter 1, p. 1.
- (a) Hilinski, E. F.; Masnovi, J. M.; Kochi, J. K. Rentzepis, P. M. J. Am. Chem. Soc. 1984, 106, 8071. (b) Miyashi, T.; Kamata, M.; Mukai, T. *ibid.* 1987, 109, 2780, and references cited therein.
- (a) Brown, P.; Cookson, R. C. Tetrahedron 1965, 21, 1977. (b) Thompson, C. C.; Holder, D. D. J. Chem. Soc., Perkin Tran. 2 1972, 257.
- 21) (a) Stewart, J. J. P. J. Comput. Chem. 1989, 10, 209. (b) Idem, QCPE Bull. 1989, 9, 10.
- (a) Preut, H.; Mayer, B.; Neumann, W. P. Acta Crystallogr. 1983, 39C, 1118. (b) Appler, H.; Gross, L. W.; Mayer, B.; Neumann, W. P. J. Organomet. Chem. 1985, 291, 9. (c) Sekiguchi, A.; Ziegler, S. S. Haller, K. J. West, R. Recl. Trav. Chim. Pays-Bas 1988, 107, 197.
- 23) Sheldrick, W. S. in ref. 17, Chapter 3, p. 227.
- 24) Bock, H.; Solouki, B., in ref. 17, Chapter 9, p. 555.
- 25) Glidewell, C.; Rhodes, C. J. J. Organomet. Chem. 1994, 471, 43.
- 26) Mattay, J.; Runsink, J.; Gersdorf, J.; Rumbach, T. Helv. Chim. Acta 1986, 69, 442.

(Received in Japan 24 September 1996; accepted 18 November 1996)