## An SO Elimination leading to Octafluorodibenzofuran

by R. D. CHAMBERS and J. A. CUNNINGHAM\* (University Science Laboratories, South Road, Durham)

We recently reported<sup>1</sup> the first synthesis of a perfluorinated compound in the dibenzo-series, octafluorodibenzothiophen (I), and we have



subsequently found that a dioxide (II) can be formed by the reaction of (I) with 85% hydrogen peroxide in trifluoroacetic anhydride, while other methods gave little or incomplete reaction. Current interest in the elimination of sulphur monoxide from sulphones<sup>2</sup> led us to examine the mass spectrum of the dioxide (II) and a large peak at 312 mass units, corresponding to P - SO was observed. Subsequently, we pyrolysed the dioxide (II) and found that the new heterocycle, octafluorodibenzofuran, m.p. 100°, could be isolated in 72% yield. The pyrolysis temperature and conditions were critical; general decomposition occurred on pyrolysis in a sealed tube but, using a flow system, products were obtained which became more simple as the temperature was raised. At 810° octafluorodibenzofuran was obtained in high

yield together with small amounts of octafluorodibenzothiophen and a minor unknown impurity. In contrast to these results, pyrolysis of decafluorodiphenyl sulphone (IV), under various conditions, gave only sulphur dioxide elimination, leading to decafluorobiphenyl.

$$(C_6F_5)_2SO_2 \rightarrow C_6F_5 \cdot C_6F_5$$

The mechanism of the SO elimination possibly involves an intramolecular rearrangement (internal nucleophilic displacement) to an unstable sulphinic ester, followed by SO loss.



Elimination of SO or  $SO_2$  from a compound would then depend on the relative rates of rearrangement (i) to C-S bond cleavage (ii). These possibilities are being investigated.

Octafluorodibenzofuran undergoes nucleophilic substitution by a variety of nucleophiles but the orientation, while of considerable interest, cannot yet be unambiguously assigned.

(Received, April 28th, 1967; Com. 405.)

<sup>1</sup> R. D. Chambers and J. Cunningham, Chem. Comm., 1966, 469.

<sup>2</sup> Results communicated by J. F. W. McOmie and by M. P. Cava at the Symposium on "Aromaticity", Sheffield, 1966.