Synthesis, Structure and Complexation Ability of Novel Metalloreceptors Containing Two Pincer Complexes of Palladium(II)

Tatsuya Nabeshima,*^[a] Daisuke Nishida,^[a] Shigehisa Akine,^[a] and Toshiyuki Saiki^[a]

Keywords: Host-guest systems / Pincer ligands / Palladium / Metalloreceptors / Crown ethers

Metalloreceptors **1a** and **1b** were synthesized by palladation of 42- and 54-membered crown ethers that have two SCSpincer ligand moieties. The X-ray crystallographic analysis reveals that the metalloreceptor **1b** has a folded structure in which the two pincer-Pd moieties are stacked in a parallel manner. The metalloreceptor **1b** recognizes the aromatic bi-

Metalloreceptors have recently attracted much attention in supramolecular chemistry because the physical and chemical properties of the metal center can be applied in guest binding, sensing, and regulation of molecular functions.^[1] In particular, the coordination of various ligands to metals is very useful for the formation of a stable supramolecular system.^[2–6] Combination of the coordinate bonding and other intermolecular interactions provides a strong and selective recognition.

Complexes between Pd and pincer ligands are useful as they provide a nice coordination site for host-guest interaction.^[7] For example, crown ethers containing a Pd center inside the ring and molecular recognition have been reported.^[8–10] Cooperative binding of the Pd unit and the crown ether moiety to nucleobases such as cytosine and adenine was observed in this system.^[11,12] However, a host molecule containing several pincer complexes as a binding site has not been synthesized although this type of host is very interesting as a multi-point receptor.

Here we report the synthesis, structure and molecular recognition ability of large macrocyclic hosts 1, which contain two pincer complexes of Pd^{II} . The metallohosts are expected to bind bidentate guests of different sizes in the cavity because the framework of these hosts is flexible so as to cause induced-fit recognition (Scheme 1).

Metallohosts 1 containing the two Pd centers were prepared according to Scheme 2. Diols $2a^{[13]}$ and $2b^{[14]}$ were treated with *p*-toluenesulfonyl chloride to afford 3a and 3b in 80% and 63% yield, respectively. Reaction of dithiol 4^[15] with 3a, 3b in the presence of K₂CO₃ afforded diols 5a, 5b,

E-mail: nabesima@chem.tsukuba.ac.jp

dentate guests 4,4'-bipyridine, pyrazine, and pyrimidine and binds to them in a bridging fashion to form the corresponding bimacrocyclic 1:1 complex.

(© Wiley-VCH Verlag GmbH & Co. KGaA, 69451 Weinheim, Germany, 2004)



Scheme 1. Recognition of bidentate guests in a bridging fashion

and the following tosylation gave **6a**, **6b**. The ditosylates **6a** and **6b** were then treated with **4** under high dilution conditions to give crown ethers **7a** and **7b** in 42% and 45% yield, respectively. Metallation of **7** with $[Pd(CH_3CN)_4](BF_4)_2^{[16]}$ yielded metallohosts **1a**, **1b**.

All spectroscopic data are consistent with the formula of 1 (see Exp. Sect.). Palladation of 7 caused broadening of the ¹H NMR signals, probably due to the enhanced rigidity of the structure, which restricts the free rotation of the methylene group. On metallation, downfield shifts of the signals for the SCH₂CH₂ protons (e.g. $\delta = 2.60$ for **7b**, $\delta =$ 3.32 for 1b, Figure 1) and the benzyl protons is observed. This is elucidated by the decreased electron-density of these protons. ESI-MS for 1b gave signals at m/z = 614.1 $(C_{52}H_{70}O_{12}S_4Pd_2, [1b-2BF_4-2CH_3CN]^{2+})$ and m/z =1247.2 ($C_{52}H_{70}FO_{12}S_4Pd_2$, [1b-2BF₄-2CH₃CN+F]⁺), (the F⁻ anion was obtained from decomposition of the BF₄⁻ anion during the ESI-MS measurements). In addition, the X-ray crystallographic analysis clarified the structure of 1b (Figure 2). The two complex moieties exhibit square-planar geometry at Pd, with three donor atoms provided by the rigid SCS chelate. The Pd-S distances are in

3779

 [[]a] Department of Chemistry, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan Fax: (internat.) +81-29-853-6503

Supporting information for this article is available on the WWW under http://www.eurjic.org or from the author.

SHORT COMMUNICATION



Scheme 2. Synthesis of metallohosts 1

the range between 2.334 Å and 2.286 Å. The two aromatic rings, which bind the palladium atoms, are stacked. The distance between the mean planes of the benzene rings (3.4 Å) suggests the existence of π - π interactions. Furthermore, the palladium atoms are located in closely proximity of the sulfur atoms. The distance between Pd2–S2 (3.371 Å) is shorter than the sum of van der Waals radii for sulfur and palladium (3.43 Å). Two CH₃CN molecules coordinate to Pd, and the Pd1–N1 and Pd2–N2 distances are 2.105 Å and 2.092 Å, respectively.



Figure 1. ¹H NMR spectral changes of **7b** upon palladation (400 MHz, $CDCl_3$); SCH_2CH_2O and benzyl protons are indicated with open and filled circles, respectively



Figure 2. Crystal structure of **1b**; thermal ellipsoids are drawn at 30% probability level; the minor component of the disordered atoms, counter anions, and hydrogen atoms are omitted for clarity

The interaction between **1b** and 4,4'-bipyridine was evaluated by ¹H NMR spectroscopy (Figure 3). Signals of free **1b** disappear when 1.0 equiv. of 4,4'-bipyridine is added, and no change is observed in the presence of excess ligand. This result supports the 1:1 complexation between **1b** and 4,4'-bipyridine in a bridging fashion as illustrated in Scheme 1. ESI-MS also supports the 1:1 complexation. A mixture of **1b** and 4,4'-bipyridine gives a signal at m/z = 692.1 (calcd. for $C_{62}H_{78}N_2O_{12}S_4Pd_2$, $[1b-2BF_4-2CH_3CN+4,4'-bipyridine]^{2+}$). In addition, the absorbance of **1b** at 304 nm increases until 1.0 equivalent of 4,4'-bipyridine is added (Figure 4). All the experiments described above indicate the 1:1 complexation between **1b** and 4,4'-

SHORT COMMUNICATION

bipyridine. Pyrazine and pyrimidine as bidentate ligands also give rise to 1:1 complexation, as seen by ¹H NMR spectroscopic experiments, although the distances involving the ligating nitrogen atoms of pyrazine and pyrimidine are much shorter than that of 4,4'-bipyridine.



Figure 3. ¹H NMR spectral changes of **1b** upon addition of 4,4'bipyridine (400 MHz, CDCl₃, [**1b**] = 2.0 mM); signals for free and complexed **1b** are indicated with open and filled circles, respectively



Figure 4. UV/Vis spectral changes of 1b upon addition of 4,4'-bipyridine (CHCl₃, [1b] = 1.0×10^{-5} M)

Complexation of **1b** with these bidentate ligands should give two small cavities which are almost half as large as that of **1b** (Scheme 1). The change in cavity size is very interesting, because this strategy may be most useful to regulate recognition ability of artificial receptors.^[17–19]

In conclusion, we have prepared metallohosts **1** bearing two Pd centers inside the cavity, and studied the structure and guest-binding ability of **1b** in detail by ¹H, ¹³C NMR spectroscopy or X-ray crystallographic analysis. We are currently investigating this bridging strategy to control molecular recognition.

Experimental Section

Synthesis of 1a: A solution of $[Pd(CH_3CN)_4](BF_4)_2$ (302 mg, 0.680 mmol) in acetonitrile (10 mL) was added to a solution of

crown ether **7a** (271 mg, 0.322 mmol) in acetonitrile (20 mL), and the mixture was stirred for 2 h at room temperature. The mixture was refluxed for 13 h, and then the solvent was evaporated. The residue was dissolved in dichloromethane (20 mL) and filtered. The filtrate was concentrated and dried in vacuo to afford **1a** (215 mg, 51%) as a yellow solid, which was further purified by recrystallization from acetonitrile to give pale yellow crystals. ¹H NMR (400 MHz, CD₃CN): δ = 3.30 (br. s, 8 H), 3.80 (t, *J* = 4.2 Hz, 8 H), 3.90 (br. t, 8 H), 4.12 (t, *J* = 4.2 Hz, 8 H), 4.29 (br. t, 8 H), 6.77 (d, *J* = 7.6 Hz, 4 H), 6.90–6.99 (m, 10 H) ppm. ¹³C NMR (100 MHz, CD₃CN): δ = 40.00, 46.78 (br.), 69.20, 70.54 (×2, CH₂), 114.78, 122.39, 123.96, 126.65 (CH), 149.49, 151.65 (br.), 156 (br., C) ppm. ESI-MS: *m*/*z* found 1071.1 [**1a**-2BF₄– 2CH₃CN+F]⁺. C₄₈H₆₀B₂F₈N₂O₈Pd₂S₄: calcd. C 44.09, H 4.62, N 2.14; found C 43.78, H 4.57, N 2.28.

Synthesis of 1b: 1b was prepared from **7b** (319 mg, 0.315 mmol) and [Pd(CH₃CN)₄](BF₄)₂ (302 mg, 0.680 mmol) in 63% yield by a procedure similar to that for **1a. 1b**: yellow oil. ¹H NMR (400 MHz, CDCl₃): $\delta = 3.32$ (br. s, 8 H), 3.67–3.69 (m, 8 H), 3.73–3.75 (m, 8 H), 3.86–3.89 (m, 8 H), 3.90–3.92 (m, 8 H), 4.16 (t, *J* = 4.0 Hz, 8 H), 4.18 (br., 8 H), 6.79 (d, *J* = 7.0 Hz, 4 H), 6.85 (t, *J* = 7.0 Hz, 2 H), 6.92 (s, 8 H) ppm. ¹³C NMR (100 MHz, CD₃CN): $\delta = 39.68$, 46.35, 69.40, 70.50 (× 2), 71.29, 71.38 (CH₂), 115.09, 122.40, 124.01, 126.65 (CH), 149.62, 151.72, 156.59 (C) ppm. ESI-MS: *m/z* found 1247.2 [**1b**–2BF₄–2CH₃CN+F]⁺. C₅₆H₇₆B₂F₈N₂O₁₂Pd₂S₄·CHCl₃: calcd. C 42.70, H 4.84, N 1.75; found C 42.30, H 5.06, N 2.08.

X-ray Crystallographic Analysis of 1b:^[20] Pale yellow crystals (0.3 \times 0.15 \times 0.1 mm) were obtained after oily **1b** was allowed to stand at room temperature. $C_{56}H_{76}B_2F_8N_2O_{12}Pd_2S_4$, M = 1483.85. Intensity data were collected at 130 K on a Rigaku R-AXIS Rapid diffractometer with Mo- K_{α} radiation ($\lambda = 0.71069$ Å). Reflection data were corrected for Lorentz and polarization factors, and for absorption using the multi-scan method. Crystal system, triclinic, a = 13.2731(12) Å, b = 14.4368(16) Å, c = 18.1120(19) Å, $\alpha =$ 71.267(4)°, $\beta = 71.424(3)°$, $\gamma = 86.580(3)°$, $V = 3112.0(6) \text{ Å}^3$, space group $P\bar{1}$, Z = 2, μ (Mo- K_{α}) = 0.797 mm⁻¹, $D_{\text{calcd.}} = 1.584$ g/cm³, $2\theta_{\text{max}} = 52^{\circ}$, collected reflections 25108, unique reflections 12053 $(R_{\rm int} = 0.0719)$. The structure was solved by Patterson methods $(DIRDIF-99)^{[21]}$ and refined by full-matrix least-squares on F^2 using SHELXL 97.^[22] The non-hydrogen atoms were refined anisotropically except for the minor component of disordered sulfur atoms (S4 and S6). Hydrogen atoms were idealized by using the riding models. Occupancy factor involving the disordered sulfur atoms was refined to be 0.888(3) for S3 and S5, and 0.112(3) for S4 and S6. $R1 = 0.0618 [I > 2\sigma(I)], wR2 = 0.1537$ (all data), GOF $(F^2) = 1.036$ for 786 parameters.

Acknowledgments

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

- [1] J. -M. Lehn, Supramolecular Chemistry, Concepts and Perspectives, VCH: Weinheim, 1995.
- ^[2] T. Nabeshima, Coord. Chem. Rev. 1996, 148, 151-169.
- ^[3] T. Nabeshima, S. Akine, T. Saiki, *Rev. Heteroat. Chem.* 2000, 22, 219–239.
- [4] D. L. Caulder, K. N. Raymond, J. Chem. Soc., Dalton Trans. 1999, 1185-1200.

SHORT COMMUNICATION

- ^[5] M. Albrecht, Chem. Rev. 2001, 101, 3457-3497.
- [6] T. Nabeshima, Y. Yoshihira, T. Saiki, S. Akine, E. Horn, J. Am. Chem. Soc. 2003, 125, 28–29.
- [7] M. Albrecht, G. van Koten, Angew. Chem. Int. Ed. 2001, 40, 3750-3781.
- [8] J. E. Kickham, S. J. Loeb, J. Chem. Soc., Chem. Commun. 1993, 1848–1850.
- ^[9] J. E. Kickham, S. J. Loeb, Inorg. Chem. 1994, 33, 4351-4359.
- ^[10] J. E. Kickham, S. J. Loeb, *Inorg. Chem.* **1995**, *34*, 5656–5665.
- ^[11] J. E. Kickham, S. J. Loeb, S. L. Murphy, J. Am. Chem. Soc. **1993**, 115, 7031–7032.
- ^[12] J. E. Kickham, S. J. Loeb, *Organometallics* **1995**, *14*, 3584–3587.
- ^[13] S. Kopolow, T. E. Hogen Esch, J. Smid, *Macromolecules* 1973, 6, 133–142.
- ^[14] S. Shinkai, K. Inuzuka, O. Miyazaki, O. Manabe, J. Am. Chem. Soc. 1985, 107, 3950–3955.
- ^[15] T. Sato, K. Nishiyama, A. Morita, Y. Iitaka, Bull. Chem. Soc. Jpn. **1985**, 58, 2366–2369.
- ^[16] R. F. Schramm, B. B. Wayland, J. Chem. Soc., Chem. Commun. 1968, 898–899.

- ^[17] T. Nabeshima, H. Furusawa, Y. Yano, *Angew. Chem. Int. Ed. Engl.* **1994**, *33*, 1750–1751.
- ^[18] T. Nabeshima, D. Nishida, *Tetrahedron Lett.* **2002**, *43*, 5719–5722.
- ^[19] T. Nabeshima, D. Nishida, T. Saiki, *Tetrahedron* **2003**, *59*, 639–647.
- ^[20] CCDC-241251 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge at http://www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: (internat.) +44-1223-336-033; E-mail: deposit@ccdc.cam.ac.uk].
- [21] P. T. Beurskens, G. Beurskens, R. de Gelder, S. Garcia-Granda, R. O. Gould, R. Israel, J. M. M. Smits, The DIRDIF-99 program system, Crystallography Laboratory, University of Nijmegen, The Netherlands, **1999**.
- [22] G. M. Sheldrick, SHELXL 97, Program for crystal structure refinement, University of Göttingen, Germany, 1997. Received June 11, 2004