



CYANOGUANIDINE BIOISOSTERES IN POTASSIUM CHANNEL OPENERS: EVALUATION OF 3,4-DISUBSTITUTED-1,2,5-THIADIAZOLE-1-OXIDES

Heinrich J. Schostarez,^{**} Theresa J. O'Sullivan,^{*} Vincent E. Groppi,[‡] Loretta A. Cipkus-Dubray[‡]

[‡]Cell Biology and Inflammation Research, and ^{*}Medicinal Chemistry Research,

Pharmacia & Upjohn, Inc., Kalamazoo, MI 49001

Abstract. Bioisosteric substitution of the cyanoguanidine group found in pinacidil (**1**) with a 3,4-diamino-1,2,5-thiadiazole-1-oxide moiety and replacement of the 4-aminopyridine group with a 3,5-dichlorophenyl group has resulted in a new structural class of potassium channel opener (PCO). Copyright © 1996 Elsevier Science Ltd

The cyanoguanidine moiety has been a major structural constituent of several drug discovery efforts, the most notable examples being the development of the antihypertensive pinacidil (**1**) and the H₂ antagonist cimetidine (**2**).^{1,2} The former is a member of a class of drugs termed potassium channel openers (PCO's) which potentially offer additional clinical utility in areas of unmet medical need (i.e., incontinence), as well as in the treatment of asthma and hair growth related disorders such as *androgenetic alopecia*.³




Figure 1.

Recent literature reports have suggested that exploitation of differences found in subtypes of the ATP sensitive potassium channel (K_{ATP}) may lead to more selective therapeutic agents.³ Our interest in the development of more *tissue selective* potassium channel openers,⁴ specifically selective peripheral vasodilators, prompted a search for structurally novel pinacidil analogues based on the bioisosteric modification the cyanoguanidine functionality found in **1**.

It is well known in the literature that a variety of H₂ antagonists have been reported that contain structurally distinct bioisosteres of the cyanoguanidine moiety found in both **1** and **2**.¹ Utilizing these reports, the 3,4-diamino-1,2,5-thiadiazole-1-oxide functionality was selected as a bioisosteric replacement for the cyanoguanidine moiety of pinacidil based on BMY-25271 (**3**), a thiadiazole-containing analogue of ranitidine.⁵ Selection of this isostere was based on two criteria: retention of the planar relationship between the two pendant

Discussion. A series of pinacidil analogues was prepared in which the cyanoguanidine functionality was replaced with a 3,4-diamino-1,2,5-thiadiazole-1-oxide moiety. At the outset, Ar = Ph was used to provide a model system to determine the experimental conditions required for this synthetic route, subsequently 3,5-dichloroaniline was incorporated onto the thiadiazole skeleton. Three classes of side chains were added: alkylamino, cycloalkylamino and alkylphenylamino.

Table 1

Product	Ar	R ₁	Yield (%)	% Change in Fluorescence (SEM) ^a		
				10 μ M	1 μ M	0.1 μ M
8c	Ph	<i>n</i> -propyl	53	-1.05(3.66)	-4.86(4.64)	-3.75(1.79)
8h	"	<i>n</i> -amyl	62	-5.23(3.23)	-7.68(1.78)	-3.76(4.74)
9a	3,5-dichloro-	H	62	-4.58(9.36)	-4.81(2.86)	+10.19(4.54)
9b	phenyl	CH ₃	58	-33.56(2.00)	-25.12(2.18)	-16.03(0.66)
9c	"	<i>n</i> -propyl	51	-11.11(3.62)	+5.85(3.29)	+8.06(0.52)
9d	"	<i>i</i> -propyl	44	-9.36(0.92)	-5.37(0.48)	+2.12(2.85)
9e	"	<i>n</i> -butyl	92	-44.66(9.94)	-10.31(2.61)	+1.29(0.38)
9f	"	<i>i</i> -butyl	88	-10.83(6.19)	-3.59(2.37)	-1.32(1.75)
9g	"	<i>sec</i> -butyl	20	-18.97(9.32)	+1.42(4.49)	-0.43(0.82)
9h	"	<i>n</i> -amyl	34	-26.32(1.03)	-10.59(3.35)	+8.48(2.73)
9i	"	<i>neo</i> -pentyl	33	+1.84(4.83)	+5.58(3.43)	+1.77(0.58)
9j	"	<i>i</i> -amyl	55	-45.55(1.58)	-12.04(3.13)	+10.71(4.44)
9k	"	CH ₂ Ph	62	+0.84(2.87)	-0.91(4.03)	+7.83(1.33)
9l	"	(CH ₂) ₂ Ph	67	-35.18(4.92)	-1.29(1.87)	-1.01(1.39)
9m	"	(CH ₂) ₃ Ph	67	+14.09(5.31)	-19.73(1.01)	-16.02(5.95)
9n	"	(CH ₂) ₄ Ph	70	+4.02(2.68)	-6.18(1.33)	-7.20(0.64)
9o	"	cyclopropyl	49	-3.96(4.23)	-1.10(3.61)	+0.95(3.86)
9p	"	cyclobutyl	66	-21.83(3.63)	-4.23(4.05)	-5.89(4.21)
9q	"	cyclopentyl	45	-4.17(5.85)	-1.36(0.68)	-5.82(1.80)
9r	"	cyclohexyl	37	-17.74(20.3)	-2.39(2.20)	+2.05(3.27)
1	4-pyridyl	CH(CH ₃)C(CH ₃) ₃		-57.37(1.43)	-45.60(5.16)	-5.89(2.69)

^aMean of triplicate assays, assay values in arbitrary fluorescent units.

The alkylamino side chains were chosen to evaluate straight-chain and branched alkyl groups in this template. Three of these analogues (**9b**: R=methyl; **9e**: R=*n*-butyl; **9j**: R=*i*-amyl) showed modest hyperpolarization at 10 μ M. Of the cycloalkyl analogues prepared, two (**9p**: R=cyclobutyl; **9r**: R=cyclohexyl) showed low to moderate activity which was not sustained at 10 μ M. The alkylphenyl analogues examined the effects of a phenyl ring "tethered" at various lengths from the thiadiazole moiety. Activity was lost at 10 μ M when R=benzyl (**9k**) and R=phenylbutyl (**9n**). Moderate activity was seen at 10 μ M when R=phenylethyl (**9l**). A low to moderate hyperpolarization effect was seen at 1 μ M when R=phenylpropyl (**9m**). Interestingly, at 10 μ M a slight *depolarizing* effect was observed. The biochemical relevance of this observation is unclear. The most active analogues at 10 μ M from this series were **9e** (R=*n*-butyl) and **9j** (R=*i*-amyl), both with a ΔF of -45%. These analogues compared favorably in activity with pinacidil (ΔF = -60% @ 10 μ M). However, **9e** and **9j** did not retain this activity at lower doses. Pinacidil (**1**) also displayed this steep dose-response between 1 μ M and 0.1 μ M in this assay. Of the thiadiazoles evaluated, **9b** exhibited the most consistent potassium channel opening activity across the 100-fold concentration range tested. Figure 2 graphically depicts the fluorescence assay results comparing **9b** and **1**. At 0.1 μ M **9b** was actually more active than pinacidil.

It is clear that a rigorous, detailed structure-activity relationship cannot be developed from the results of this three concentration screen. The objective of this effort was the demonstration of the utility of the 3,4-diamino-1,2,5-thiadiazole-1-oxide as a bioisosteric replacement of the cyanoguanidine group. The significance of these data lie in the confirmation of this hypothesis. Additionally, the utility of the 3,5-dichlorophenyl group as a replacement of the 4-pyridyl moiety found in **1** was confirmed.

In summary, a new structural class of K_{ATP} openers has been prepared which offer insight to the requirements for binding to the ATP sensitive potassium channel in vascular smooth muscle.³ Further studies regarding the more comprehensive evaluation of these compounds in various tissues as well as the further use of the bioisosteric replacement technique will be reported in due course.




Figure 2. Comparison of 9b (upper) and 1 (lower) at various concentrations in the A-10 smooth muscle cell assay. Drug added at 7 min.¹⁰ Results are reported as a percent change in fluorescence (ΔF) where 30 mM KCl, by definition, induces a 100% change.

Acknowledgment: The authors gratefully acknowledge the assistance of the Structural, Analytical & Medicinal Chemistry Unit of Pharmacia & Upjohn in compound characterization.

References and Notes:

1. Cooper, D.G.; Young, R.C.; Durant, G.J.; Ganellin, C.R. In *Comprehensive Medicinal Chemistry*; Hansch, C., Ed.; Pergamon: Oxford, 1990; Vol. 3, pp 323-411.
2. Petersen, H.J.; Nielsen, C.K.; Arrigoni-Martelli, E. *J. Med. Chem.* **1978**, *21*, 773.
3. Longman, S.D.; Hamilton, T.C. *Medicinal Research Reviews* **1992**, *12*, 73. Empfield, J.R.; Russell, K. *Ann. Reports Med. Chem.* **1995**, *30*, 81. Lawson, K. *Pharmacol. Ther.* **1996**, *70*, 39.
4. Recent examples of tissue selective K_{ATP} modulators include: Grant, T.L.; Ohnmacht, C.J.; Howe, B.B. *Trends Pharmacol. Sci.* **1994**, *15*, 402. Atwal, K.S. *J. Cardiovasc. Pharmacol.* **1994**, *24*, S12.
5. Lumma Jr., W.C.; Anderson, P.S.; Baldwin, J.J.; Bolhofer, W.A.; Habecker, C.N.; Hirschfield, J.M.; Pietruszkiewicz, A.M.; Randall, W.C.; Torchiana, M.L.; Britcher, S.F.; Clineschmidt, B.V.; Denny, G.H.; Hirschmann, R.; Hoffman, J.M.; Phipps, B.T.; Streeter, K.B. *J. Med. Chem.* **1982**, *25*, 207. Algieri, A.A.; Luke, G.M.; Standridge, R.T.; Brown, M.; Partyka, R.A.; Crenshaw, R.R. *J. Med. Chem.* **1982**, *25*, 210.
6. Morita, T.; Yoshiizuma, K.; Nisihimura, N.; Goto, K.; Sukamoto, T.; Yoshino, K. European Patent Application 405 525 A2.
7. Amato, J.S.; Karady, S.; Reamer, R.A.; Schlegel, H.B.; Springer, J.P.; Weinstock, L.M. *J. Am. Chem. Soc.* **1982**, *104*, 1375. Lumma Jr., W.C.; Anderson, P.S.; Baldwin, J.J.; Bolhofer, W.A.; Habecker, C.N.; Hirschfeld, J.M.; Pietruszkiewicz, A.M.; Randall, W.C.; Torchiana, M.L.; Britcher, S.F.; Clineschmidt, B.V.; Denny, G.H.; Hirschman, R.; Hoffman, J.M.; Phillips, B.T.; Streeter, K.B. *J. Med. Chem.* **1982**, *25*, 207. Muller, D.; Umbricht, G.; Weber, B.; Pfaltz, A. *Helv. Chim. Acta.* **1991**, *74*, 232.
8. Karady, S.; Amato, J.S.; Dortmund, D.; Weinstock, L.M. *Heterocycles* **1981**, *16*, 1561. Karady, S.; Amato, J.S.; Dortmund, D.; Reamer, R.A.; Weinstock, L.M. *Heterocycles* **1981**, *16*, 1565.
9. All compounds exhibited satisfactory spectral and elemental analysis.
10. Gadwood, R. C.; Kamdar, B. V.; Cipkus-Dubray, L. A.; Wolfe, M. L.; Smith, M. P.; Watt, W.; Mizesak, S. A.; Groppi, V. E. *J. Med. Chem.* **1993**, *36*, 1480.

(Received in USA 27 November 1995; accepted 16 August 1996)