

Preliminary communication

SYNTHESIS AND STRUCTURAL CHARACTERIZATION OF $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2$ AND RELATED ALKYL AND ARYL THIOLATO COMPLEXES

CLIVE E. BRIANT, GLYN R. HUGHES, PETER C. MINSHALL and
 D. MICHAEL P. MINGOS*


*Inorganic Chemistry and Chemical Crystallography Laboratories, University of Oxford,
 South Parks Road, Oxford OX 3QR (Great Britain)*

(Received September 8th, 1980)

Summary


The synthesis and spectroscopic characterization of the complexes $\text{cis-Pt}(\text{SR})_2(\text{PPh}_3)_2$ ($\text{R} = \text{H, Me, n-Bu and Ph}$) are reported, together with the single crystal X-ray crystallographic characterization of $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2$. The ability of these complexes to act as bidentate ligands towards other metal centres is also briefly described.

In 1971 Ugo and his coworkers [1] reported that $\text{Pt}(\text{PPh}_3)_n$ ($n = 2$ or 3) reacts with H_2S to give an adduct $\text{Pt}(\text{PPh}_3)_2(\text{H}_2\text{S})$, which in solution appeared to exist as a mixture of the tautomeric forms I and II.



A weak interaction between H_B and the platinum d_{z^2} orbital was initially proposed [2] to account for the high field chemical shift of H_B , although this was later discarded in favour of diamagnetic ring current effect arising from the proximity of H_B to a phenyl ring [2]. Subsequently there have been reports of other SH complexes of the platinum metals, but to date none of the complexes have been structurally characterized in the solid state [3,4].

We have found that chloroform solution of the dioxygen complex, $\text{Pt}(\text{O}_2)(\text{PPh}_3)_2$, react readily with H_2S and RSH ($\text{R} = \text{Me}$, $n\text{-Bu}$ and Ph) according to the following equation



to give high yields of the *cis*-dithiolato complexes as off white ($\text{R} = \text{H}$) or pale yellow ($\text{R} = \text{Me}$, $n\text{-Bu}$ or Ph) crystalline complexes. Crystals of $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2 \cdot \text{CHCl}_3$ suitable for crystallographic investigation were obtained by recrystallisation from $\text{CHCl}_3/\text{EtOH}$.

Crystal data. The structure was determined from 5234 unique reflections having $I > 3\sigma(I)$ and refined to $R = 0.0450$. $\text{C}_{37}\text{H}_{33}\text{S}_2\text{Cl}_3\text{P}_2\text{Pt}$, $M = 904.5$, triclinic a 11.198(2), b 11.876(3) and c 14.462(4) Å, α 91.10(2), β 98.43(2) and γ 107.70(2)°, U 1808 Å³, $Z = 2$, D_c 1.66 g cm⁻³, $\mu(\text{Mo-K}\alpha)$ 45.4 cm⁻¹, $F(000) = 824$, $\text{Mo-K}\alpha$ 0.71069 Å, Space group $P\bar{1}$.

The molecular structure of $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2$ illustrated in Fig. 1 confirms that it has a *cis*-square planar structure with few significant distortions. The Pt, S(1), S(2), P(1) and P(2) atoms all lie within 0.06 Å of the best least squares planes through these atoms, and the deviations from 90° for S(1)—Pt—S(2) and P(1)—Pt—P(2) can be readily accounted for in terms of the very different steric requirements of SH and PPh_3 . The hydrogen atoms on the SH ligands appear to be crystallographically disordered and therefore were not unambiguously located.

The ^1H and $^{31}\text{P}\{^1\text{H}\}$ NMR data for the complexes $\text{cis-Pt}(\text{SR})_2(\text{PPh}_3)_2$ ($\text{R} = \text{H}$, Me or $n\text{-Bu}$) given in Table 1 confirm that they all have *cis*-geometries. The ^1H resonance associated with the SH ligands in $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2$ occurs at




Fig. 1. Molecular structure of $\text{cis-Pt}(\text{SH})_2(\text{PPh}_3)_2$, Pt—S(1) 2.360(2), Pt—S(2) 2.340(2), Pt—P(1) 2.286(2), Pt—P(2) 2.279(2) Å; S(1)—Pt—S(2) 83.14(8), S(1)—Pt—P(1) 89.28(7), S(2)—Pt—P(2) 90.06(8), P(1)—Pt—P(2) 97.65(7)°.

TABLE 1

NMR PARAMETERS^a FOR *cis*-Pt(SR)₂(PPh₃)₂ IN CD₂Cl₂ AT 293 K

Complex	δ (SH)	δ (SMe)	δ (³¹ P) ^b	² J(Pt—H)	³ J(P—H)	³ J(Pt—H)	⁴ J(Pt—H)	¹ J(Pt—P)
<i>cis</i> -Pt(SH) ₂ (PPh ₃) ₂	0.1		21.37	50	8.5			2981
<i>cis</i> -Pt(SMe) ₂ (PPh ₃) ₂		2.25	25.52			54	6.5	2862
<i>cis</i> -Pt(SBu) ₂ (PPh ₃) ₂			21.38					2863

^a δ in ppm, J in Hz. ^b To high frequency of trimethylphosphate.

0.18 (cf. -1.58 for II [1] and *trans*-Pt(SH)₂(PEt₃)₂ [3] and -1.18 for *trans*-PtH(SH)(PEt₃)₂ [3]).

The *cis*-arrangement of dithiolato ligands in the complexes described in this paper suggests that they may be able to function as bidentate ligands towards other transition metal centres in a manner analogous to that reported previously for M(η-C₅H₅)₂(SR)₂ (M = Ti or Nb; R = Me or Ph) [5,6].

cis-Pt(SPh)₂(PPh₃)₂ reacts readily with Mo(CO)₄(η⁴-C₇H₈) in CHCl₃ solutions to give (Ph₃P)₂Pt(μ-SPh)₂Mo(CO)₄ as a brick red crystalline solid (m.p. 134–137°C). The ν(CO) stretching frequencies for this mixed metal complex observed in CHCl₃ solutions at 2000, 1940, 1905, 1845 cm⁻¹, are consistent with its formulation as *cis*-bridged thiolato complex [5,6]. The complexes *cis*-Pt(SR)₂(PPh₃)₂ (R = n-Bu, or Ph) also react with the complexes Pd(PhCN)₂Cl₂ and Pt(COD)Cl₂ (COD = cyclooctadiene) to give the complexes (Ph₃P)₂Pt(SR)₂MCl₂ (M = Pd or Pt). The structural, spectroscopic and electrochemical properties of these complexes are currently being investigated. However, these preliminary results indicate that the complexes *cis*-Pt(SR)₂(PPh₃)₂ may prove to be important precursors for binuclear mixed metal thiolato complexes.

We thank the S.R.C. for financial support and Johnson–Mathey for the loan of platinum metals.

References

- 1 R. Ugo, G. La Monica, S. Cenini, A. Segre and F. Conti, J. Chem. Soc. A, (1971) 522.
- 2 D. Morelli, A. Segre, R. Ugo, G. La Monica, S. Cenini, F. Bonati and F. Conti, Chem. Commun., (1967) 524.
- 3 I.M. Blacklaws, E.A.V. Ebsworth, D.W.H. Rankin and H.E. Robertson, J. Chem. Soc. Dalton Trans., (1978) 753.
- 4 T. Miyamoto, J. Organometal. Chem., 134 (1977) 335.
- 5 P.S. Braterman, V.A. Wilson and K.K. Joshi, J. Chem. Soc. A, (1971) 191.
- 6 W.E. Douglas and M.L.H. Green, J. Chem. Soc. Dalton Trans., (1972) 1796.