

Tetrahedron Letters, Vol. 37, No. 5, pp. 699-702, 1996 Copyright © 1996 Elsevier Science Ltd Printed in Great Britain. All rights reserved 0040-4039/96 \$15.00 + 0.00

0040-4039(95)02246-5

Azobenzene-Porphyrins

Christopher A. Hunter* and Luke D. Sarson

Krebs Institute for Biomolecular Science, Department of Chemistry, University of Sheffield, Sheffield S3 7HF, UK and Department of Chemistry, University of Otago, PO Box 56, Dunedin, New Zealand

Abstract: A series of covalently-connected azobenzene-porphyrin derivatives have been synthesised and their photochemical properties investigated. The photochemistry of the porphyrin components is essentially unaltered, but photochemical isomerisation of the azobenzene components can not be detected.

The potential for molecules to function as components in information processing systems is attracting increasing attention.¹ In principle, well-designed molecules or supramolecular assemblies could perform switching and gating functions in place of conventional silicon circuitry.² Progress in this field requires the synthesis of compounds which can process electronic or optical inputs and generate a usable output. We have therefore investigated the properties of covalently-connected azobenzene-porphyrin conjugates, 1-3.³



Azobenzenes and porphyrins absorb in different regions of the visible spectrum, and hence the two chromophores should be independently addressable. In addition, 2 contains hydrogen-bonding sites which should confer molecular recognition properties on the system, and the long alkyl chains on 3 may lead to mesomorphism. Irradiation of the azobenzene chromophore should induce a photochemical isomerisation of the N-N double bond, which would modulate the photophysical, molecular recognition or phase properties of these compounds (Scheme 1), i.e. these are potential molecular logic gates.



Scheme 1 Schematic illustration of the proposed photochemical switching of the molecular recognition properties of M2. This would in turn affect the photochemical properties of the porphyrin chromophore.

Synthesis

The porphyrins were synthesised from the corresponding azobenzene aldehydes using the Lindsey condensation procedure (Scheme 2).⁴ This route allowed the synthesis of gram quantities of the azobenzene-porphyrins H_21a and H_21b .⁵ The simple model chromophores H_27 and 8 were synthesised using similar procedures (see Figure 1).



Scheme 2 Reagents and conditions: i, NaNO₂, HCl; ii. Me₂SO₄, NaH, DMF; iii, MeOH, H₂SO₄; iv, LiAlH₄, ether; v, DMSO, (COCl)₂, -78°C; vi, CH₂Cl₂, EtOH, BF₃•OEt₂; vii, DDQ, NEt₃

The subsequent derivatisation of H_21b is outlined in Scheme 3. Cleavage of both tetramethyl ethers, H_21a and H_21b , was attempted with boron tribromide.⁶ However, H_21a could not be cleanly demethylated due to the low solubility of the intermediate diol and triol. The use of isopropyl groups dramatically increased the solubility of the compounds so that even the fully demethylated tetraol H_22b was readily soluble in nonpolar organic solvents.



700

Photochemical Experiments

The UV/Visible absorption spectra of the azobenzene-porphyrins are almost a summation of the spectra of the component chromophores (Figure 1(a)), which suggests that there is little electronic communication between the azobenzene and the porphyrin components. Thus it is possible to address the two chromophores separately using different wavelength light (340 nm for the azobenzene and 430 nm for the porphyrin).



Figure 1 UV/Visible Absorption Spectra in CH_2Cl_2 (arbitrary absorption (A) scales) (a) The azobenzene-porphyrin H_2lb and the two model compounds H_27 and 8 (b) The change in the spectrum of 8 caused by photochemical isomerisation.

The fluorescence spectra of the azobenzene-porphyrins (on irradiation at the Soret maximum at 430 nm) are similar to that of the model porphyrin H_27 . Thus covalent connection of the two chromophores does not greatly perturb the excited state properties of the porphyrin component. However, the same is not true of the azobenzene components where there is a significant change in the excited state properties. The simple model azobenzene 8 was readily converted from the thermodynamically more stable (E) isomer to the (Z) isomer by irradiation with a mercury-lamp at 313 nm. In the absence of visible light, the (Z) isomer was stable for hours, and Figure 1(b) shows the change in the UV/Visible absorption spectrum caused by photochemical isomerisation. Similar experiments with the azobenzene-porphyrins produced no detectable change in the UV/Visible absorption spectrum. Thus in these systems, either the azobenzene excited state decays via a process which is faster than the isomerisation reaction, or the thermal relaxation of the (Z) to the

(E) isomer is so fast that the (Z) isomer was not detected on the timescale at which we were working (minutes). Porphyrins have been reported to act as intermolecular sensitisers of the (E) to (Z) isomerisation of azobenzenes, but that does not appear to be the case in this system.⁷ When excited at the azobenzene absorption maximum at 340 nm, the fluorescence spectra of the azobenzene-porphyrins are essentially identical to those of the model compounds H_27 and Zn7, so the mechanism of relaxation is neither direct fluorescent emission nor energy transfer to the porphyrin. This behaviour was not substantially affected by metallation of the porphyrins with zinc, coordination of ligands (pyridine or quinuclidine) to the metal or changes in solvent polarity (dichloromethane or hexane).

In conclusion, we have shown that the two different chromophores in covalently-connected azobenzene-porphyrins can be separately addressed by different wavelengths of light. While the excited state properties of the porphyrin components are not affected by connection to the azobenzenes, the excited state properties of the azobenzene components are significantly perturbed in these compounds: they no longer undergo photochemical isomerisation. More detailed studies of the photochemistry of these compounds are in progress.

Acknowledgements

We thank the University of Otago, University of Sheffield, Glaxo, the CVCP (LDS) and the Nuffield Foundation (CAH) for financial support. We thank Rob Hyde for some of the fluorescence studies.

References

- 1. Balzani, V. Supramolecular Photochemistry, D. Riedel, Dordrecht, 1991, 207.
- 2. Day, P. Chem. Br., 1990, 26, 52.
- Autret, M.; Le Plouzennec, M.; Moinet, C.; Simonneaux, G. J. Chem. Soc., Chem. Commun., 1994, 1169; Hombrecher, H. K.; Lüdtke, K. Tetrahedron, 1993, 49, 9489; Neumann, K. H.; Vögtle, F. J. Chem. Soc., Chem. Commun., 1988, 520.
- 4. Lindsey, J. S.; Wagner, R. W. J. Org. Chem., 1989, 54, 828.
- 5. Data for H₂1a: ¹H NMR (CDCl₃): δ (ppm) = 8.96 (8H, s, pyrrolic H); 8.38 (8H, d, ArH); 8.29 (8H, d, ArH); 8.11 (8H, d, ArH); 7.12 (8H, d, ArH); 3.96 (12H, s, OCH₃); -2.68 (2H, s, broad, pyrrolic NH). UV/VIS (CH₂Cl₂): λ_{max} = 346 nm (ε: 1.0 x 10⁵); 430 nm (ε: 5.1 x 10⁵); 518 nm (ε: 4.0 x 10⁴); 558 nm (ε: 3.2 x 10⁴); 592 nm (ε: 1.6 x 10⁴); 630 nm (ε: 1.6 x 10⁴); 690 nm (ε: 1.2 x 10⁴). MS (+ve FAB): 1152 (M⁺). Mp: >400°C. Data for H₂1b: ¹H NMR (CDCl₃): δ (ppm) = 8.98 (8H, s, pyrrolic H); 8.39 (8H, d, ArH); 8.30 (8H, d,

ArH); 7.90 (8H, d, ArH); 3.86 (12H, s, OCH₃); 3.46 (8H, sept, CH(CH₃)₂); 1.40 (48H, d, CHCH₃); -2.67 (2H, s, pyrrolic NH). UV/VIS (CH₂Cl₂): $\lambda_{max} = 338$ nm (ϵ : 9.7 x 10⁴); 430 nm (ϵ : 5.0 x 10⁵); 521 nm (ϵ : 3.3 x 10⁴); 558 nm (ϵ : 2.8 x 10⁴); 592 nm (ϵ : 1.3 x 10⁴); 650 nm (ϵ : 1.2 x 10⁴). MS (+ve FAB): 1486 (M⁺). Mp: >400°C.

- 6. Milgrom, L. R. J. Chem. Soc. Perkin Trans. I, 1983, 2535.
- 7. Wildes, P. D.; Pacifici, J. G.; Irick, G.; Whitten, D. G. J. Am. Chem. Soc., 1971, 93, 2004.

(Received in UK 26 October 1995; revised 23 November 1995; accepted 24 November 1995)