


Reaction of a moderately stabilized bismuthonium ylide with 1,2-dicarbonyl compounds. A novel ring enlargement of *ortho*-quinones to 3-hydroxytropones

Yoshihiro Matano and Hitomi Suzuki*

Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo-ku, Kyoto 606-01, Japan

Triphenylbismuthonium 2-oxoalkylide **2**, generated *in situ* from salt **1** in THF at low temperature, readily reacts with 1,2-dicarbonyl compounds **3** to give 2,3-diacyloxiranes **4** or 2-acyl-3-hydroxytropones **5** depending on the structure of the electrophiles involved.

Organic transformations based on bismuth reagents have proved to be of synthetic utility.¹ However, in spite of the well-established uses of phosphonium, arsonium and stibonium ylides as reagents for carbon–carbon bond formation,^{2–4} the bismuthonium ylide has received attention only recently.^{5–7} Moderately stabilized bismuthonium 2-oxoalkylides **2**, generated from the corresponding 2-oxoalkyl(triphenyl)bismuthonium salts **1**, readily react with aldehydes⁶ and activated imines⁷ to give the respective α,β -epoxy and α,β -aziridino ketones in good yields. This mode of reaction leading to oxirane and aziridine is characteristic of the bismuth, since the analogous ylides of phosphorus, arsenic and antimony all undergo Wittig-type olefination with the same substrates to give α,β -unsaturated ketones. Here we report that triphenylbismuthonium 2-oxoalkylides **2** can easily convert 1,2-dicarbonyl compounds to 2,3-diacyloxiranes **4** or 2-acyl-3-hydroxytropones **5**, depending on the structure of the substrates employed.



Scheme 1 Reagents and conditions: i, base, THF, $-78\text{ }^{\circ}\text{C}$ to room temp.

Triphenylbismuthonium 2-oxoalkylides **2a,b**, generated *in situ* from the corresponding bismuthonium salts **1a,b** and a base in THF at $-78\text{ }^{\circ}\text{C}$, did not react with simple ketones such as acetophenone and benzophenone or with an ester such as ethyl acetate. However, they readily underwent a carbon–carbon bond forming reaction with an α -keto ester, ethyl pyruvate **3a**, at low temperature to give the 2,3-difunctionalized oxiranes **4a,b** and triphenylbismuthane in good yields,[†] probably owing to the enhanced electrophilicity of the α -carbonyl carbon atom in **3a** through the adjacent electron withdrawing ethoxycarbonyl moiety. Potassium *tert*-butoxide (KO^tBu), potassium bis(trimethylsilyl)amide [$\text{KN}(\text{SiMe}_3)_2$] and sodium bis(trimethylsilyl)amide [$\text{NaN}(\text{SiMe}_3)_2$] were the bases used, and *trans* selectivity was always prominent in the oxiranes (Table 1, runs 1–5).[‡] When diacetyl **3b** was used as a 1,2-dicarbonyl counterpart, 2,3-diacyloxirane **4c** was obtained as the single stereoisomer (run 6). To our knowledge, there have been very few reported synthesis of 2,3-diacyloxiranes from 1,2-dicarbonyl compounds and an ylide bearing the 2-oxoalkylidene group.^{9,10} Therefore, the present epoxidation of 1,2-dicarbonyl compounds with bismuthonium 2-oxoalkylides **2** should provide a new addition to the existing synthesis of 2,3-difunctionalized oxiranes.

The reaction of bismuthonium 2-oxoalkylides **2a,b** with *ortho*-quinones **3c,d** led to a quite unexpected result. Treatment of the salts **1** with KO^tBu in the presence of 9,10-phenanthrenequinone **3c** afforded 2-acyl-3-hydroxytropones **5a,b** in good yield (runs 7–9).[§] A similar reaction of the salt **1b** with *ortho*-chloranil **3d** gave rise to a tropone **5c** in 33% yield (run 10). In the ^1H NMR spectra of tropones **5** in CDCl_3 , the hydroxy group was observed as a sharp singlet at δ_{H} 16.78–18.16, suggesting the operation of strong intramolecular hydrogen bonding in these molecules. The hydrogen bonding became weaker in a polar solvent such as $(\text{CD}_3)_2\text{SO}$; resonances of the hydroxy and dibenzo ring protons of **5a,b** turned into a broad singlet and a symmetrical multiplet pattern, respectively. Direct ring expansion of *ortho*-quinones to functionalized tropones based on the use of the bismuthonium ylides **2** may provide a new attractive route to some types of substituted tropones.¹²

Table 1 Reaction of 2-oxoalkyl(triphenyl)bismuthonium salts **1** with 1,2-dicarbonyl compounds **3** in the presence of a base

Entry	Salt	Base	Substrate	Product ^a	Yield (%) (<i>cis</i> : <i>trans</i>)
1	1a	KO^tBu	3a	4a	70 (12 : 88)
2	1a	$\text{KN}(\text{SiMe}_3)_2$	3a	4a	79 (10 : 90)
3	1a	$\text{NaN}(\text{SiMe}_3)_2$	3a	4a	82 (34 : 66)
4	1b	$\text{KN}(\text{SiMe}_3)_2$	3a	4b	62 (13 : 87)
5	1b	$\text{NaN}(\text{SiMe}_3)_2$	3a	4b	62 (9 : 91)
6	1b	$\text{NaN}(\text{SiMe}_3)_2$	3b	4c	26 (0 : 100)
7	1a	KO^tBu	3c	5a	71
8	1a	$\text{KN}(\text{SiMe}_3)_2$	3c	5a	75
9	1b	KO^tBu	3c	5b	66
10	1b	KO^tBu	3d	5c	33

^a All products gave satisfactory spectral and analytical data. In all reactions triphenylbismuthane was recovered in good yield.

Although we are not in a position to explain the mechanism of the novel ring enlargement, this coupling mode of the bismuthonium ylide **2** leading to compound **4** or **5** is in marked contrast with the well established behaviour of phosphonium ylides, which undergo the Wittig reaction with 1,2-diketones and *ortho*-quinones to yield α,β -unsaturated carbonyl compounds.^{2,13} It is noteworthy that two different types of carbonyl compounds **4** and **5** are accessible from the same bismuthonium ylides **2** via an appropriate modification of the structure of the 1,2-dicarbonyl compounds. Compounds like **4** and **5** have great synthetic potential, since their highly functionalized structures are readily subject to either ring opening or ring enlargement or new ring framework construction.

This work was supported by a Grant-in-Aid for Scientific Research (No. 08740492) from the Ministry of Education, Science, Sports and Culture, Japan.

Footnotes

† Typical procedure: to a stirred suspension of the salt **1a** (125 mg, 0.2 mmol) in THF (5 cm³) was added NaN(SiMe₃)₂ (1.0 mol dm⁻³ THF solution; 0.2 cm³, 0.2 mmol) at -78 °C. After 10 min, ethyl pyruvate **3a** (23 mg, 0.2 mmol) was added and the resulting mixture was gradually allowed to warm to room temp. Evaporation of the solvent under reduced pressure gave an oily residue, which was diluted with benzene (15 cm³) and the insoluble matter was filtered off through a Celite bed. The filtrate was concentrated under reduced pressure to leave an oily residue, the composition of which was estimated by ¹H NMR integration. The product was subjected to chromatographic separation on silica gel using hexane-ethyl acetate (100:0 to 80:20) as the solvent to elute triphenylbismuthane (85 mg, 97%), *cis*-**4a** (12 mg, 54%) and *trans*-**4a** (23 mg, 28%) in this order. Selected data for *cis*-**4a**: $\delta_{\text{H}}(\text{CDCl}_3)$ 1.22 (9 H, s), 1.32 (3 H, t, $J = 7.2$ Hz), 1.44 (3 H, s), 4.17 (1 H, s) and 4.27 (2 H, q, $J = 7.2$ Hz); m/z (EI) 141 ($M^+ - 73$); $\nu_{\text{max}}(\text{neat})$ 1744 (C=O) and 1717 (C=O) cm⁻¹. Found: C, 61.51; H, 8.42. C₁₁H₁₈O₄ requires C, 61.66; H, 8.47%. For *trans*-**4a**: $\delta_{\text{H}}(\text{CDCl}_3)$ 1.23 (9 H, s), 1.27 (3 H, t, $J = 7.2$ Hz), 1.67 (3 H, s), 3.79 (1 H, s) and 4.19 (2 H, q, $J = 7.2$ Hz); m/z (EI) 141 ($M^+ - 73$); $\nu_{\text{max}}(\text{neat})$ 1740 (C=O) and 1719 (C=O) cm⁻¹. Found: C, 61.65; H, 8.42. C₁₁H₁₈O₄ requires C, 61.66; H, 8.47%.

‡ The major product was assigned as the *trans* isomer by ¹H NMR spectral comparison with analogous compounds.^{8,9}

§ Selected data for **5a**: mp 154–155 °C; $\delta_{\text{H}}(\text{CDCl}_3)$ 1.40 (9 H, s), 7.43–7.65 (6 H, m), 7.67–7.75 (1 H, m), 8.15 (1 H, d, $J = 7.5$ Hz) and 18.16 (1 H, s); $\delta_{\text{H}}[(\text{CD}_3)_2\text{SO}]$ 1.20 (9 H, s), 7.62 (2 H, t, $J = 7.6$ Hz), 7.71 (2 H, t, $J = 7.4$ Hz), 7.75–7.90 (2 H, br s), 7.88 (2 H, d, $J = 7.8$ Hz) and 13.2 (1 H, br s); m/z (EI) 306 (M^+). Found: C, 78.61; H, 5.94. C₂₀H₁₈O₃ requires C, 78.41; H, 5.92%. For **5b**: mp 118–119 °C (lit.,¹¹ 117.5 °C); $\delta_{\text{H}}(\text{CDCl}_3)$ 7.40–7.82 (12 H, m), 8.27–8.35 (1 H, m) and 17.17 (1 H, s); $\delta_{\text{H}}[(\text{CD}_3)_2\text{SO}]$ 7.42–7.62 (5 H, m), 7.66 (2 H, t, $J = 7.3$ Hz), 7.76 (2 H, t, $J = 7.3$ Hz), 7.85 (2 H,

br d), 7.88 (2 H, d, $J = 7.9$ Hz) and 15.9 (1 H, br s); m/z (EI) 326 (M^+). For **5c**: mp 138–139 °C; $\delta_{\text{H}}(\text{CDCl}_3)$ 7.46 (2 H, t, $J = 7.7$ Hz), 7.52–7.60 (3 H, m) and 16.78 (1 H, s); m/z (EI) 364 (M^+). Found: C, 46.26; H, 1.63. C₁₄H₆Cl₄O₃ requires C, 46.19; H, 1.66%.

References

- 1 For reviews of organic transformations based on bismuth reagents, see: L. D. Freedman and G. O. Doak, *Chem. Rev.*, 1982, **82**, 15; J.-P. Finet, *Chem. Rev.*, 1989, **89**, 1487; M. Wada and H. Ohki, *J. Synth. Org. Chem., Jpn.*, 1989, **47**, 425; Y. Matano and H. Suzuki, *Bull. Chem. Soc. Jpn.*, 1996, **69**, 2673; H. Suzuki, T. Ikegami and Y. Matano, *Synthesis*, in the press.
- 2 A. W. Johnson, *Ylides and Imines of Phosphorus*, Wiley, Chichester, 1993.
- 3 D. Lloyd, I. Gosney and R. A. Ormiston, *Chem. Soc. Rev.*, 1987, **16**, 45.
- 4 Y.-Z. Huang, *Acc. Chem. Res.*, 1992, **25**, 182.
- 5 For the C–C bond formations based on stabilized bismuthonium ylides, see: H. Suzuki, T. Murafuji and T. Ogawa, *Chem. Lett.*, 1988, 847; T. Ogawa, T. Murafuji and H. Suzuki, *Chem. Lett.*, 1989, 849; T. Ogawa, T. Murafuji and H. Suzuki, *J. Chem. Soc., Chem. Commun.*, 1989, 1749; T. Ogawa, T. Murafuji, K. Iwata and H. Suzuki, *Chem. Lett.*, 1989, 325.
- 6 Y. Matano, *J. Chem. Soc., Perkin Trans. 1*, 1994, 2703.
- 7 Y. Matano, M. Yoshimune and H. Suzuki, *J. Org. Chem.*, 1995, **60**, 4663.
- 8 W. I. Awad, Z. S. Salih and Z. H. Aiube, *J. Chem. Soc., Perkin Trans. 1*, 1977, 1280.
- 9 Reaction between the ester **3a** and Me₂S=CHCO₂Et in boiling acetone has been reported to give diethyl 2-methyl-2,3-epoxysuccinate in 42% yield. G. B. Payne, *J. Org. Chem.*, 1968, **33**, 3517.
- 10 B. M. Trost and H. C. Arndt, *J. Org. Chem.*, 1973, **38**, 3140.
- 11 W. Ried and R. Conte, *Chem. Ber.*, 1971, **104**, 1573.
- 12 For the known ring expansion reactions of *ortho*-quinones to tropones and tropolones, see: B. Eistert and L. Klein, *Chem. Ber.*, 1968, **101**, 391; M. Sato, J. Tsunetsugu and S. Ebine, *Bull. Chem. Soc. Jpn.*, 1972, **45**, 638; H. Kogler, H.-W. Fehlhaber, K. Leube and W. Durckheimer, *Chem. Ber.*, 1989, **122**, 2205.
- 13 Triphenylphosphonium 2-oxoalkylides undergo the Wittig reaction with 9,10-phenanthroquinone to give α,β -unsaturated ketones as the initial product, see D. N. Nicolaides, S. G. Adamopoulos, D. A. Lefkaditis, K. E. Litinas and P. V. Tarantili, *J. Chem. Soc., Perkin Trans. 1*, 1992, 283; D. N. Nicolaides, S. G. Adamopoulos, D. A. Lefkaditis and K. E. Litinas, *J. Chem. Soc., Perkin Trans. 1*, 1990, 2127; D. N. Nicolaides, D. A. Lefkaditis, P. S. Lianis and K. E. Litinas, *J. Chem. Soc., Perkin Trans. 1*, 1989, 2329; H. J. Bestmann and H. J. Lang, *Tetrahedron Lett.*, 1969, 2101.

Received, 5th September 1996; Com. 6/06103F